AIRR:ARR-2026-0Y9HAB5A1R9KMARG v1 [Number theory] 27 Nov 2026 · AIRR.SCIENCE

Birational Quartic Descent, Infinite Duplication Towers, Elementary 2-Descent Rank-Zero Proofs, Group-Law Duality, and Constructive Fermat Descent for Pythapotent Hypotenuse–Leg Pairs

Lluis Eriksson
Independent Researcher
ORCID: 0009-0006-7494-9040

November 27, 2026

Abstract

Halbeisen, Hungerbühler, and Shamsi Zargar [4] recently introduced *pythapotent hypotenuse–leg pairs* (hl-pairs) of degree h: positive integer pairs (c,d) with $c^2-d^2=e^2>0$ admitting a Pythagorean pair (k,l) $\not\sim$ (d,e) $(k^2+l^2=\square)$ such that $(c^hk)^2-(d^hl)^2=\square>0$. Reducing degree-$h$ pythapotency to positive Mordell–Weil rank of the elliptic curve $\Gamma_{c^h,d^h}: y^2=x(x+c^{2h})(x-d^{2h})$, they proved universal pythapotency for $h\in\{1,3\}$ (Corollaries 8–9), gave a quartic sufficient family for $h=2$ (Corollary 10), and reported Magma rank computations on $(5,3)$ and $(5,4)$ (including $\operatorname{rank}(\Gamma_{5^2,4^2}(\mathbb{Q}))=0$). We establish four self-contained, elementary, and machine-checked structural theorems resolving all unproved algebraic assertions, Magma-only rank-zero barriers, group-law ambiguities, and external descent dependencies in [4], backed by 20 constructive Lean 4.36.0 theorems (zero external imports, zero `sorry`, zero `Classical.choice`, zero warnings), 16 AST-audited zero-float SHA-256 certificates, and 56 adversarial tests:

1. **Explicit Birational Isomorphism $\Psi: Q_1 \xrightarrow{\sim} E_{12}$, Degree-4 Morphism $\Phi=-[2]\circ\Psi$, Quartic Unification, & Infinite Duplication Tower ($h=2$):** Closing the unproved claim on p. 11 of [4], we exhibit the genuine birational isomorphism $\Psi(m,n,q)=\left(\frac{2(q+n^2)}{m^2},\frac{4n(q+n^2)}{m^3}\right)$ from $Q_1: 3m^4+n^4=q^2$ to $E_{12}: Y^2=X^3-12X$ with rational inverse $\Psi^{-1}(X,Y)=\left(\frac{m}{n}=\frac{2X}{Y},\frac{q}{n^2}=\frac{2X^3}{Y^2}-1\right)$ (and show the degree-4 descent morphism $\Phi(m,n,q)=\left(\frac{q^2}{m^2n^2},\frac{q(3m^4-n^4)}{m^3n^3}\right)$ factors as $\Phi=-[2]\circ\Psi$), unify all four Corollary 10 quartics via Remark 2's $(m,n)\mapsto(m+n,m-n)$ identity, and prove that the polynomial duplication endomorphism $(m',n',q')=(2mnq,|3m^4-n^4|,9m^8+18m^4n^4+n^8)$ preserves $\gcd(m,n)=1$, opposite parity, and $3\nmid n$ while strictly growing $q'>q$ and $c'>c$ (e.g. $(2,1,7)\mapsto(28,47,2593)\mapsto\cdots$), constructively generating an infinite primitive degree-2 pythapotent tower.
2. **4-Adic/8-Adic Corollary 10 Obstructions & Elementary Magma-Free 2-Descent Rank-Zero Theorem for $\Gamma_{5^2,4^2}$:** We prove mod-4 obstructions blocking opposite-parity Case 1 quartics and a mod-8 obstruction blocking every primitive Case 2 hl-pair with $b\equiv 4 \pmod 8$ (i.e. $4\mid b$ and $8\nmid b$, explaining why $(5,4)$ fails Corollary 10). Using elementary complete 2-descent, we prove that whenever $C=p^k$ ($p\equiv 5 \pmod 8$ prime), $A=2^s$ ($s\geq 3$), and $Q=C^2+A^2$ is prime (automatically $Q\equiv 1 \pmod 8$ and $(Q/p)=+1$), every non-torsion 2-descent homogeneous space fails locally at 2 or p, forcing $\dim_{\mathbb{F}_2}\mathrm{Sel}^{(2)}(\Gamma_{C,A})=2$ and $\operatorname{rank}(\Gamma_{C,A}(\mathbb{Q}))=0$—giving the first elementary, Magma-free proof that $(5,4)$ is not pythapotent of degree 2 ($C=25, A=16, Q=881$).
3. **Exact Lemma 5–6 Group-Law Identity ($R=+[2]P$), Crossed Norm Conservation, Symmetry Dichotomy, & Non-2-Divisibility Barrier:** We prove that the canonical algebraic composition of Lemmas 5 and 6 equals $+[2]P$ identically in both x- and y-coordinates (so positive-root Lemma 6 yields $\operatorname{sgn}(jwl)[2]P$), formalize Remark 1's

norm conservation law, contrast Corollary 8's $(d \leftrightarrow e)$-symmetry with Corollary 9's hypotenuse ordering $2c^2 - a^2 < 2c^2 - b^2 \iff b < a$, and prove that the universal points of Corollaries 8 (preimage Q), 9 ($P_0, P_0 + T_0, P_0 + T_1$), and 10 ($P$) never lie in $[2]\Gamma_{c^h,d^h}(\mathbb{Q})$.

4. **Constructive Fermat Infinite Descent, Elementary Lemma 2 ($h = 1$ and $2 \mid h$), & Torsion-3/4 Algebraic Boundary:** Proving both $x^4 + y^4 \neq z^2$ and $x^4 - y^4 \neq z^2$ constructively in Lean 4 from Euclidean gcd square-factoring, we prove Lemma 2 of [4] for $h = 1$ and all even $h \geq 2$ and $cd \neq \square$ without Darmon–Merel [1], certify the absence of rational 4-torsion constructively, and prove the exact 3-division identity $\psi_3(x) = 4x^2(x+C^2)(x-A^2)-(x^2+C^2A^2)^2$ ruling out rational 3-torsion via square-divisor elimination.

AI Assistance Disclosure. This research manuscript, its constructive Lean 4.36.0 formalization, and its exact-rational Python verification suite were developed by Lluis Eriksson with interactive assistance from Google Antigravity for proof exploration, formal verification, and typesetting. All mathematical proofs, formal Lean theorems, and exact-rational certificates were mechanically verified; the author takes full responsibility for all claims.

1 Introduction and Preliminaries

Extending their earlier studies of paired Pythagorean pairs [2, 3], Halbeisen, Hungerbühler, and Shamsi Zargar [4] defined a *hypotenuse–leg pair* (hl-pair) to be a pair $(c, d) \in \mathbb{Z}^2_{>0}$ with $c^2 - d^2 = e^2$ for some integer $e > 0$ (so (d, e, c) is a Pythagorean triple), and called (c, d) *pythapotent of degree* $h \geq 1$ if there exists a Pythagorean pair $(k, l) \in \mathbb{Z}^2_{>0}$ ($k^2 + l^2 = \square$) not a rational multiple of (d, e) such that $(c^h k, d^h l)$ is an hl-pair:

$$c^2 - d^2 = e^2 > 0, \qquad k^2 + l^2 = j^2, \qquad (c^h k)^2 - (d^h l)^2 = w^2 > 0. \tag{1}$$

To every hl-pair (c, d) and degree $h \geq 1$, [4] associates the elliptic curve

$$\Gamma_{C,A} \; : \; y^2 \; = \; x^3 + (C^2 - A^2)x^2 - C^2A^2x \; = \; x(x + C^2)(x - A^2), \qquad C := c^h, \;\; A := d^h. \tag{2}$$

Using Mazur's torsion theorem [5], Darmon–Merel [1], and Mordell's presentation of Euler's quartic [6], Proposition 3 and Theorem 4 of [4] establish $\Gamma_{c^h,d^h}(\mathbb{Q})_{\text{tors}} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and prove that (c, d) is pythapotent of degree h if and only if $\text{rank}(\Gamma_{c^h,d^h}(\mathbb{Q})) > 0$. Corollaries 8 and 9 show every hl-pair is pythapotent of degrees $h = 1$ and $h = 3$, while Corollary 10 establishes degree-2 pythapotency whenever $c^2 + \varepsilon cd + d^2 = q^2$ ($\varepsilon \in \{1, -1\}$), remarking without proof (p. 11) that the resulting quartics are birational to $Y^2 = X^3 - 12X$ of rank 1, and relying on Magma to assert $\text{rank}(\Gamma_{5^2,4^2}(\mathbb{Q})) = 0$ (p. 8).

Table 1 summarizes how our four main theorems close these algebraic, arithmetic, group-law, and constructive-descent gaps.

2 Birational Descent, Quartic Unification, and Infinite Duplication Tower for Corollary 10

Writing a primitive Pythagorean triple as $(a, b, c) = (m^2 - n^2, 2mn, m^2 + n^2)$ ($m > n > 0$, $\gcd(m, n) = 1$, $m \not\equiv n \pmod 2$), Corollary 10 of [4] reduces $c^2 + \varepsilon cd + d^2 = q^2$ to four quartics: $Q_1 : 3m^4 + n^4 = q^2$ and $Q_2 : m^4 + 3n^4 = q^2$ in Case 1 ($d = a$), and $Q_3^{\pm} : m^4 \pm 2m^3n + 6m^2n^2 \pm 2mn^3 + n^4 = q^2$ in Case 2 ($d = b$).

Theorem 2.1 (Birational Descent to E_{12}, Quartic Unification, and Constructive Duplication Tower)**.**

Table 1: Summary of results extending Halbeisen, Hungerbühler, and Shamsi Zargar [4].

Topic	Halbeisen et al. [4]	This Paper
Corollary 10 ($h = 2$) quartics	States birationality to E_{12} on p. 11 without explicit maps or proof	Explicit birational isomorphism $\Psi : Q_1 \xrightarrow{\sim} E_{12}$ (with rational Ψ^{-1}) and $\Phi = -[2] \circ \Psi$, Remark-2 unification of all 4 quartics, and invariant-preserving infinite duplication tower (Thm. 2.1)
Rank 0 of $\Gamma_{5^2,4^2}$	Reported via Magma computation (p. 8); no local obstruction analysis	Mod-4 / mod-8 obstructions for Cor. 10; Magma-free complete 2-descent proving $\mathrm{rank}(\Gamma_{C,A}(\mathbb{Q})) = 0$ for all $C = p^k$ ($p \equiv 5 \bmod 8$), $A = 2^s$ ($s \geq 3$), $Q = C^2 + A^2$ prime, including $\Gamma_{5^2,4^2}$ (Thm. 3.1)
Lemmas 5–6 & Cor. 8–10 points	Maps $P \mapsto (k, l) \mapsto R$ ($x(R) = x([2]P)$); y-sign and 2-divisibility unanalyzed	Exact algebraic identity $R = +[2]P$ (both coordinates); Remark-1 norm law; Cor. 8 symmetry vs. Cor. 9 asymmetry; non-2-divisibility barrier for Cor. 8, 9, 10 (Thm. 4.1)
Lemma 2 & Prop. 3 torsion	Invokes Darmon–Merel [1], Mazur [5], and Euler [6]	Constructive Lean 4 Fermat descent ($x^4 \pm y^4 \neq z^2$) proving Lemma 2 ($h = 1$, $2 \mid h$), $cd \neq \square$, no 4-torsion, and exact $\psi_3(x)$ identity (Thm. 5.1)

1. ***Explicit Birational Isomorphism*** $\Psi : Q_1 \xrightarrow{\sim} E_{12}$ ***& Degree-4 Morphism*** $\Phi = -[2] \circ \Psi$ (Lean `thm01`–`thm02`, Cert. `C01`)***:*** *For any nonzero $m, n, q \in \mathbb{Z}$ satisfying $3m^4 + n^4 = q^2$ (with $q + n^2 \neq 0$), the map*

$$\Psi(m, n, q) := \left(\frac{2(q + n^2)}{m^2}, \frac{4n(q + n^2)}{m^3} \right) \tag{3}$$

is a genuine birational isomorphism from $Q_1 : 3m^4 + n^4 = q^2$ onto $E_{12} : Y^2 = X^3 - 12X$, with rational inverse

$$\Psi^{-1}(X, Y) : \frac{m}{n} = \frac{2X}{Y}, \qquad \frac{q}{n^2} = \frac{2X^3}{Y^2} - 1. \tag{4}$$

Moreover, the degree-4 rational descent morphism

$$\Phi(m, n, q) := \left(\frac{q^2}{m^2 n^2}, \frac{q(3m^4 - n^4)}{m^3 n^3} \right) \tag{5}$$

maps $Q_1 \to E_{12}$ and factors identically as $\Phi = -[2] \circ \Psi$ (with squared-coordinate recovery $\frac{m^2}{n^2} = \frac{s^3+Y}{6s}$, $\frac{n^2}{m^2} = \frac{s^3-Y}{2s}$ for $s = q/(mn)$).

2. ***Remark-2 Quartic Unification*** (Lean `thm03`, Cert. `C02`)***:*** *Under the linear involution $(m, n) \mapsto (m + n, m - n)$,*

$$\begin{aligned} 3(m+n)^4 + (m-n)^4 &= 4(m^4 + 2m^3 n + 6m^2 n^2 + 2mn^3 + n^4), \\ (m+n)^4 + 3(m-n)^4 &= 4(m^4 - 2m^3 n + 6m^2 n^2 - 2mn^3 + n^4), \end{aligned} \tag{6}$$

establishing explicit rational equivalences $Q_3^+ \cong Q_1$ and $Q_3^- \cong Q_2 \cong Q_1$ (swapping $m \leftrightarrow n$).

3. ***Invariant-Preserving Infinite Duplication Tower*** (Lean `thm04`–`thm05`, Certs. `C01`, `C03`–`C04`)***:*** *For every positive integer solution (m, n, q) to $3m^4 + n^4 = q^2$ with $3m^4 \neq n^4$, define*

$$m' := 2mnq, \qquad n' := |3m^4 - n^4|, \qquad q' := 9m^8 + 18m^4 n^4 + n^8. \tag{7}$$

Then $3(m')^4+(n')^4=(q')^2$, $q'>q$, *and the triple of arithmetic invariants* ($\gcd(m,n)=1$, $m+n\equiv 1 \pmod 2$, $3\nmid n$) *is preserved identically at every step. Consequently, setting* $c_k:=m_k^2+n_k^2$, $a_k:=|m_k^2-n_k^2|$, *and* $\varepsilon_k:=\operatorname{sgn}(m_k-n_k)\in\{1,-1\}$ *yields* $c_k^2+\varepsilon_k c_k a_k+a_k^2=q_k^2$; *iterating from* $(m_0,n_0,q_0)=(2,1,7)\mapsto(28,47,2593)\mapsto\cdots$ *constructively generates infinitely many distinct primitive degree-2 pythapotent hl-pairs (beginning* $(5,3)$ *and* $(2993,1425)$*).*

Proof. (1) Set $M:=m^4$, $N:=n^4$, and $S:=q+n^2$. Since $q^2=3M+N$, we have $S^2-3M=q^2+2n^2q+N-(q^2-N)=2n^2S$, whence $(2S)^3-12(2S)M=8S(S^2-3M)=16n^2S^2=(4nS)^2$ (Lean `thm01`); dividing by m^6 verifies $\Psi(m,n,q)\in E_{12}$. For the rational inverse, $\frac{2X}{Y}=\frac{4S/m^2}{4nS/m^3}=\frac{m}{n}$ and $\frac{2X^3}{Y^2}-1=\frac{16S^3/m^6}{16n^2S^2/m^6}-1=\frac{S-n^2}{n^2}=\frac{q}{n^2}$ (Lean `thm02`), confirming $\Psi^{-1}\circ\Psi=\mathrm{id}$. Conversely, for any $(X,Y)\in E_{12}$ with $Y\neq 0$ (so $Y^2=X^3-12X$), setting $t:=2X/Y$ and $u:=2X^3/Y^2-1=(X^3+12X)/Y^2$ satisfies $3t^4+1=(48X^4+(X^3-12X)^2)/Y^4=(X^3+12X)^2/Y^4=u^2$ (hence $\Psi^{-1}(E_{12})\subset Q_1$), and $2(u+1)/t^2=(4X^3/Y^2)/(4X^2/Y^2)=X$, $4(u+1)/t^3=(8X^3/Y^2)/(8X^3/Y^3)=Y$, verifying $\Psi\circ\Psi^{-1}=\mathrm{id}$ with zero square roots. For Φ, $(3M-N)^2=(3M+N)^2-12MN=q^4-12MN$ gives $q^2(3M-N)^2=q^6-12q^2MN$ (Lean `thm01`), and direct evaluation of the duplication formula on E_{12} at $(X,Y)=\Psi(m,n,q)$ gives $x([2]\Psi)=\frac{(3X^2-12)^2}{4Y^2}-2X=\frac{q^2}{m^2n^2}$ and $y([2]\Psi)=-\frac{q(3m^4-n^4)}{m^3n^3}$, so $\Phi=-[2]\circ\Psi$ (Cert. `C01`), with $q^3\pm q(3M-N)=6qM,2qN$ (Lean `thm02`).

(2) Expanding $(m\pm n)^4=m^4\pm 4m^3n+6m^2n^2\pm 4mn^3+n^4$ and forming $3(m+n)^4+(m-n)^4$ and $(m+n)^4+3(m-n)^4$ gives (6) directly (Lean `thm03`, Cert. `C02`).

(3) Setting $X:=m^4, Y:=n^4$ so $q^2=3X+Y$, we have $3(m')^4+(n')^4=48XY(3X+Y)^2+(3X-Y)^4=81X^4+324X^3Y+342X^2Y^2+36XY^3+Y^4=(9X^2+18XY+Y^2)^2=(q')^2$, and $q'=(3X+Y)^2+12XY=q^4+12XY>q$ (Lean `thm04`). Moreover, since $c_k^2=(m_k^2+n_k^2)^2\in[\frac{1}{3}q_k^2,2q_k^2]$ (as $m_k^4+n_k^4\le c_k^2\le 2(m_k^4+n_k^4)$ and $\frac{1}{3}q_k^2\le m_k^4+n_k^4\le q_k^2$) while $q_{k+1}>q_k^4$, we have $c_{k+1}^2\ge\frac{1}{3}q_{k+1}^2>\frac{1}{3}q_k^8>2q_k^2\ge c_k^2$ (for all $q_k\ge 7$), so the hypotenuses c_k are strictly increasing and all generated primitive hl-pairs (c_k,a_k) are pairwise distinct. For invariant preservation: since $m'=2mnq$ is even while $n'=|3m^4-n^4|\equiv 3m^4-n^4\equiv m+n\equiv 1 \pmod 2$, opposite parity is preserved; since $n'=|3m^4-n^4|\equiv\pm n^4\equiv\pm 1\not\equiv 0 \pmod 3$ whenever $3\nmid n$, $3\nmid n'$ is preserved (Lean `thm04`). For coprimality, let p be a prime dividing $\gcd(m',n')$. Since n' is odd and $3\nmid n'$, $p\notin\{2,3\}$. Thus $p\mid mnq$. If $p\mid m$ or $p\mid n$, then $p\mid(3m^4-n^4)$ forces p to divide both m and n, contradicting $\gcd(m,n)=1$. If $p\mid q$, then $p\mid(3m^4+n^4)$ and $p\mid(3m^4-n^4)$ force $p\mid 6m^4$ and $p\mid 2n^4$; as $p\notin\{2,3\}$, $p\mid\gcd(m,n)=1$, contradiction. Hence $\gcd(m',n')=1$. Finally, $(m^2+n^2)^2+(m^2+n^2)(m^2-n^2)+(m^2-n^2)^2=3m^4+n^4$ (Lean `thm05`), so replacing m^2-n^2 by $\varepsilon|m^2-n^2|$ ($\varepsilon=\operatorname{sgn}(m-n)$) gives $c_k^2+\varepsilon_k c_k a_k+a_k^2=q_k^2$, satisfying Corollary 10 of [4] (Certs. `C01`, `C03`). □

3 Modular Obstructions and Elementary 2-Descent Rank-Zero Theorem for $\Gamma_{C,A}$

Halbeisen et al. [4] (p. 8) used Magma to report $\operatorname{rank}(\Gamma_{5^2,4^2}(\mathbb{Q}))=0$, proving $(5,4)$ is not pythapotent of degree 2. We first isolate the elementary congruential obstructions explaining why Corollary 10 fails on $(5,4)$ and half of all Case 1 parameters, and then prove an infinite family of elementary, Magma-free complete 2-descent rank-zero theorems covering $\Gamma_{5^2,4^2}$.

Theorem 3.1 (Modular Obstructions and Elementary 2-Descent Rank-Zero Family)**.**

1. ***Corollary 10 Congruential Obstructions*** (Lean `thm06`–`thm07`, Cert. `C05`)***:***
   - *If m is odd and n is even, then $3m^4+n^4\equiv 3 \pmod 4$, so Q_1 has no integer solution; symmetrically, if m is even and n is odd, then $m^4+3n^4\equiv 3 \pmod 4$, so Q_2 has no integer solution.*



Figure 1: (a) Quadruple-exponential height growth ($\log_{10} q_k$ and $\log_{10} c_k$) of the invariant-preserving duplication tower (7) seeded at $(2, 1, 7)$ (Theorem 2.1). (b) Genuine birational image $\Psi(2, 1, 7) = (4, 4)$ and degree-4 morphism images $\Phi(2, 1, 7) = -[2]\Psi(2, 1, 7) = (\frac{49}{4}, \frac{329}{8})$ and $\Phi(28, 47, 2593)$ on $E_{12} : Y^2 = X^3 - 12X$.

- *For every primitive Case-2 hl-pair (c, b) with $b \equiv 4 \pmod 8$ (i.e. $4 \mid b$ and $8 \nmid b$, equivalently $2 \mid mn$ and $4 \nmid mn$, such as $(5, 4)$), $c^2 \pm cb + b^2 \equiv 5 \pmod 8$, so Corollary 10 fails for both $\varepsilon = \pm 1$.*

2. ***Elementary Complete 2-Descent Rank-Zero Theorem*** (Lean `thm08`–`thm10`, Certs. `C06`–`C08`)**:** *Let $p \equiv 5 \pmod 8$ be prime, $k \geq 1$, $s \geq 3$, $C := p^k$, $A := 2^s$, and suppose $Q := C^2 + A^2$ is prime. Then automatically $Q \equiv 1 \pmod 8$ and $(Q/p) = +1$, and among all 16 real-solvable 2-descent homogeneous spaces of $\Gamma_{C,A} : y^2 = x(x + C^2)(x - A^2)$, all 12 non-torsion cosets are locally unsolvable at 2 or p:*
   - *every coset with $\delta \in \{\pm 2, \pm 2p\}$ fails modulo p (since $(2/p) = -1$ and $(Q/p) = +1$);*
   - *every coset with $\delta \in \{\pm p, \pm 2p\}$ fails 2-adically across all three coprime valuation regimes ($2 \nmid u$, $2 \parallel u$, and $4 \mid u, 2 \nmid v$).*

   Hence $|\mathrm{Sel}^{(2)}(\Gamma_{C,A})| = 4 = |\Gamma_{C,A}(\mathbb{Q})[2]|$, forcing $\mathrm{rank}(\Gamma_{C,A}(\mathbb{Q})) = 0$.

3. ***Magma-Free Proof for*** $(5, 4)$ ***at*** $h = 2$ (Lean `thm10`, Cert. `C06`)**:** *Specializing Part (2) to $(p, k, s) = (5, 2, 4)$ gives $C = 25$, $A = 16$, and $Q = 25^2 + 16^2 = 881$ (prime, $881 \equiv 1 \pmod 8$, $881 \equiv 1 \pmod 5$), proving unconditionally and without Magma that $\mathrm{rank}(\Gamma_{5^2,4^2}(\mathbb{Q})) = 0$, so $(5, 4)$ is not pythapotent of degree 2.*

Proof. (1) Every odd integer satisfies $x^4 \equiv 1 \pmod 4$ and every even integer satisfies $x^4 \equiv 0 \pmod 4$, whereas squares are 0 or 1 (mod 4) (Lean `thm06`). For c odd and $b \equiv 4 \pmod 8$, we have $c^2 \equiv 1$, $cb \equiv 4$, and $b^2 \equiv 0 \pmod 8$, giving $c^2 \pm cb + b^2 \equiv 5 \pmod 8$, impossible since squares mod 8 lie in $\{0, 1, 4\}$ (Lean `thm07`).

(2) Note that $C = p^k$ is odd and $8 \mid A = 2^s$ ($s \geq 3$), so $Q = C^2 + A^2 \equiv 1 + 0 = 1 \pmod 8$ and $Q \equiv (2^s)^2 \pmod p$, whence the Legendre symbol $(Q/p) = +1$ holds automatically (Cert. `C08`). For complete 2-descent on $\Gamma_{C,A} : y^2 = x(x + C^2)(x - A^2)$ with $\gcd(C, A) = 1$ (Silverman–Tate [7]), any rational point $P = (x, y)$ ($y \neq 0$) has x with square denominator (since any prime q with negative odd $v_q(x)$ would force odd $v_q(y^2) = 3v_q(x)$), so writing $x = \delta u^2/v^2$ with $\gcd(u, v) = 1$, $\gcd(\delta, v) = 1$ ($v > 0$) and squarefree (δ, d_1, d_2) ($\delta d_1 d_2 = \square$) gives the homogeneous space

$$\delta u^2 + C^2 v^2 \;=\; d_1 w_1^2, \qquad \delta u^2 - A^2 v^2 \;=\; d_2 w_2^2. \tag{8}$$

Because $\gcd(u, v) = \gcd(\delta, v) = 1$, the pairwise gcd bounds $\gcd(\delta u^2, \delta u^2 + C^2 v^2) \mid C^2 = p^{2k}$, $\gcd(\delta u^2, \delta u^2 - A^2 v^2) \mid A^2 = 2^{2s}$, and $\gcd(\delta u^2 + C^2 v^2, \delta u^2 - A^2 v^2) \mid Q$ (with Q prime, $Q \nmid 2p$)

restrict the prime support of the squarefree triples:

- 2 divides neither d_1 (if $2 \mid u$ then $2 \nmid v$ and C is odd, so $\delta u^2 + C^2 v^2$ is odd) nor Q, hence $v_2(\delta) = v_2(d_2) \in \{0, 1\}$ and $v_2(d_1) = 0$;
- p divides neither d_2 (if $p \mid u$ then $p \nmid v$ and $p \nmid A$, so $\delta u^2 - A^2 v^2$ is coprime to p) nor Q, hence $v_p(\delta) = v_p(d_1) \in \{0, 1\}$ and $v_p(d_2) = 0$;
- Q cannot divide δ, so $v_Q(d_1) = v_Q(d_2) \in \{0, 1\}$.

By real solvability of (8): if $\delta > 0$, then $\delta u^2 + C^2 v^2 > 0$ forces $d_1 > 0$, whence $d_2 > 0$ (since $\delta d_1 d_2 = \square > 0$); if $\delta < 0$, then $\delta u^2 - A^2 v^2 < 0$ forces $d_2 < 0$, whence $d_1 > 0$. Thus there are exactly $2 \times 2^3 = 16$ real-solvable triples (δ, d_1, d_2): 4 come from the 2-torsion subgroup $\Gamma_{C,A}(\mathbb{Q})[2]$—namely $\mathcal{O} \mapsto (1, 1, 1)$, $(0, 0) \mapsto (-1, 1, -1)$, $(A^2, 0) \mapsto (1, Q, Q)$, and $(-C^2, 0) \mapsto (-1, Q, -Q)$—and 12 are non-torsion ($\delta \in \{\pm 2, \pm p, \pm 2p\}$, $v_Q \in \{0, 1\}$).

To check $\mathbb{Q}_\ell$-solvability ($\ell \in \{2, p\}$) of the projective curve (8) in $(u : v : w_1 : w_2)$: first, the hyperplane at infinity $v = 0$ requires $\delta/d_1, \delta/d_2 \in (\mathbb{Q}_\ell^\times)^2$, which fails for all 12 non-torsion cosets (for $\delta = \pm 2$, $(\delta/d_1 \bmod p) = (\pm 2/p) = -1$; for $\delta = \pm p$, $\delta/d_2 = pQ^{-v_Q} \equiv 5 \pmod 8 \notin (\mathbb{Q}_2^\times)^2$; for $\delta = \pm 2p$, $v_2(\delta/d_1) = 1$ is odd). Second, any $\mathbb{Q}_\ell$-point with $v \neq 0$ scales homogeneously to $(u, v) \in \mathbb{Z}_\ell^2$ with $\min(v_\ell(u), v_\ell(v)) = 0$, which forces $w_1, w_2 \in \mathbb{Z}_\ell$ (since d_1, d_2 are squarefree, so $v_\ell(d_i w_i^2) \geq 0 \implies v_\ell(w_i) \geq 0$). We now eliminate all 12 non-torsion triples in $\mathbb{Z}_\ell$:

(a) *Mod-p elimination of $\delta = \pm 2$ (4 cosets):* Here $d_1 \in \{1, Q\}$ satisfies $(d_1/p) = +1$, while $d_2 = \operatorname{sgn}(\delta) \cdot 2Q^{v_Q}$ satisfies $(d_2/p) = (\pm 2/p) = -1$ (since $p \equiv 5 \pmod 8$, so $(-1/p) = +1$ and $(2/p) = -1$). If $p \nmid u$, reducing the first equation $\pm 2u^2 + C^2 v^2 = d_1 w_1^2$ modulo p gives $\pm 2u^2 \equiv d_1 w_1^2 \pmod p$, contradicting $(\pm 2u^2/p) = -1$ versus $(d_1 w_1^2/p) \in \{0, +1\}$ (Lean `thm08`). If $p \mid u$, then $p \nmid v$ (as $\gcd(u, v) = 1$), so reducing the second equation $\pm 2u^2 - A^2 v^2 = d_2 w_2^2$ modulo p gives $-A^2 v^2 \equiv d_2 w_2^2 \pmod p$, contradicting $(-A^2 v^2/p) = (-1/p) = +1$ (as $p \nmid A$) versus $(d_2 w_2^2/p) \in \{0, -1\}$. Thus both $p \nmid u$ and $p \mid u$ fail locally at p.

(b) *2-adic elimination of $\delta = \pm p$ (4 cosets):* Here $d_1 \in \{p, pQ\} \equiv 5 \pmod 8$ and $d_2 = \operatorname{sgn}(\delta) \cdot Q^{v_Q} \equiv \operatorname{sgn}(\delta) \pmod 8$. Multiplying the second equation by $\operatorname{sgn}(\delta)$ yields $pu^2 - \operatorname{sgn}(\delta) A^2 v^2 = |d_2| w_2^2$ with $|d_2| \equiv 1 \pmod 8$ and $64 \mid A^2$. We split into the three exhaustive coprime 2-adic valuation cases (Lean `thm09`):

- $2 \nmid u$: $pu^2 \equiv 5 \pmod 8$, forcing $|d_2| w_2^2 \equiv 5 \pmod 8$, impossible since $|d_2| \equiv 1 \pmod 8$ and $w_2^2 \bmod 8 \in \{0, 1, 4\}$.
- $2 \parallel u$: writing $u = 2u_1$ ($2 \nmid u_1$), $4 \mid (pu^2 - \operatorname{sgn}(\delta) A^2 v^2)$ forces $2 \mid w_2$ ($w_2 = 2w_2'$). Dividing by 4 gives $pu_1^2 - \operatorname{sgn}(\delta)(A/2)^2 v^2 = |d_2|(w_2')^2$; since $4 \mid (A/2)$ (as $8 \mid A$), this reduces mod 8 to $5 \equiv |d_2|(w_2')^2 \pmod 8$, impossible.
- $4 \mid u$: by $\gcd(u, v) = 1$, v is odd, so $8 \mid pu^2$ and $C^2 v^2 \equiv 1 \pmod 8$. The first equation then gives $d_1 w_1^2 \equiv 1 \pmod 8$, impossible since $d_1 \equiv 5 \pmod 8$ implies $d_1 w_1^2 \bmod 8 \in \{0, 4, 5\}$.

(c) *2-adic elimination of $\delta = \pm 2p$ (4 cosets):* Here $d_1 \in \{p, pQ\} \equiv 5 \pmod 8$ and $d_2 = \operatorname{sgn}(\delta) \cdot 2Q^{v_Q}$ ($|d_2|/2 \equiv 1 \pmod 8$). If $2 \nmid u$, the second equation $\pm 2pu^2 - A^2 v^2 = d_2 w_2^2$ divided by 2 gives $pu^2 - \operatorname{sgn}(\delta)(A^2/2) v^2 = (|d_2|/2) w_2^2 \implies 5 \equiv (|d_2|/2) w_2^2 \pmod 8$ (as $32 \mid A^2/2$), impossible. If $2 \mid u$, then v is odd and $8 \mid 2pu^2$, so the first equation gives $d_1 w_1^2 \equiv C^2 v^2 \equiv 1 \pmod 8$, impossible since $d_1 \equiv 5 \pmod 8$.

Thus only the 4 torsion cosets survive local solvability, proving $|\mathrm{Sel}^{(2)}(\Gamma_{C,A})| = 4$ and $\operatorname{rank}(\Gamma_{C,A}(\mathbb{Q})) = 0$.

(3) Immediate from Part (2) with $p = 5$, $k = 2$, $s = 4$, $C = 25$, $A = 16$, $Q = 881$ (Lean `thm10`, Cert. `C06`). □



Figure 2: (a) Complete 16-coset 2-descent elimination matrix on $\Gamma_{5^2,4^2}$: all 12 non-torsion cosets fail locally at 2 or 5, proving rank = 0 without Magma (Theorem 3.1). (b) Certified rank-zero instances of the generalized family $C = p^k$ ($p \equiv 5 \pmod 8$), $A = 2^s$ ($s \geq 3$), $Q = C^2 + A^2$ prime.

4 Exact Lemma 5–6 Group-Law Identity, Crossed Duality, and Non-2-Divisibility Barrier

Theorem 4 of [4] links rational points on $\Gamma_{C,A}$ to Pythagorean pairs (k, l) via Lemma 5 ($P \mapsto (k, l)$) and Lemma 6 ($(k, l) \mapsto R$), observing that $x(R) = x([2]P)$ without establishing the y-coordinate identity or checking whether the universal points of Corollaries 8–10 are divisible by 2 in $\Gamma_{c^h,d^h}(\mathbb{Q})$.

Theorem 4.1 (Group-Law Identity, Crossed Norm Conservation, Symmetry Dichotomy, and Non-2-Divisibility)**.**

1. ***Exact Lemma 5–6 Group-Law Identity*** (Lean `thm11`, Cert. `C09`)*: Let* $P = (x_1, y_1) \in \Gamma_{C,A}(\mathbb{Q})$ *with* $y_1 \neq 0$*, and define the signed polynomials*

$$k := x_1^2 + C^2A^2, \quad l := 2Cy_1, \quad j := x_1^2 + 2C^2x_1 - C^2A^2, \quad w := C(x_1^2 - 2A^2x_1 - C^2A^2).$$

   Then $k^2 + l^2 = j^2$, $(Ck)^2 - (Al)^2 = w^2$*, and the Lemma 6 point* $R := (\frac{C^2k^2}{l^2}, \frac{C^2kjw}{l^3})$ *equals* $+[2]P$ *identically in both coordinates (when positive roots* $|j|, |w|, |l|$ *are taken in Lemma 6,* $R_+ = \mathrm{sgn}(jwl)\,[2]P$*).*

2. ***Crossed Remark-1 Norm Conservation & Corollary 8–9 Symmetry Dichotomy*** (Lean `thm12`–`thm13`, Certs. `C10`–`C11`)*: Under Remark 1's duality* $(l', k') := (Ck, Al)$*, setting* $(\hat{k}, \hat{l}) := (Ck', Al') = (CAl, CAk)$ *satisfies* $(Ck')^2 + (Al')^2 = \hat{k}^2 + \hat{l}^2 = C^2A^2(k^2 + l^2)$*, while* $(Ck)^2 - (Al)^2 = w^2$ *implies* $(l')^2 - (k')^2 = w^2$*. Moreover, Corollary 8's universal pair* $(d^4 + d^2e^2 + e^4, 2c^2de)$ *is strictly* $(d \leftrightarrow e)$*-symmetric (identical for Case 1 and Case 2), whereas Corollary 9's pair* $(d^2, 2ce)$ *has hypotenuse* $2c^2 - d^2$*, strictly breaking symmetry via* $2c^2 - a^2 < 2c^2 - b^2 \iff b < a$*.*

3. ***Non-2-Divisibility Barrier for Corollaries 8, 9, and 10*** (Lean `thm14`–`thm15`, Cert. `C12`)*: For every hl-pair* (c, d)*:*

   - *Corollary 8's preimage* $Q = (-cd, -cde) \in \Gamma_{c,d}(\mathbb{Q})$ *does not lie in* $[2]\Gamma_{c,d}(\mathbb{Q})$*;*
   - *Corollary 9's universal point* $P_0 = (c^4d^2, c^4d^2e(c^2+d^2))$ *and its torsion translates* $P_0+(0,0)$ *and* $P_0 + (-c^6, 0)$ *do not lie in* $[2]\Gamma_{c^3,d^3}(\mathbb{Q})$*;*
   - *Corollary 10's point* $P = (-\varepsilon cd^3, cd^3eq)$ *does not lie in* $[2]\Gamma_{c^2,d^2}(\mathbb{Q})$ *for either* $\varepsilon \in \{1, -1\}$*.*



Figure 3: (a) Corollary 9 symmetry breaking: Case 1 (c, a) vs. Case 2 (c, b) hypotenuse $2c^2 - d^2$ across primitive Pythagorean triples (Theorem 4.1). (b) The 3-division polynomial $\psi_3(x) = 4x^2(x + C^2)(x - A^2) - (x^2 + C^2A^2)^2$ on $\Gamma_{5,3}$, strictly negative at all three 2-torsion points (Theorem 5.1).

Proof. (1) Expanding $j^2 - k^2 = 4C^2(x_1^3 + (C^2 - A^2)x_1^2 - C^2A^2x_1) = 4C^2y_1^2 = l^2$ and $(Ck)^2 - (Al)^2 = C^2(k^2 - 4A^2y_1^2) = w^2$ (Lean `thm11`) verifies the pair identities. For the elliptic duplication formula on $y^2 = x^3 + (C^2 - A^2)x^2 - C^2A^2x$, the tangent slope is $\lambda = \frac{3x_1^2+2(C^2-A^2)x_1-C^2A^2}{2y_1}$, giving $x([2]P) = \lambda^2 - (C^2 - A^2) - 2x_1 = \frac{(x_1^2+C^2A^2)^2}{4y_1^2} = \frac{C^2k^2}{l^2}$ and

$$\begin{aligned} y([2]P) &= \lambda(x_1 - x([2]P)) - y_1 \\ &= \frac{(x_1^2 + C^2A^2)(x_1^2 + 2C^2x_1 - C^2A^2)(x_1^2 - 2A^2x_1 - C^2A^2)}{8y_1^3} = \frac{C^2kjw}{l^3}, \end{aligned}$$

proving $R = +[2]P$ identically (Cert. `C09`).

(2) Both algebraic norm identities and $(d^2)^2 + (2ce)^2 = (2c^2 - d^2)^2$ follow directly from $d^2 + e^2 = c^2$ (Lean `thm12`–`thm13`, Certs. `C10`–`C11`).

(3) By the duplication identities (2) of [4], every point $S = (x_S, y_S) \in [2]\Gamma_{C,A}(\mathbb{Q})$ must have x_S, $x_S + C^2$, and $x_S - A^2$ all rational squares (in particular $x_S \geq 0$).

- For Corollary 8's $Q = (-cd, -cde)$: $x(Q) = -cd < 0$ is not a square in $\mathbb{Q}$ (Lean `thm15`).
- For Corollary 9's $P_0 = (c^4d^2, \dots)$ on Γ_{c^3,d^3}: $x(P_0) + c^6 = c^4(c^2 + d^2)$ being a rational square forces $c^2 + d^2 = \square$ (Lean `thm14`), contradicting Theorem 5.1(2). For the torsion translates: $x(P_0 + (0,0)) = -c^6d^6/(c^4d^2) = -c^2d^4 < 0$, and $x(P_0 + (-c^6, 0)) = -c^6 + \frac{c^6(c^6+d^6)}{c^4(c^2+d^2)} = -c^2d^2(c^2 - d^2) = -c^2d^2e^2 < 0$, neither a square.
- For Corollary 10's $P = (-\varepsilon cd^3, \dots)$ on Γ_{c^2,d^2}: if $\varepsilon = +1$, $x(P) = -cd^3 < 0$; if $\varepsilon = -1$, $x(P) = cd^3 = (cd)d^2$ being a rational square forces $cd = \square$ (Lean `thm15`), contradicting Theorem 5.1(2). □

5 Constructive Fermat Descent, Elementary Lemma 2, and Torsion-3/4 Algebraic Boundary

Lemma 2 and Proposition 3 of [4] cite Darmon–Merel [1], Mazur [5], and Mordell [6] even in the elementary regimes $h = 1$ and $2 \mid h$. We prove Fermat's two quartic descents completely constructively inside Lean 4.36.0 (using only Euclidean gcd coprime square-factoring and primitive Pythagorean parametrization, with zero external libraries), and derive self-contained proofs of

Lemma 2 ($h = 1,\ 2 \mid h$), $cd \neq \square$, the absence of 4-torsion, and the exact 3-division algebraic boundary.

Theorem 5.1 (Constructive Fermat Descent, Elementary Lemma 2, and Torsion-3/4 Certification)**.**

1. ***Constructive Fermat Infinite Descent*** (Lean `thm16`–`thm17`, Cert. `C13`)*: Neither* $x^4 + y^4 = z^2$ *nor* $x^4 - y^4 = z^2$ *has any solution in positive integers* $x, y, z \in \mathbb{Z}_{>0}$.
2. ***Elementary Lemma 2 (*$h = 1$ *and* $2 \mid h$*) & Non-Square Product*** (Lean `thm18`, Cert. `C14`)*: For every hl-pair* (c, d) *and every* $h \in \{1\} \cup 2\mathbb{Z}_{>0}$, $c^{2h} + d^{2h} \neq \square$, *and* $cd \neq \square$ *(with no reliance on Darmon–Merel [1]).*
3. ***Constructive 4-Torsion Elimination & 3-Division Polynomial Identity*** (Lean `thm19`–`thm20`, Certs. `C15`–`C16`)*: On* $\Gamma_{C,A}$ *(*$C, A > 0$*):*
   - *No rational point has order 4 whenever* $C^2 + A^2 \neq \square$ *(in particular, for every hl-pair and* $h \in \{1\} \cup 2\mathbb{Z}_{>0}$*).*
   - *The 3-division polynomial* $\psi_3(x) := 3x^4 + 4(C^2 - A^2)x^3 - 6C^2A^2x^2 - C^4A^4$ *satisfies the exact identity*
   $$\psi_3(x) \;=\; 4x^2(x + C^2)(x - A^2) - (x^2 + C^2A^2)^2. \tag{9}$$
   Consequently, any rational point $P = (x, y)$ *of order* 3 *on* $\Gamma_{C,A}$ *satisfies* $4xy^2 = (x^2 + C^2A^2)^2$*, forcing* $x = t^2 > 0$ *to be an integer square with* $t \mid C^2A^2$ *(ruling out 3-torsion by finite square-divisor evaluation on any concrete curve, and reducing generally to Euler's* $r^4 + r^2s^2 + s^4 = \square$ *via Remark 3 of [4]).*

Proof. (1) First, if $\gcd(a, b) = 1$ and $ab = n^2$ $(a, b, n \in \mathbb{N})$, setting $g_1 := \gcd(a, n)$ gives $g_1^2 \mid a$ (since $g_1^2 \mid n^2 = ab$ and $\gcd(g_1^2, b) = 1$) and $a \mid g_1^2$ (writing $a = g_1a', n = g_1n'$ with $\gcd(a', n') = 1$ forces $a' \mid g_1$), proving $a = g_1^2$ and symmetrically $b = g_2^2$ with zero prime factorization. Applying this to primitive Pythagorean triples $a^2 + b^2 = c^2$ $(2 \mid b)$ via $\frac{c+a}{2} \cdot \frac{c-a}{2} = (b/2)^2$ yields coprime $m > n > 0$ with $a = m^2 - n^2$, $b = 2mn$, $c = m^2 + n^2$.

- For $x^4+y^4 = z^2$ (strong induction on z, Lean `thm16`): stripping $\gcd(x, y)$ reduces to primitive $(x^2)^2+(y^2)^2 = z^2$ (WLOG $2 \mid y$). Parametrizing gives $x^2+n^2 = m^2$, $y^2 = 2mn$, $z = m^2+n^2$; parametrizing $x^2 + n^2 = m^2$ gives $n = 2rs$, $m = r^2 + s^2$ with m, r, s pairwise coprime and $m(rs) = (y/2)^2$. Hence $m = (z')^2$, $r = (x')^2$, $s = (y')^2$, yielding $(x')^4 + (y')^4 = (z')^2$ with $z' \leq m < z$.
- For $y^4 + z^2 = x^4$ (strong induction on x, Lean `thm17`): stripping $\gcd(x, y)$ reduces to $\gcd(y^2, z) = 1$. If $2 \mid z$, parametrizing gives $y^2 = m^2 - n^2$, $x^2 = m^2 + n^2$, yielding $n^4 + (xy)^2 = m^4$ with $m < x$. If $2 \nmid z$ (so $2 \mid y$), parametrizing gives $y^2 = 2mn$, $x^2 = m^2+n^2$, so $\{m, n\} = \{s^2, 2r^2\}$ and $(s^2)^2+(2r^2)^2 = x^2$. Parametrizing a second time gives $s^2 = a^2-b^2$, $r^2 = ab$, $x = a^2 + b^2$, whence $a = u^2, b = v^2$, and $v^4 + s^2 = u^4$ with $u < x$.

(2) If $c^2 - d^2 = e^2$ and $c^2 + d^2 = z^2$, multiplying gives $c^4 - d^4 = (ez)^2$, contradicting Part (1). For even $h = 2k$, $c^{2h} + d^{2h} = (c^k)^4 + (d^k)^4 \neq \square$ directly by Part (1). If $cd = z^2$, letting $g = \gcd(c, d)$ and $c = gc_1, d = gd_1, e = ge_1$ yields $c_1d_1 = (z/g)^2$ with $\gcd(c_1, d_1) = 1$, hence $c_1 = u^2, d_1 = v^2$, and $v^4 + e_1^2 = u^4$, again contradicting Part (1) (Lean `thm18`, Cert. `C14`).

(3) If $P = (x_1, y_1)$ has order 4, then $[2]P \in \{(0, 0), (-C^2, 0), (A^2, 0)\}$ with $x([2]P) = \frac{(x_1^2+C^2A^2)^2}{4y_1^2} > 0$, ruling out $(0, 0)$ and $(-C^2, 0)$ immediately (Lean `thm19` certifies $(x_1^2+C^2A^2)^2 > 0$, $-C^2 < 0$, and the $A^2 + C^2 = C^2 + A^2$ reduction); at $(A^2, 0)$, the duplication identity (2) of [4] forces $x([2]P) + C^2 = A^2 + C^2 = \square$, impossible by Part (2). Finally, expanding $4x^2(x + C^2)(x - A^2) - (x^2 + C^2A^2)^2$ gives (9) identically (Lean `thm20`, Cert. `C15`). Since $x(x + C^2)(x - A^2) = y^2$ on $\Gamma_{C,A}$, $\psi_3(x) = 0 \iff 4xy^2 = (x^2 + C^2A^2)^2$, forcing $x > 0$ to be a rational square. By the rational root theorem on $3x^4 + \cdots - C^4A^4 = 0$, the denominator of x divides 3; since 3 is not a square, $x = t^2 \in \mathbb{Z}_{>0}$ with $t \mid C^2A^2$ (verified with zero roots on $\Gamma_{5,3}, \Gamma_{5,4}, \Gamma_{13,5}, \Gamma_{25,16}$ in Cert. `C16`). □

6 Formal Verification Architecture and Reproducibility

All algebraic identities, modular and 2-adic obstructions, Euclidean square-factoring, primitive Pythagorean parametrization, Fermat quartic infinite descents, and torsion reductions are verified across three layers (while Mordell–Weil complete 2-descent homomorphisms and rational-root divisibility arguments are proved in full in Sections 2–5):

1. **Constructive Lean 4.36.0 Formalization:** 20 theorems (`thm01`–`thm20`) in `lean/PythagoreanHLPairsBarriers.lean` compiling under `lean -DwarningAsError=true` with zero external imports, zero `sorry`, zero `grind`, zero `Classical.choice`, and zero warnings.
2. **AST-Audited Zero-Float Python Verifier:** Pure standard-library code (`int`, `Fraction`) in `verification/verify_pythagorean_hl_pairs.py` self-auditing its AST (0 float violations) and emitting 16 canonical SHA-256 certificates (`C01`–`C16`) across four modes, with manifest SHA-256 `640d215f71c572c37cec8cf0722e7a8e1494ca0ea51cec6f6bb48ee3df278ac1`.
3. **Adversarial Test Suite:** 56 pytest checks in `tests/test_adversarial.py` covering positive verification and adversarial boundary cases across all four theorems.

References

[1] Henri Darmon and Loïc Merel. Winding quotients and some variants of Fermat's last theorem. *Journal für die Reine und Angewandte Mathematik*, 490:81–100, 1997.

[2] Lorenz Halbeisen and Norbert Hungerbühler. Pairing pythagorean pairs. *Journal of Number Theory*, 233:467–480, 2022.

[3] Lorenz Halbeisen, Norbert Hungerbühler, and Arman Shamsi Zargar. Pairing powers of pythagorean pairs. *Indagationes Mathematicae*, 36:903–911, 2025.

[4] Lorenz Halbeisen, Norbert Hungerbühler, and Arman Shamsi Zargar. Pairing powers of Pythagorean pairs II. *arXiv preprint arXiv:2611.34902v1*, November 2026.

[5] Barry Mazur. Modular curves and the Eisenstein ideal. *Publications Mathématiques de l'IHÉS*, 47:33–186, 1977.

[6] Louis J. Mordell. *Diophantine Equations*. Academic Press, London, 1969.

[7] Joseph H. Silverman and John Tate. *Rational Points on Elliptic Curves*. Springer-Verlag, New York, 2nd edition, 2015.