EXACT ALL-COUPLING NON-PERTURBATIVE BOUNDS FOR SU(2) AND SU(N) LATTICE YANG–MILLS THEORY WITH HEAT-KERNEL (MENOTTI–ONOFRI) AND MANTON GEODESIC ACTIONS

LLUÍS ERIKSSON

Abstract. Beyond Wilson's cosine plaquette action, the Menotti–Onofri heat-kernel (Villain) action and Manton's geodesic-distance-squared action are the two canonical lattice discretizations of non-Abelian Yang–Mills theory on compact Lie groups. Without invoking polymer or cluster expansions, we establish four self-contained non-perturbative theorems on periodic hypercubic tori $\mathbb{T}_L^d$ ($d \geq 2$):

(1) *Exact 2D Heat-Kernel, Casimir scaling, and zero-quadrature Manton thermodynamics:* Dual character and Jacobi–Poisson series evaluate $K_t^{\mathrm{SU}(2)}(\theta)$ to arbitrary precision across all $t > 0$, proving exact all-coupling 2D Casimir scaling $\sigma_R^{\mathrm{2D,HK}}/\sigma_F^{\mathrm{2D,HK}} = C_2(R)/C_2(F)$ for all SU(N) ($N \geq 2$), while a purely positive backward moment recurrence $M_j(\beta) = (e^{-a} + 2aM_{j+1}(\beta))/(2j+1)$ evaluates exact 2D SU(2) Manton free energies and string tensions with zero quadrature.

(2) *Sharp pointwise Wilson–Manton domination and universal $\mathbb{Z}_2$ confinement:* Pointwise $1-\cos\theta \leq \frac{1}{2}\theta^2 \leq \frac{\pi^2}{4}(1-\cos\theta)$ on SU(2) sandwiches the d-dimensional Manton free energy $f^{\mathrm{W},0}_{\mathbb{T}^d_L}(\frac{\pi^2}{4}\beta) \leq f^{\mathrm{M},0}_{\mathbb{T}^d_L}(\beta) \leq f^{\mathrm{W},0}_{\mathbb{T}^d_L}(\beta)$ at every finite L, d, and $\beta \geq 0$; duplicate-variable Ginibre $\mathbb{Z}_2$ domination plus Weighted-Kraft FIFO link queues certify $\underline{\sigma}_{1/2} > 0$ for both Heat-Kernel ($\tau_{\mathrm{HK}}(t) < \rho_d^4$) and Manton ($\tanh(\frac{\pi^2}{4}\beta) < \rho_d^4$) SU(2) in $d \geq 3$.

(3) *All-coupling d-dimensional Heat-Kernel semigroup enclosures matching exact one-loop entropy:* Transfer-matrix positivity proves exact dimensional monotonicity $f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) \geq f^{\mathrm{HK}}_{\mathbb{T}^{d-1}_L}(t) \geq \cdots \geq f^{\mathrm{HK}}_{\mathbb{T}^2_L}(t) > 0$ at every finite $L \geq 2$, while d-way Hölder's inequality, axial gauge, and $\int_G K_t^G(U)^2\,dU = K_{2t}^G(I)$ prove $\overline{f}^{\mathrm{HK}}_{\mathrm{3D}}(t) \leq \ln K_{2t}^G(I)$ and $\overline{f}^{\mathrm{HK}}_{\mathrm{4D}}(t) \leq 3\ln K_{2t}^G(I)$ for *every* compact connected Lie group G. On SU(2), a Schur-averaged geodesic-ball Gibbs variational bound on $\underline{f}^{\mathrm{HK}}_{-d}(t)$ matches the exact one-loop entropy coefficient $\binom{d-1}{2}\frac{3}{2}\ln(1/t)$ with bounded $O(1)$ gaps ([3.95, 3.99] in 3D over $t \in [1/160, 1/20]$; [6.20, 6.30] in 4D over $t \in [1/80, 1/10]$).

(4) *Exact Heat-Kernel Bakry–Émery curvature:* Exploiting $\mathrm{Ric}_{\mathrm{SU}(2)} = 1 \cdot g_{\mathrm{HS}}$, exact SU(2) Dynkin indices $T(\mathbf{m}) = \frac{m(m^2-1)}{6}$, and non-negativity of $(K_t'/K_t)^2 \geq 0$ certifies $\kappa_{\mathrm{HK}}(d,t) = 1 - 8(d-1)A_2(t)/K_t^{\mathrm{lb}}(t) > 0$, volume-uniform LSI ($\rho_{\mathrm{LS}} \geq \kappa_{\mathrm{HK}}$), and spectral gap $\lambda_1 \geq \kappa_{\mathrm{HK}}$ up to $\beta_{\mathrm{HK}} = 2/t = 4/11 \approx 0.3636$ in 4D (4.36× deeper than Wilson SU(2)'s $\beta < 1/12$).

All 24 benchmarks are certified at true 140-digit `mpmath.iv` precision (`mp.mp.dps = mp.iv.dps = 140`) and mechanically verified in a standalone zero-import, zero-axiom Lean 4.36.0 kernel (`HeatKernelMantonCerts.lean`).

1. Introduction and Honest Scope

On a periodic hypercubic torus $\mathbb{T}_L^d$ ($d \geq 2$, $V = L^d$, $|E| = dV$, $|P| = \binom{d}{2}V$, unoriented elementary plaquettes $p = (x; \mu < \nu)$), let G be a compact connected Lie group with normalized Haar measure dU and bi-invariant Hilbert–Schmidt metric $\langle X, Y\rangle_{\mathrm{HS}} = \mathrm{Re}\,\mathrm{Tr}(X^\dagger Y)$ (for $G = \mathrm{SU}(2)$, every $U \in \mathrm{SU}(2)$ has eigenvalues $e^{\pm i\theta}$ with unique Cartan angle $\theta \in [0,\pi]$ and geodesic distance $d_{\mathrm{geo}}(I_2, U) = \sqrt{2}\,\theta$). Alongside the 0-normalized Wilson plaquette action $s_W(U) := 1 - \frac{1}{2}\mathrm{Re}\,\mathrm{Tr}(U) = 1 - \cos\theta$ [9], we analyze the two canonical geometric alternatives:

Date: November 29, 2026.

(1) *Manton geodesic-distance-squared action* [5]: $s_M(U) := \frac{1}{4} d_{\mathrm{geo}}(I_2, U)^2 = \frac{1}{2}\theta^2$ on SU(2) (matching $s_W(U) = \frac{1}{2}\theta^2 + O(\theta^4)$ near I_2), with partition function and free energy

$$Z^{\mathrm{M},0}_{\mathbb{T}^d_L}(\beta) := \int_{\mathrm{SU}(2)^{|E|}} e^{-\beta \sum_{p\in P} s_M(U_p)}\, dU, \qquad f^{\mathrm{M},0}_{\mathbb{T}^d_L}(\beta) := \frac{1}{V} \ln Z^{\mathrm{M},0}_{\mathbb{T}^d_L}(\beta) \;\le\; 0,$$

and analogously $f^{\mathrm{W},0}_{\mathbb{T}^d_L}(\beta)$ for s_W.

(2) *Menotti–Onofri heat-kernel (Villain) action* [3, 6]: for diffusion time $t > 0$ (matching Wilson coupling $\beta_{\mathrm{HK}} := 2/t$ on SU(2)), define the normalized heat kernel on G (with $\int_G K^G_t(U)\, dU = 1$ and $K^G_t * K^G_s = K^G_{t+s}$)

$$K^G_t(U) := \sum_{\lambda\in\widehat{G}} d_\lambda\, \chi_\lambda(U)\, e^{-\frac{t}{2}C_2(\lambda)}, \qquad K^{\mathrm{SU}(2)}_t(\theta) = \sum_{m=1}^{\infty} m\, \frac{\sin(m\theta)}{\sin\theta}\, e^{-\frac{t}{4}(m^2-1)}, \tag{1}$$

where on SU(2) the spin-j irrep ($m = 2j+1$) has Casimir $C_2(j) = \frac{m^2-1}{2}$ (so $\frac{t}{2}C_2(j) = \frac{t}{4}(m^2-1) = tj(j+1)$). Define the unnormalized and 0-normalized heat-kernel free energies

$$f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) := \frac{1}{V} \ln \int_{G^{|E|}} \prod_{p\in P} K^G_t(U_p)\, dU, \qquad f^{\mathrm{HK},0}_{\mathbb{T}^d_L}(t) := f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) - \binom{d}{2} \ln K^G_t(I). \tag{2}$$

Remark 1.1 (Honest Scope and Non-Claims). All results hold at fixed lattice spacing ($a = 1$). Theorems 2.1, 3.1(1), and 4.1 hold at *all* couplings ($\beta > 0$, $t > 0$), whereas Theorem 3.1(2) holds in the strong-coupling slab $\tau < \rho_d^4$ and Theorem 5.1 holds for t sufficiently large that $8(d-1)A_2(t) < K^{\mathrm{lb}}_t(t)$ ($t \ge 5.5$, $\beta_{\mathrm{HK}} \le 4/11$ in 4D), certifying a volume-uniform log-Sobolev inequality and spectral gap rather than a 4D continuum limit. This paper does *not* claim any resolution of the Clay Millennium Yang–Mills Existence and Mass Gap Problem.

2. Exact 2D Heat-Kernel, Casimir Scaling, and Manton Thermodynamics

For $\beta > 0$, define the single-plaquette SU(2) Manton integrals $c^{\mathrm{M}}_0(\beta) := \frac{2}{\pi}\int_0^\pi \sin^2\theta\, e^{-\frac{\beta}{2}\theta^2}\, d\theta$ and $c^{\mathrm{M}}_1(\beta) := \frac{2}{\pi}\int_0^\pi \sin\theta\sin(2\theta)\, e^{-\frac{\beta}{2}\theta^2}\, d\theta$, and the positive Gaussian moments $M_j(\beta) := \int_0^1 u^{2j} e^{-au^2}\, du$ ($a := \frac{\beta\pi^2}{2}$, $j \in \mathbb{N}$).

Theorem 2.1 (Jacobi–Poisson Duality, Exact 2D Casimir Scaling, and Manton Backward Recurrence)**.**

(1) (Jacobi–Poisson identity on SU(2)) *For every $t > 0$ and $\theta \in (0, \pi)$ (with $\theta = 0$ and $\theta = \pi$ obtained by L'Hôpital limits $\theta \to 0^+$ and $\theta \to \pi^-$, pairing n with $-1-n$ at $\theta \to \pi^-$),*

$$\begin{aligned} K^{\mathrm{SU}(2)}_t(\theta) &= e^{t/4}\frac{2\sqrt{\pi}}{t^{3/2}} \sum_{n\in\mathbb{Z}} \frac{\theta + 2\pi n}{\sin\theta}\, e^{-\frac{(\theta+2\pi n)^2}{t}}, \\ K^{\mathrm{SU}(2)}_t(0) &= e^{t/4}\frac{2\sqrt{\pi}}{t^{3/2}} \sum_{n\in\mathbb{Z}} \left(1 - \frac{8\pi^2 n^2}{t}\right) e^{-\frac{4\pi^2 n^2}{t}}. \end{aligned} \tag{3}$$

(2) (Exact 2D Heat-Kernel thermodynamics and Casimir scaling for all SU(N)) *For every $N \ge 2$, $L \ge 2$, and $t > 0$, on $\mathbb{T}^2_L$ the exact Migdal–Rusakov partition function is $Z^{\mathrm{HK}}_{\mathbb{T}^2_L}(t) = \sum_{\lambda\in\widehat{\mathrm{SU}(N)}} e^{-\frac{t}{2}C_2(\lambda)L^2}$, whence $f^{\mathrm{HK}}_{\mathrm{2D}}(t) = 0$ and $f^{\mathrm{HK},0}_{\mathrm{2D}}(t) = -\ln K^{\mathrm{SU}(N)}_t(I)$ in the thermodynamic limit. On open/infinite 2D planes, every irreducible representation R has exact string tension $\sigma^{\mathrm{2D,HK}}_R(t) = \frac{t}{2}C_2(R)$, yielding exact all-coupling Casimir scaling $\sigma^{\mathrm{2D,HK}}_R(t)/\sigma^{\mathrm{2D,HK}}_F(t) = C_2(R)/C_2(F)$ (for* SU(2), $\sigma^{\mathrm{2D,HK}}_{1/2} = \frac{3}{4}t = \frac{3}{2\beta_{\mathrm{HK}}}$ *and* $\sigma_1/\sigma_{1/2} = \frac{8}{3}$ *identically).*

(3) (Exact zero-quadrature 2D SU(2) Manton thermodynamics) *For every* $\beta > 0$, $f_{\mathrm{2D}}^{\mathrm{M},0}(\beta) = \ln c_0^{\mathrm{M}}(\beta)$ *and* $\sigma_{1/2}^{\mathrm{2D,M}}(\beta) = -\ln \frac{c_1^{\mathrm{M}}(\beta)}{2c_0^{\mathrm{M}}(\beta)} > 0$, *where* $M_j(\beta)$ *satisfies the strictly positive backward recurrence*

$$M_j(\beta) = \frac{e^{-a} + 2a\, M_{j+1}(\beta)}{2j+1} \qquad \Big(0 < M_j(\beta) \le \frac{1}{2j+1},\ a = \frac{\beta\pi^2}{2}\Big), \tag{4}$$

and the exact β*-independent-radius alternating series*

$$c_0^{\mathrm{M}}(\beta) = \sum_{j=1}^{\infty} \frac{(-1)^{j-1}(2\pi)^{2j}}{(2j)!}\, M_j(\beta), \qquad c_1^{\mathrm{M}}(\beta) = \sum_{j=1}^{\infty} \frac{(-1)^{j-1}\big((3\pi)^{2j} - \pi^{2j}\big)}{(2j)!}\, M_j(\beta). \tag{5}$$

Proof. (1) Apply the Poisson summation formula to the even function $g(y) = y\sin(y\theta)e^{-ty^2/4}$ on $\mathbb{Z}$: its Fourier transform is $\widehat{g}(\xi) = +\frac{d}{d\xi}\Big[\sqrt{\frac{4\pi}{t}}\,\frac{e^{-(\xi-\theta)^2/t} - e^{-(\xi+\theta)^2/t}}{2}\Big]$, yielding $\sum_{m=1}^{\infty} m\sin(m\theta)e^{-tm^2/4} = \frac{1}{2}\sum_{n\in\mathbb{Z}}\widehat{g}(2\pi n) = \frac{2\sqrt{\pi}}{t^{3/2}}\sum_{n\in\mathbb{Z}}(\theta + 2\pi n)e^{-(\theta+2\pi n)^2/t}$. Multiplying by $e^{t/4}/\sin\theta$ proves (3) on $(0,\pi)$; letting $\theta \to 0^+$ via L'Hôpital's rule gives the $\theta = 0$ identity, and pairing n with $-1-n$ yields the $\theta \to \pi^-$ limit.

(2) By orthogonality of irreducible characters in 2D [7, 8], gluing L^2 plaquettes on the genus-1 torus $\mathbb{T}_L^2$ yields $Z_{\mathbb{T}_L^2}^{\mathrm{HK}}(t) = \sum_\lambda d_\lambda^0 (e^{-\frac{t}{2}C_2(\lambda)})^{L^2}$ (since $\chi(\mathbb{T}_L^2) = 0$), which satisfies $1 \le Z_{\mathbb{T}_L^2}^{\mathrm{HK}}(t) \le \sum_\lambda e^{-\frac{t}{2}C_2(\lambda)L^2} = 1 + O(e^{-\frac{t}{2}C_2(F)L^2})$, giving $f_{\mathrm{2D}}^{\mathrm{HK}}(t) = 0$. On an infinite plane, an $R \times T$ Wilson loop in representation R has expectation $(e^{-\frac{t}{2}C_2(R)})^{RT}$, proving exact Casimir scaling.

(3) On $\mathbb{T}_L^2$, character orthogonality yields $Z_{\mathbb{T}_L^2}^{\mathrm{M},0}(\beta) = \sum_{m=1}^{\infty}(c_{m-1}^{\mathrm{M}}(\beta)/m)^{L^2}$ with $c_{m-1}^{\mathrm{M}}(\beta) := \frac{2}{\pi}\int_0^\pi \sin(m\theta)\sin\theta\, e^{-\frac{\beta}{2}\theta^2}\, d\theta$. Since $|\sin(m\theta)| < m\sin\theta$ on $(0,\pi)$ for every $m \ge 2$, strict domination $|c_{m-1}^{\mathrm{M}}(\beta)| < m c_0^{\mathrm{M}}(\beta)$ holds. Integrating by parts twice against $\sin(m\theta)$ (the first boundary term vanishes as $\sin\theta\, e^{-\frac{\beta}{2}\theta^2} = 0$ at $\theta = 0, \pi$) gives $|c_{m-1}^{\mathrm{M}}(\beta)| \le C(\beta)m^{-2}$, whence the ratios $r_m(\beta) := |c_{m-1}^{\mathrm{M}}(\beta)/(m c_0^{\mathrm{M}}(\beta))|$ obey $r_m(\beta) \le \min(r(\beta), C'(\beta)m^{-3})$ with $r(\beta) < 1$; thus $\sum_{m\ge 2} r_m(\beta)^{L^2} \to 0$ as $L \to \infty$ and $f_{\mathrm{2D}}^{\mathrm{M},0}(\beta) = \ln c_0^{\mathrm{M}}(\beta)$. Substituting $\theta = \pi u$ in $c_0^{\mathrm{M}}(\beta) = \int_0^1 (1 - \cos(2\pi u))e^{-au^2}\, du$ and $c_1^{\mathrm{M}}(\beta) = \int_0^1 (\cos(\pi u) - \cos(3\pi u))e^{-au^2}\, du$ and expanding the cosines in Maclaurin series (uniformly convergent on $[0,1]$) gives (5); integrating $M_j(\beta) = \int_0^1 u^{2j}e^{-au^2}\, du$ by parts once ($\frac{u^{2j+1}}{2j+1}e^{-au^2}\big|_0^1 + \frac{2a}{2j+1}\int_0^1 u^{2j+2}e^{-au^2}\, du$) proves (4). Positivity of $c_1^{\mathrm{M}}(\beta) = \frac{4}{\pi}\int_0^{\pi/2}\sin^2\theta\cos\theta(e^{-\frac{\beta}{2}\theta^2} - e^{-\frac{\beta}{2}(\pi-\theta)^2})\, d\theta > 0$ and $|\cos\theta| < 1$ a.e. give $\sigma_{1/2}^{\mathrm{2D,M}}(\beta) > 0$. □

3. Sharp Action Domination and Universal $\mathbb{Z}_2$ Confinement

Recall the odd-subset Weighted-Kraft polynomials $F_d(\rho) := \frac{\rho}{2}[(1+\rho)^{2d-3} - (1-\rho)^{2d-3}]$ (with $F_3(0.55) = 0.99900625 \le 1$ and $F_4(0.39) = 0.995362843761 \le 1$) and $F_{C,d}(\rho) := \frac{1}{2}[(1+\rho)^{2(d-1)} - (1-\rho)^{2(d-1)}]$. Define the elementary character lower bound

$$K_t^{\mathrm{lb}}(t) := 1 - \sum_{m=2}^{\infty} m^2\, e^{-\frac{t}{4}(m^2-1)}. \tag{6}$$

Theorem 3.1 (Sharp Wilson–Manton Domination and $\mathbb{Z}_2$ Confinement for Heat-Kernel & Manton SU(2))**.**

(1) (Exact finite-L Wilson–Manton free-energy sandwich) *For every* $U \in \mathrm{SU}(2)$, $s_W(U) \le s_M(U) \le \frac{\pi^2}{4}s_W(U)$ *(lower bound sharp at* $U = +I_2$*; upper bound sharp at* $U = -I_2$*).*

Consequently, for every $d \geq 2$, every $L \geq 2$, and every $\beta \geq 0$,

$$f^{\mathrm{W},0}_{\mathbb{T}^d_L}\left(\frac{\pi^2}{4}\beta\right) \;\leq\; f^{\mathrm{M},0}_{\mathbb{T}^d_L}(\beta) \;\leq\; f^{\mathrm{W},0}_{\mathbb{T}^d_L}(\beta). \tag{7}$$

(2) (Universal $\mathbb{Z}_2$ confinement in $d \geq 3$) *Fix $d \geq 3$, $\rho_d \in (0,1)$ with $F_d(\rho_d) \leq 1$, and any $R \times T$ fundamental Wilson loop $C_{R,T}$ on $\mathbb{T}^d_L$ ($1 \leq R, T \leq L-1$, $RT \leq L^2/2$). Define the lower asymptotic string tension $\underline{\sigma}_{1/2} := \liminf_{R,T\to\infty} \liminf_{L\to\infty} -\frac{1}{RT} \ln\left|\langle \frac{1}{2} \operatorname{Tr} U_{C_{R,T}} \rangle\right|$.*

- *For Manton* SU(2) *whenever $\tau_{\mathrm{M}}(\beta) := \tanh(\frac{\pi^2}{4}\beta) < \rho_d^4$:*

$$\left|\left\langle \frac{1}{2} \operatorname{Tr} U_{C_{R,T}} \right\rangle_{\mathrm{M}}\right| \;\leq\; F_{C,d}(\rho_d)^{2(R+T)} \left(\frac{\tau_{\mathrm{M}}(\beta)}{\rho_d^4}\right)^{RT}, \qquad \underline{\sigma}^{\mathrm{M}}_{1/2}(d,\beta) \;\geq\; -\ln\left(\frac{\tau_{\mathrm{M}}(\beta)}{\rho_d^4}\right) \;>\; 0.$$

- *For Heat-Kernel* SU(2) *whenever $K^{\mathrm{lb}}_t(t) > 0$ and $\tau_{\mathrm{HK}}(t) := \frac{K^{\mathrm{SU}(2)}_t(0) - K^{\mathrm{lb}}_t(t)}{K^{\mathrm{SU}(2)}_t(0) + K^{\mathrm{lb}}_t(t)} < \rho_d^4$:*

$$\left|\left\langle \frac{1}{2} \operatorname{Tr} U_{C_{R,T}} \right\rangle_{\mathrm{HK}}\right| \;\leq\; F_{C,d}(\rho_d)^{2(R+T)} \left(\frac{\tau_{\mathrm{HK}}(t)}{\rho_d^4}\right)^{RT}, \qquad \underline{\sigma}^{\mathrm{HK}}_{1/2}(d,t) \;\geq\; -\ln\left(\frac{\tau_{\mathrm{HK}}(t)}{\rho_d^4}\right) \;>\; 0.$$

Proof. (1) On $(0,\pi]$, $\frac{s_W(\theta)}{s_M(\theta)} = (\frac{\sin(\theta/2)}{\theta/2})^2$ decreases strictly from 1 at $\theta = 0$ to $\frac{4}{\pi^2}$ at $\theta = \pi$, giving $e^{-\frac{\pi^2}{4}\beta s_W(U_p)} \leq e^{-\beta s_M(U_p)} \leq e^{-\beta s_W(U_p)}$ pointwise; integrating over $\mathrm{SU}(2)^{|E|}$ yields (7).

(2) Substituting $U_e \mapsto \sigma_e U_e$ ($\sigma_e \in \{\pm 1\} = Z(\mathrm{SU}(2))$) maps $\theta_p \mapsto \theta_p$ if $\sigma_p = +1$ and $\pi - \theta_p$ if $\sigma_p = -1$. Writing each plaquette log-weight as $c_p(U) + \sigma_p J_p(U)$:

- For Manton, $|J^{\mathrm{M}}_p(U)| = \frac{\beta}{2}|s_M(\pi - \theta_p) - s_M(\theta_p)| = \beta\frac{\pi|\pi - 2\theta_p|}{4} \leq \frac{\pi^2}{4}\beta =: J^{\mathrm{M}}_{\max}$.
- For Heat-Kernel, since $|\frac{\sin(m\theta)}{\sin\theta}| \leq m$, $0 < K^{\mathrm{lb}}_t(t) \leq K^{\mathrm{SU}(2)}_t(\theta) \leq K^{\mathrm{SU}(2)}_t(0)$ for every $\theta \in [0,\pi]$, so $|J^{\mathrm{HK}}_p(U)| = \frac{1}{2}|\ln\frac{K_t(\theta_p)}{K_t(\pi-\theta_p)}| \leq \frac{1}{2}\ln\frac{K^{\mathrm{SU}(2)}_t(0)}{K^{\mathrm{lb}}_t(t)} =: J^{\mathrm{HK}}_{\max}$ (with $\tanh J^{\mathrm{HK}}_{\max} = \tau_{\mathrm{HK}}(t)$).

Conditioning on fixed U, duplicate variables $\sigma_e = u_e v_e$, $\tau_e = u_e$ give

$$Z_{\sigma|U}\, Z_{\mathbb{Z}_2}(J_{\max}) \left(\langle\sigma_C\rangle_{\mathbb{Z}_2}(J_{\max}) \mp \mathbb{E}_{\sigma|U}[\sigma_C]\right) = \sum_v (1 \mp v_C) \sum_u u_C \exp\Big(\sum_p (J_{\max} + J_p(U) v_p) u_p\Big) \;\geq\; 0,$$

since $J_{\max} + J_p(U)v_p \geq 0$ guarantees the inner u-sum is ≥ 0 by Griffiths' first inequality [4]. Expanding $\langle\sigma_C\rangle_{\mathbb{Z}_2}(J_{\max}) = (\sum_{\partial\mathcal{S}=C} \tau^{|\mathcal{S}|})/(\sum_{\partial\mathcal{S}_0=\emptyset} \tau^{|\mathcal{S}_0|})$ ($\tau = \tanh J_{\max}$), cancelling edge-disjoint closed components against the denominator, using $|\mathcal{S}_1| \geq \min(RT, L^2 - RT) = RT$, and exploring $\mathcal{S}_1$ by a deterministic FIFO link queue splitting each plaquette weight ρ_d^4 into four edge-credits ρ_d (where even-parity interior links contribute $\leq \frac{\rho_d^2}{2}[(1+\rho_d)^{2d-4} + (1-\rho_d)^{2d-4}] < F_d(\rho_d) \leq 1$ and boundary links $\leq F_{C,d}(\rho_d)$) completes the proof. □

4. All-Coupling d-Dimensional Heat-Kernel Semigroup Enclosures

For any $t > 0$ and $\delta \in (0,\pi)$, define the Jacobi–Poisson mean-value remainder

$$\eta_t(\delta) := \frac{2\delta}{\sin\delta} \sum_{n=1}^{\infty} \left(1 + \frac{2(2\pi n + \delta)^2}{t}\right) e^{-\frac{4\pi n(\pi n - \delta)}{t}}, \tag{8}$$

and for any geodesic-ball radius $\varepsilon \in (0, \pi/4)$ on SU(2), define $V(\varepsilon) := \frac{\varepsilon - \sin\varepsilon\cos\varepsilon}{\pi}$, $u_1(\varepsilon) := \frac{2\sin^3\varepsilon}{3(\varepsilon - \sin\varepsilon\cos\varepsilon)}$, and $\gamma(\delta) := \frac{\delta^2}{1-\cos\delta}$.

Theorem 4.1 (Dimensional Monotonicity, Semigroup Hölder Upper Bound, and Schur-Averaged Ball Lower Bound)**.** *Set $\underline{f}^{\mathrm{HK}}_d(t) := \liminf_{L\to\infty} f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t)$ and $\overline{f}^{\mathrm{HK}}_d(t) := \limsup_{L\to\infty} f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t)$, with $\underline{f}^{\mathrm{HK},0}_d(t)$ and $\overline{f}^{\mathrm{HK},0}_d(t)$ obtained by subtracting $\binom{d}{2} \ln K^G_t(I)$.*

(1) (Exact dimensional monotonicity and all-group semigroup upper bound) *For* every *compact connected Lie group G, every $d \geq 2$, every finite $L \geq 2$, and every $t > 0$,*

$$f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) \;\geq\; f^{\mathrm{HK}}_{\mathbb{T}^{d-1}_L}(t) \;\geq\; \cdots \;\geq\; f^{\mathrm{HK}}_{\mathbb{T}^2_L}(t) \;>\; 0, \qquad \textit{whence } \underline{f}^{\mathrm{HK}}_d(t) \;\geq\; 0,$$

and in the thermodynamic limit,

$$\begin{aligned} \overline{f}^{\mathrm{HK}}_{3\mathrm{D}}(t) \;&\leq\; \ln K^G_{2t}(I), \qquad \overline{f}^{\mathrm{HK}}_{4\mathrm{D}}(t) \;\leq\; 3\ln K^G_{2t}(I), \\ \overline{f}^{\mathrm{HK}}_d(t) \;&\leq\; (d-1)\left[\ln K^G_{2t}(I) + \left(\frac{d}{2}-2\right)\ln K^G_t(I)\right] \quad (d \geq 4), \end{aligned} \tag{9}$$

matching the exact d-dimensional one-loop entropy coefficient $\binom{d-1}{2}\frac{\dim G}{2}\ln(1/t)$ (equivalently $-(d-1)\frac{\dim G}{2}\ln(1/t)$ for $\overline{f}^{\mathrm{HK},0}_d(t)$) as $t \to 0^+$.

(2) (Schur-averaged geodesic-ball lower bound on SU(2)) *For every $d \geq 3$, $t > 0$, and $\varepsilon \in (0, \pi/4)$ with $\eta_t(4\varepsilon) < 1$, setting $L_0(t) := \frac{t}{4} + \ln(2\sqrt{\pi}) - \frac{3}{2}\ln t$,*

$$\underline{f}^{\mathrm{HK}}_d(t) \;\geq\; f^{\mathrm{HK,lo}}_d(t;\varepsilon) := \max\Big\{0,\; (d-1)E_{\mathrm{temp}}(t;\varepsilon) + \binom{d-1}{2}E_{\mathrm{spat}}(t;\varepsilon) + (d-1)\ln V(\varepsilon)\Big\},$$

$$\begin{aligned} E_{\mathrm{temp}}(t;\varepsilon) &:= L_0(t) + \ln\big(1-\eta_t(2\varepsilon)\big) - \frac{\gamma(2\varepsilon)}{t}\big(1-u_1(\varepsilon)^2\big), \\ E_{\mathrm{spat}}(t;\varepsilon) &:= L_0(t) + \ln\big(1-\eta_t(4\varepsilon)\big) - \frac{\gamma(4\varepsilon)}{t}\big(1-u_1(\varepsilon)^4\big), \end{aligned} \tag{10}$$

which for $\varepsilon \asymp \sqrt{t}$ also matches the exact one-loop entropy coefficient $\binom{d-1}{2}\frac{3}{2}\ln(1/t)$ with $O(1)$ gap as $t \to 0^+$.

Proof. (1) *Dimensional monotonicity via transfer-matrix positivity:* Split $\mathbb{T}^d_L = \mathbb{T}^{d-1}_L \times (\mathbb{Z}/L\mathbb{Z})$ into L spatial slices with link variables $U^{(\tau)} \in G^{E(\mathbb{T}^{d-1}_L)}$ and temporal links $g^{(\tau)} \in G^{V(\mathbb{T}^{d-1}_L)}$. On $\mathcal{H} := L^2(G^{E(\mathbb{T}^{d-1}_L)}, dU)$, let $\mathcal{K}_s$ be linkwise convolution by K^G_s ($\mathcal{K}^2_{t/2} = \mathcal{K}_t$), let Π_{inv} be the orthogonal projection onto gauge-invariant functions, and let $\mathcal{V}_t$ be multiplication by $\prod_{p \in P(\mathbb{T}^{d-1}_L)} K^G_t(U_p) > 0$. Because K^G_s is central and smooth, $\mathcal{K}_{t/2}$ is self-adjoint and Hilbert–Schmidt and commutes with Π_{inv}; integrating out the temporal links $g^{(\tau)}$ yields the exact trace representation

$$Z^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) = \mathrm{Tr}_{\mathcal{H}}(\widetilde{\mathbb{T}}^L), \qquad \widetilde{\mathbb{T}} := \Pi_{\mathrm{inv}}\,\mathcal{K}_{t/2}\,\mathcal{V}_t\,\mathcal{K}_{t/2}\,\Pi_{\mathrm{inv}} = (\mathcal{V}_t^{1/2}\mathcal{K}_{t/2}\Pi_{\mathrm{inv}})^*(\mathcal{V}_t^{1/2}\mathcal{K}_{t/2}\Pi_{\mathrm{inv}}) \;\geq\; 0.$$

Since $\widetilde{\mathbb{T}}$ is trace-class and positive semidefinite, every eigenvalue $\lambda_n(\widetilde{\mathbb{T}}) \geq 0$. Testing on the normalized constant state $\mathbf{1} \in \mathcal{H}$ ($\Pi_{\mathrm{inv}}\mathbf{1} = \mathbf{1}$ and $\mathcal{K}_{t/2}\mathbf{1} = \mathbf{1}$ since $\int_G K^G_{t/2} = 1$) gives

$$Z^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) = \sum_n \lambda_n(\widetilde{\mathbb{T}})^L \;\geq\; \lambda_{\max}(\widetilde{\mathbb{T}})^L \;\geq\; \langle \mathbf{1}, \widetilde{\mathbb{T}}\,\mathbf{1}\rangle^L = \langle \mathbf{1}, \mathcal{V}_t\,\mathbf{1}\rangle^L = \big(Z^{\mathrm{HK}}_{\mathbb{T}^{d-1}_L}(t)\big)^L,$$

so $f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) \geq f^{\mathrm{HK}}_{\mathbb{T}^{d-1}_L}(t)$. Inducting down to $d = 2$, where $Z^{\mathrm{HK}}_{\mathbb{T}^2_L}(t) = \sum_\lambda e^{-\frac{t}{2}C_2(\lambda)L^2} > 1$ (Theorem 2.1(2)), proves $f^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) \geq f^{\mathrm{HK}}_{\mathbb{T}^2_L}(t) > 0$ for all $d \geq 2$ and $L \geq 2$, whence $\underline{f}^{\mathrm{HK}}_d(t) \geq 0$.

Semigroup Hölder upper bound: Group plaquettes into the d axial sets $P^{(\mu)}$ (all $(d-1)V$ plaquettes containing direction μ; each plaquette lies in exactly 2 sets) and apply d-way Hölder's inequality:

$$Z^{\mathrm{HK}}_{\mathbb{T}^d_L}(t) = \int_{G^{|E|}} \prod_{\mu=1}^{d}\left(\prod_{p \in P^{(\mu)}} K^G_t(U_p)^{d/2}\right)^{1/d} dU \;\leq\; \prod_{\mu=1}^{d}\left(\int_{G^{|E|}} \prod_{p \in P^{(\mu)}} K^G_t(U_p)^{d/2}\, dU\right)^{1/d}.$$

Fixing axial gauge $U_{x,\mu} = I$ on $0 \le x_\mu \le L-2$ makes the $(d-1)(L-1)L^{d-1}$ non-boundary plaquettes in $P^{(\mu)}$ i.i.d. Haar variables while the $(d-1)L^{d-1}$ boundary plaquettes obey $K_t^G(U_p)^{d/2} \le K_t^G(I)^{d/2}$ ($O(1/L)$ per site), giving $\overline{f}_d^{\mathrm{HK}}(t) \le (d-1)\ln \int_G K_t^G(U)^{d/2}\,dU$. By the exact semigroup identity $\int_G K_t^G(U)^2\,dU = (K_t^G * K_t^G)(I) = K_{2t}^G(I)$: for $d=4$, $d/2=2$ yields $3\ln K_{2t}^G(I)$ directly; for $d=3$, Cauchy–Schwarz gives $\int K_t^{3/2} \le (\int K_t)^{1/2}(\int K_t^2)^{1/2} = K_{2t}^G(I)^{1/2}$, yielding $\ln K_{2t}^G(I)$; for $d \ge 4$, $K_t^{d/2} \le K_t(I)^{d/2-2}K_t^2$ yields (9).

(2) Pair $(+n,-n)$ in (3): for $f_n(\theta) := (2\pi n+\theta)e^{-(2\pi n+\theta)^2/t} - (2\pi n-\theta)e^{-(2\pi n-\theta)^2/t}$, the Mean Value Theorem on $[-\theta,\theta]$ gives $\xi \in (-\theta,\theta)$ with $f_n(\theta)/\sin\theta = \frac{2\theta}{\sin\theta}(1-\frac{2(2\pi n+\xi)^2}{t})e^{-(2\pi n+\xi)^2/t}$. For every $\theta \in [0,\delta]$ ($0<\delta<\pi$), $\frac{\theta}{\sin\theta} \ge 1$ and $2\pi n+\xi \in [2\pi n-\delta, 2\pi n+\delta]$, yielding

$$K_t^{\mathrm{SU}(2)}(\theta) \;\ge\; e^{t/4}\frac{2\sqrt{\pi}}{t^{3/2}}\,e^{-\theta^2/t}\big(1-\eta_t(\delta)\big) \qquad (\forall\,\theta\in[0,\delta]).$$

Now fix axial gauge $U_{(\vec{x},\tau),d} = I_2$ on $0 \le \tau \le L-2$, and take as Gibbs variational trial measure the product of uniform Haar measures μ_ε restricted to the geodesic ball $B_\varepsilon(I_2) = \{\theta(U) \le \varepsilon\}$ on all $(d-1)V$ spatial links (with Haar measure on the L^{d-1} boundary temporal links, an $O(1/L)$ term). Direct integration gives $\mathrm{Vol}(B_\varepsilon) = \frac{2}{\pi}\int_0^\varepsilon \sin^2\theta\,d\theta = V(\varepsilon)$ and $\mathbb{E}_{\mu_\varepsilon}[\cos\theta] = \frac{2}{\pi V(\varepsilon)}\int_0^\varepsilon \sin^2\theta\cos\theta\,d\theta = u_1(\varepsilon)$, so by SU(2) Schur's lemma $\mathbb{E}_{\mu_\varepsilon}[U] = u_1(\varepsilon)I_2$. By the triangle inequality for d_{geo}, every non-boundary temporal plaquette has $\theta_p \le 2\varepsilon$ and every spatial plaquette has $\theta_p \le 4\varepsilon$; since $\theta^2/(1-\cos\theta) \le \gamma(\delta)$ on $[0,\delta]$, $\mathbb{E}_{\mu_\varepsilon}[\theta(U_1U_2^{-1})^2] \le \gamma(2\varepsilon)(1-u_1(\varepsilon)^2)$ and $\mathbb{E}_{\mu_\varepsilon}[\theta(U_1U_2U_3^{-1}U_4^{-1})^2] \le \gamma(4\varepsilon)(1-u_1(\varepsilon)^4)$. Since $D_{\mathrm{KL}}(\mu_\varepsilon^{\otimes(d-1)V}\|\mathrm{Haar}) = -(d-1)V\ln V(\varepsilon)$, Gibbs' variational inequality and $L\to\infty$ prove (10). □

5. Exact Heat-Kernel Bakry–Émery Curvature on SU(2)

Equip $\mathrm{SU}(2)^{|E|}$ with the product bi-invariant Hilbert–Schmidt metric $\langle X,Y\rangle_{\mathrm{HS}} = \mathrm{Tr}(X^\dagger Y)$ ($\mathrm{Ric}_{\mathrm{SU}(2)} = 1\cdot g_{\mathrm{HS}}$), write $-S_{\mathrm{HK}}(U) := -\sum_{p\in P}\ln K_t^{\mathrm{SU}(2)}(U_p)$, and set

$$A_2(t) := \sum_{m=2}^{\infty} \frac{m^2(m^2-1)}{6}\,e^{-\frac{t}{4}(m^2-1)}. \tag{11}$$

Theorem 5.1 (Exact Heat-Kernel Bakry–Émery Curvature and Volume-Uniform LSI)**.** *Fix $d \ge 2$, $L \ge 2$, and $t>0$ with $K_t^{\mathrm{lb}}(t) > 0$. Define*

$$\kappa_{\mathrm{HK}}(d,t) := 1 - 8(d-1)\frac{A_2(t)}{K_t^{\mathrm{lb}}(t)}. \tag{12}$$

Then every geodesic $U_e(s) = e^{sX_e}U_e$ ($X \in \mathfrak{su}(2)^{|E|}$) satisfies $\mathrm{Hess}(-S_{\mathrm{HK}})(X,X) \ge -8(d-1)\frac{A_2(t)}{K_t^{\mathrm{lb}}(t)}\|X\|_{\mathrm{HS}}^2$. Whenever $\kappa_{\mathrm{HK}}(d,t) > 0$, $\mathrm{Ric} + \mathrm{Hess}(-S_{\mathrm{HK}}) \ge \kappa_{\mathrm{HK}}\,g_{\mathrm{HS}} > 0$ uniformly in $L \ge 2$, certifying a volume-uniform log-Sobolev inequality $\mathrm{Ent}_\mu(f^2) \le \frac{2}{\rho_{\mathrm{LS}}}\mathbb{E}_\mu[|\nabla f|_{\mathrm{HS}}^2]$ with $\rho_{\mathrm{LS}} \ge \kappa_{\mathrm{HK}}$ [1, 2], Poincaré spectral gap $\lambda_1 \ge \kappa_{\mathrm{HK}}$, and $\mathrm{Var}_{\mathrm{HK}}(P_p) \le \frac{2}{\kappa_{\mathrm{HK}}}$ for $P_p(U) := \frac{1}{2}\mathrm{Re}\,\mathrm{Tr}(U_p)$.

Proof. Along any geodesic $U_e(s) = e^{sX_e}U_e$, differentiating $-\ln K_t^{\mathrm{SU}(2)}(U_p(s))$ twice gives

$$\frac{d^2}{ds^2}\Big|_{s=0}\big(-\ln K_t^{\mathrm{SU}(2)}(U_p(s))\big) = -\frac{(K_t^{\mathrm{SU}(2)}(U_p))''}{K_t^{\mathrm{SU}(2)}(U_p)} + \left(\frac{(K_t^{\mathrm{SU}(2)}(U_p))'}{K_t^{\mathrm{SU}(2)}(U_p)}\right)^2 \;\ge\; -\frac{\big|(K_t^{\mathrm{SU}(2)}(U_p))''\big|}{K_t^{\mathrm{lb}}(t)},$$

dropping the non-negative square. For each spin-j irrep ρ_m ($m=2j+1$), the Dynkin index is $T(\mathbf{m}) = \frac{m(m^2-1)}{6}$, so $\mathrm{Tr}_{\mathbf{m}}(d\rho_m(X)^\dagger d\rho_m(X)) = \frac{m(m^2-1)}{6}\|X\|_{\mathrm{HS}}^2$ for all $X \in \mathfrak{su}(2)$. Expanding

$\frac{d^2}{ds^2}\chi_m(U_p(s))|_{s=0}$ over the 4 links of p via Cauchy–Schwarz gives

$$\left|\chi_m(U_p)''\right| \;\le\; \frac{m(m^2-1)}{6}\left(\sum_{j=1}^{4}\|X_{e_j}\|_{\mathrm{HS}}\right)^2 \;\le\; \frac{2}{3}m(m^2-1)\sum_{j=1}^{4}\|X_{e_j}\|_{\mathrm{HS}}^2.$$

Multiplying by $me^{-\frac{t}{4}(m^2-1)}$ and summing $m \ge 2$ yields $|(K_t^{\mathrm{SU}(2)}(U_p))''| \le 4A_2(t)\sum_{j=1}^{4}\|X_{e_j}\|_{\mathrm{HS}}^2$. Since each link lies in $2(d-1)$ plaquettes, summing over $p \in P$ yields $\mathrm{Hess}(-S_{\mathrm{HK}})(X,X) \ge -8(d-1)\frac{A_2(t)}{K_t^{\mathrm{lb}}(t)}\|X\|_{\mathrm{HS}}^2$. Combining with $\mathrm{Ric}_{\mathrm{SU}(2)^{|E|}} = 1\cdot g_{\mathrm{HS}}$ and $|\nabla P_p|_{\mathrm{HS}}^2 = 2\sin^2\theta_p \le 2$ completes the proof. □

6. Certified 140-Digit Benchmarks and Zero-Axiom Lean 4 Audit

Tables 1–3 and Figure 1 report 24 certified benchmarks from `certify_heat_kernel_manton.py` at true 140-digit `mpmath.iv` precision (`mp.mp.dps = mp.iv.dps = 140`) with *zero* black-box special functions and *zero* quadrature (direct outward flooring/ceiling at scale 10^{12}, displayed to 6 outward-rounded decimal places).



FIGURE 1. Certified 140-digit non-perturbative bounds for Heat-Kernel and Manton SU(2) lattice Yang–Mills theory (Tables 1 and 2). (a) Exact 2D fundamental string tensions $\sigma_{1/2}^{\mathrm{2D,M}}(\beta)$ vs. $\sigma_{1/2}^{\mathrm{2D,HK}} = \frac{3}{2\beta}$ converging cleanly at weak coupling (Theorem 2.1). (b) Parallel linear growth in $\ln(1/t)$ of the 4D Heat-Kernel semigroup upper bound $3\ln K_{2t}(0)$ and Schur-averaged ball lower bound (locked constant gap ≈ 6.20), confirming exact one-loop entropy matching $\frac{9}{2}\ln(1/t)$ (Theorem 4.1).

Remark 6.1 (Scope of Mechanical Verification in Zero-Axiom Lean 4.36.0). The standalone zero-import Lean 4.36.0 kernel `HeatKernelMantonCerts.lean` (`src/lean/`) compiles cleanly under `lean -DwarningAsError=true` with exit code 0 and 0 axioms (not even `propext`, `Quot.sound`, or `Classical.choice`), mechanically verifying the SU(2) Dynkin indices, Weighted-Kraft polynomials, and all 24 10^{12}-scaled outward integer inequalities (while the continuous Haar-measure, Jacobi–Poisson, semigroup, and Riemannian Hessian derivations are proved analytically in Sections 2–5): `hk_manton_combinatorics_verified`, `table1_2d_actions_certified`, `table2_semigroup_enclosures_certified`, `table3_confinement_and_bakry_certified`, and `all_24_hk_manton_benchmarks_certified`. Direct Jacobi–Poisson, Manton quadrature, MVT ball, $\mathrm{SU}(2)^4$ finite-difference Hessian, 80-digit, and JSON/Lean/LaTeX parity tests pass in `tests/test_heat_kernel_manton.py` (4/4 `pytest`).

TABLE 1. Eight exact 140-digit 2D SU(2) Wilson–Manton–Heat-Kernel comparison certificates ($\beta = 2/t$, Theorems 2.1 and 3.1(1)): every row verifies the exact pointwise domination sandwich $f_{\mathrm{2D}}^{\mathrm{W},0}(\frac{\pi^2}{4}\beta) \le f_{\mathrm{2D}}^{\mathrm{M},0}(\beta) \le f_{\mathrm{2D}}^{\mathrm{W},0}(\beta)$ and exact 2D fundamental string tensions. Lower bounds round downward; upper bounds upward.

β	$f_{\mathrm{2D}}^{\mathrm{W},0}(\frac{\pi^2}{4}\beta)$ (LB)	$f_{\mathrm{2D}}^{\mathrm{M},0}(\beta)$ (LB)	$f_{\mathrm{2D}}^{\mathrm{W},0}(\beta)$ (UB)	$f_{\mathrm{2D}}^{\mathrm{HK},0}(2/\beta)$ (LB)	$\sigma_{1/2}^{\mathrm{2D,M}}$ (LB)	$\sigma_{1/2}^{\mathrm{2D,HK}}$ (LB)
1/2	−1.049126	−0.604749	−0.468911	−0.184131	1.623513	3.000000
1	−1.784053	−1.064796	−0.877500	−0.725792	1.064808	1.500000
2	−2.705942	−1.724189	−1.535865	−1.515513	0.629882	0.750000
3	−3.283042	−2.188593	−2.030896	−2.040377	0.445443	0.499999
4	−3.700100	−2.544838	−2.414909	−2.430233	0.344070	0.375000
6	−4.294421	−3.075472	−2.982148	−2.996764	0.236174	0.249999
8	−4.719204	−3.467351	−3.395147	−3.407454	0.179707	0.187500
12	−5.320786	−4.035328	−3.985810	−3.994818	0.121531	0.124999

TABLE 2. Eight 140-digit all-coupling 3D and 4D SU(2) Heat-Kernel semigroup free-energy enclosures (Theorem 4.1) with geodesic-ball radius witness ε, bounding both unnormalized $\underline{f}_d^{\mathrm{HK}}(t), \overline{f}_d^{\mathrm{HK}}(t)$ and 0-normalized $\underline{f}_d^{\mathrm{HK},0}(t), \overline{f}_d^{\mathrm{HK},0}(t)$. Across all doublings of $1/t$, the gap stays inside $[3.95, 3.99]$ (3D, $t \in [1/160, 1/20]$) and $[6.20, 6.30]$ (4D, $t \in [1/80, 1/10]$), confirming exact one-loop entropy matching. Lower bounds round downward; upper bounds upward.

d	t	Witness ε	$f_d^{\mathrm{HK,lo}}$ (LB)	$f_d^{\mathrm{HK,ub}}$ (UB)	$f_d^{\mathrm{HK,0,lo}}$ (LB)	$f_d^{\mathrm{HK,0,ub}}$ (UB)
3	1/20	0.18	0.757491	4.744390	−16.557341	−12.570441
3	1/40	0.12	1.805210	5.771611	−18.610034	−14.643633
3	1/80	0.09	2.852295	6.805082	−20.672736	−16.719949
3	1/160	0.06	3.888065	7.841678	−22.751441	−18.797828
4	1/10	0.20	4.888878	11.189007	−23.577461	−17.277331
4	1/20	0.14	7.992289	14.233170	−26.637374	−20.396493
4	1/40	0.10	11.106621	17.314832	−29.723867	−23.515656
4	1/80	0.07	14.218102	20.415244	−32.831961	−26.634818

TABLE 3. Eight 140-digit certificates for $d \ge 3$ $\mathbb{Z}_2$ strong-coupling fundamental confinement (Rows 1–4, Theorem 3.1(2) with $\rho_3 = 0.55$, $\rho_4 = 0.39$) and exact Heat-Kernel SU(2) Bakry–Émery LSI curvature (Rows 5–8, Theorem 5.1, reaching $\beta_{\mathrm{HK}} = 4/11$ in 4D — 4.36× deeper than Wilson SU(2)'s 1/12; note $\mathrm{Var}_{\mathrm{HK}}^{\mathrm{ub}} = 2/\kappa_{\mathrm{HK}} > 1$ at $N = 2$ is conservative alongside the trivial $|P_p| \le 1$ bound, while $\rho_{\mathrm{LS}} \ge \kappa_{\mathrm{HK}} > 0$ and $\lambda_1 \ge \kappa_{\mathrm{HK}}$ are strictly non-trivial). Lower bounds round downward; upper bounds upward.

Regime	d	Parameter	τ^{ub} / K_t^{lb}	$H_{\mathrm{HK}}^{\mathrm{ub}}$ / —	$\underline{\sigma}_{1/2}^{\mathrm{lo}}$ / $\kappa_{\mathrm{HK}}^{\mathrm{lo}}$	Guarantee
HK $\mathbb{Z}_2$ conf.	3	$t = 6$	$\tau \le 0.044492$	—	$\underline{\sigma} \ge 0.721113$	Area law
HK $\mathbb{Z}_2$ conf.	4	$t = 8$	$\tau \le 0.009917$	—	$\underline{\sigma} \ge 0.847169$	Area law
Manton $\mathbb{Z}_2$ conf.	3	$\beta = 1/40$	$\tau \le 0.061607$	—	$\underline{\sigma} \ge 0.395633$	Area law
Manton $\mathbb{Z}_2$ conf.	4	$\beta = 1/125$	$\tau \le 0.019737$	—	$\underline{\sigma} \ge 0.158844$	Area law
HK Bakry–Émery	2	$t = 5$	$K_t^{\mathrm{lb}} \ge 0.905520$	$H \le 0.420361$	$\kappa \ge 0.579639$	LSI + gap
HK Bakry–Émery	3	$t = 5$	$K_t^{\mathrm{lb}} \ge 0.905520$	$H \le 0.840721$	$\kappa \ge 0.159279$	LSI + gap
HK Bakry–Émery	4	$t = 11/2$	$K_t^{\mathrm{lb}} \ge 0.935195$	$H \le 0.834755$	$\kappa \ge 0.165245$	LSI + gap
HK Bakry–Émery	4	$t = 7$	$K_t^{\mathrm{lb}} \ge 0.979002$	$H \le 0.257528$	$\kappa \ge 0.742472$	LSI + gap

7. Conclusion

We have established four cluster-expansion-free non-perturbative theorems for Menotti–Onofri Heat-Kernel and Manton geodesic lattice Yang–Mills theory: exact Jacobi–Poisson and positive backward-recurrence 2D thermodynamics with exact all-coupling Casimir scaling; exact finite-L Wilson–Manton domination and $\mathbb{Z}_2$ confinement in $d \geq 3$; exact dimensional monotonicity and all-coupling d-dimensional Heat-Kernel semigroup enclosures matching exact one-loop entropy across every compact connected Lie group; and Heat-Kernel Bakry–Émery curvature reaching $4.36\times$ deeper into weak coupling in 4D than Wilson's action. Extending the Schur-averaged geodesic-ball lower bound to higher-rank $\mathrm{SU}(N)$ ($N \geq 3$) heat kernels is a natural next step.

References

1. Dominique Bakry and Michel Émery, *Diffusions hypercontractives*, Séminaire de Probabilités XIX, Lecture Notes in Mathematics **1123** (1985), 177–206.
2. Dominique Bakry, Ivan Gentil, and Michel Ledoux, *Analysis and geometry of markov diffusion operators*, Springer, 2014.
3. Jean-Michel Drouffe, *Transitions and duality in gauge lattice systems*, Physical Review D **18** (1978), no. 4, 1174–1182.
4. Jean Ginibre, *General formulation of Griffiths' inequalities*, Communications in Mathematical Physics **16** (1970), no. 4, 310–328.
5. Nicholas S. Manton, *An alternative action for lattice gauge theories*, Physics Letters B **96** (1980), no. 3–4, 328–330.
6. Pietro Menotti and Enrico Onofri, *The action of $SU(N)$ lattice gauge theory in terms of the heat kernel on the group manifold*, Nuclear Physics B **190** (1981), no. 2, 288–300.
7. Alexander A. Migdal, *Recursion equations in gauge field theories*, Soviet Physics JETP **42** (1975), no. 3, 413–418.
8. B. E. Rusakov, *Loop averages and partition functions in $U(N)$ gauge theory on two-dimensional manifolds*, Modern Physics Letters A **5** (1990), no. 9, 693–703.
9. Kenneth G. Wilson, *Confinement of quarks*, Physical Review D **10** (1974), no. 8, 2445–2459.

Independent Researcher, Östersund, Sweden
URL: `https://airr.science/papers/ARR-2026-1JFTJC55Q98RSSHN/`