Universal $O(1)$ Closed-Form Hypergraph Bounds, Dual Gap-Closing BIBD/RBIBD Families, and Even-t Residual-Degree Refinement for $(n, t, w, w-2)$-Cooling Codes

Lluis Eriksson
Independent Researcher (lluiseriksson@gmail.com)
Public Repository: https://github.com/arr-research/arr-research.github.io

November 28, 2026

AIRR:ARR-2026-3Z52WEEHN48QF96M | v1 | 2026-11-28

Abstract

Liu, Zhang, Yu, Zhang, and Ji (arXiv:2611.34386v1) recently derived new upper bounds and optimal design constructions for $(n, t, w, w-2)$-constant-power error-correcting cooling (CPECC) codes across both the small-t graph regime ($t < w-3$, Theorem 3.4) and the large-t hypergraph regime ($t \geq w-3$, Theorem 4.6), explicitly leaving two open problems in Section 5: (1) constructing optimal $(n, t, w, w-2)$-CPECC codes for *even* $t < w-3$, and (2) evaluating the hypergraph penalty $z_{w-1}(t+1)$ and constructing optimal codes inside the Frobenius gap $w-3 \leq t < (w-3)\lfloor (w-2)/3 \rfloor - 1$. We prove four new theorems directly resolving Open Problem 2 and advancing Open Problem 1: (1) a *Universal $O(1)$ Closed-Form Theorem* proving $z_{w-1}(x) = \min\big(x - p(w-1),\ \max(0, (p+1)(w-2) - x)\big)$ with $p = \lfloor x/(w-1) \rfloor$ unconditionally for *all* $w \geq 4, x \geq 0$ (formally proved in Lean 4.36.0 T01–T05), eliminating Liu et al.'s Frobenius-number restriction entirely; (2) a *Dual Gap-Closing Optimal BIBD/RBIBD Theorem* proving exact attainment of the Theorem 4.6 bound $C'(n, t, w, w-2) = \frac{n(n-1)}{2((2w-3)(t+1)+\ell)}$ inside Liu et al.'s open gap across two infinite coprime families—unconditional *Family A* ($t+1 = \ell(w-3)$, $2 \leq \ell \leq \lfloor (w-1)/3 \rfloor$, $\gcd(\ell, 2w-5) = 1$, $(n, 2w-4, 1)$-RBIBDs, entering the gap for all $w \geq 11$) and a **brand-new Dual Family B** ($t+1 = \ell w$, $1 \leq \ell \leq \lfloor (w-2)/3 \rfloor$, $(n, 2w-1, 1)$-BIBDs at $\ell = 1$ with no resolvability or coprimality restriction—entering the open gap for *every* $w \geq 8$—and $(n, 2w-1, 1)$-RBIBDs for $\ell \geq 2$ with $\gcd(\ell, 2w-2) = 1$); (3) an *Even-t Residual-Degree Jump and Exact h_3-Attaining Transversal-Design Theorem* proving $B(w, 2r) = (w+r-1)^2$, giving a self-contained proof that any codeword graph with $\delta(G) \geq w-1$, Property $A(2r)$, and $\Delta(G) \leq w+r-2$ forces $|V(G)| \geq n_{\min} := \lceil (2r+1)(w+r-1)/r \rceil$ with an exact boundary identity $\frac{w-1}{2} n_{\min} = (w+r-1)^2$ at $w = 2r^2+1$ and strict edge jump $\lceil (w-1)n_{\min}/2 \rceil > B(w, 2r)$ for *all* $w \geq 2r^2+2$ ($3\binom{w}{2} > w^2$ for all $w \geq 6$ at $t = 2$), constructing an **explicit unconditional infinite family** of optimal $t = 2$ codes from projective transversal designs $\mathrm{TD}(q+1, q)$ ($q = w+1 \geq 7$ odd prime power, attaining exact equality with Liu et al.'s Theorem 3.4 bound h_3 with zero asymptotic existence hypothesis!), and proving a sharp non-existence dichotomy at $r = 2$ ($t = 4$); and (4) an *Explicit Finite Open-Gap Certification Theorem* constructing and verifying exact optimal finite affine-plane codes including $\mathrm{AG}(2, 17)$ ($n = 289, t = 8, w = 9, M = 306$) and $\mathrm{AG}(2, 19)$ ($n = 361, t = 9, w = 10, M = 380$) strictly inside Liu et al.'s open gap. All 20 core theorems (T01–T20) are formally proved in Lean 4.36.0 (0 sorry, 0 grind, 0 custom axioms), paired with an AST-pure zero-float verifier, 16 SHA-256 certificates, and 70 passing tests.

1 Introduction

Constant-power error-correcting cooling (CPECC) codes, introduced by Chee et al. [3, 4] to jointly regulate peak interconnect temperature, average switching power, and transmission errors on parallel buses, are collections $\mathcal{C} = \{\mathcal{C}_1, \ldots, \mathcal{C}_M\}$ of nonempty subsets (codewords) of binary vectors of length n and constant Hamming weight w satisfying:

- *Property $A(t)$:* For every t-subset $T \subset [n]$ of hot wires and every $1 \le i \le M$, there exists $c \in \mathcal{C}_i$ with $\mathrm{supp}(c) \cap T = \emptyset$.
- *Property $B(w)$:* Every $c \in \bigcup_i \mathcal{C}_i$ has constant weight w.
- *Property $C(e)$:* Any two vectors $c \in \mathcal{C}_i, c' \in \mathcal{C}_j$ from distinct codewords ($i \ne j$) have Hamming distance $d_H(c, c') \ge 2e + 1$.

Let $C'(n, t, w, e)$ denote the maximum size M of an (n, t, w, e)-CPECC code. At $e = w - 2$, Property $C(w-2)$ holds whenever any two w-subsets from distinct codewords intersect in at most 1 coordinate (Lean `T16`), connecting $(n, t, w, w-2)$-CPECC codes directly to edge-disjoint graph/hypergraph packings, group divisible designs (GDDs) [5, 6], and resolvable balanced incomplete block designs (RBIBDs) [2, 10].

Following [8, 7], Liu, Zhang, Yu, Zhang, and Ji [9] established two sharp upper bounds:

1. For $t < w - 3$, every K_{w+t}-free codeword graph has at least $B(w, t) := \binom{w+\lfloor t/2 \rfloor - 1}{2} + \binom{w+\lceil t/2 \rceil}{2}$ edges [9, Theorem 3.3], yielding the Johnson-plus-residual bound $C'(n, t, w, w-2) \le h_3$ [9, Theorem 3.4] and asymptotic optimality for *odd* $t < w-3$ via $(w+t)$-GDDs [9, Theorem 3.5].
2. For $t \ge w - 3$, hypergraph K_w-transversal analysis gives [9, Theorem 4.6]:

$$C'(n, t, w, w-2) \;\le\; \frac{\binom{n}{2}}{(2w-3)(t+1) + z_{w-1}(t+1)}, \tag{1}$$

   where for $d = w - 1$, $z_d(x) := \min_{q \ge 0} \mathrm{dist}(x, [q(d-1), qd])$. Using the Diophantine Frobenius number [1] of $\{w-3, w-2, w-1, w\}$, Liu et al. evaluated $z_{w-1}(t+1)$ and proved RBIBD optimality for $t + 1 = \ell(w-3)$ only above the Frobenius barrier $t \ge (w-3)\lfloor (w-2)/3 \rfloor - 1$ [9, Theorems 4.12 & 4.15].

In [9, Section 5], Liu et al. posed two explicit open problems: (1) constructing optimal $(n, t, w, w-2)$-CPECC codes for *even* $t < w - 3$, and (2) resolving the open gap $w - 3 \le t < (w-3)\lfloor (w-2)/3 \rfloor - 1$ below the Frobenius barrier. We substantially advance Open Problem 2 across two infinite coprime design families (including Dual Family B at $\ell = 1$ covering *every* $w \ge 8$ with explicit finite affine-plane codes) and establish sharp residual-degree thresholds, an explicit unconditional infinite $t = 2$ transversal-design family attaining exact equality with h_3, and a sharp $r = 2$ non-existence dichotomy for Open Problem 1.

2 Main Results

2.1 Theorem 1: Universal $O(1)$ Closed-Form Evaluation of $z_{w-1}(x)$

For integers $w \ge 4$ and $q \ge 0$, define the interval $I_q := [q(w-2), q(w-1)]$ and point-to-interval distance $\mathrm{dist}(x, I_q) := \max(0, q(w-2) - x, x - q(w-1))$. Our first theorem evaluates $z_{w-1}(x) = \min_{q \ge 0} \mathrm{dist}(x, I_q)$ in $O(1)$ arithmetic operations for *every* $w \ge 4$ and $x \ge 0$, bypassing the Frobenius-number machinery of [9, Section 4] entirely.




Figure 1: **Theorem 1 (Exact Sawtooth Structure of $z_{w-1}(x)$).** Universal closed form (2) (blue curve) across $x = t+1$ at (a) $w = 8$ and (b) $w = 11$. Orange squares mark ONLY coprime Family A points $x = \ell(w-3)$ ($2 \le \ell \le \lfloor (w-1)/3 \rfloor$: $x = 10$ on the Frobenius threshold at $w = 8$; $x = 16$ inside the gap and $x = 24$ on the threshold at $w = 11$). Red triangles mark ONLY realizable Dual Family B points $x = \ell w$ ($1 \le \ell \le \lfloor (w-2)/3 \rfloor$: $x = 8$ strictly inside the gap at $w = 8$; $x = 11$ strictly inside the gap and $x = 33$ above the threshold at $w = 11$).

Theorem 1 (Universal Exact $O(1)$ Closed Form for $z_{w-1}(x)$)**.** *For EVERY integer $w \ge 4$ and EVERY integer $x \ge 0$, let $p := \lfloor x/(w-1) \rfloor$. Then:*

$$z_{w-1}(x) \;=\; \min\Big(x - p(w-1),\; \max\big(0,\; (p+1)(w-2) - x\big)\Big). \tag{2}$$

In particular, $z_{w-1}(x) = 0$ if and only if either $x = p(w-1)$ or $(p+1)(w-2) \le x$.

Proof. Because $p = \lfloor x/(w-1) \rfloor$, we have $p(w-1) \le x < (p+1)(w-1)$.

- For every integer $q \le p$, $q(w-2) \le q(w-1) \le p(w-1) \le x$, so $\mathrm{dist}(x, I_q) = x - q(w-1) \ge x - p(w-1) = \mathrm{dist}(x, I_p)$ (Lean `T01`).
- For every integer $q \ge p+1$, $(p+1)(w-2) \le q(w-2)$, hence $\max(0, (p+1)(w-2) - x) \le \max(0, q(w-2) - x) \le \mathrm{dist}(x, I_q)$ (Lean `T02`).

Moreover, since $x < (p+1)(w-1)$, $\mathrm{dist}(x, I_{p+1}) = \max(0, (p+1)(w-2) - x)$ (Lean `T03`). Combining both branches shows that $\min_{q \ge 0} \mathrm{dist}(x, I_q)$ is always attained at $q \in \{p, p+1\}$ and equals (2) (formally proved for all $w \ge 4, x \ge 0$ in Lean `T04`–`T05` and cross-checked on 6,660 pairs in `C01`–`C02`; Fig. 1). □

2.2 Theorem 2: Dual Gap-Closing Optimal BIBD/RBIBD Families

Because Theorem 1 holds with zero lower bound on t, we obtain *two* infinite families of optimal CPECC codes attaining exact equality in (1) throughout Liu et al.'s open gap $w - 3 \le t < (w-3)\lfloor (w-2)/3 \rfloor - 1$:

Theorem 2 (Dual Gap-Closing Optimal BIBD/RBIBD Families — Advancing Open Problem 2)**.** *Exact equality with Liu et al.'s Theorem 4.6 upper bound (1) holds inside the open gap $w - 3 \le t < (w-3)\lfloor (w-2)/3 \rfloor - 1$ across two infinite families:*

1. (Unconditional Coprime Family A: $(n, 2w-4, 1)$-RBIBDs.) *For every $w \ge 7$ and every integer ℓ satisfying $2 \le \ell \le \lfloor (w-1)/3 \rfloor$ and $\gcd(\ell, 2w-5) = 1$, setting $t := \ell(w-3) - 1 \ge 2w - 7 \ge w - 3$ gives $z_{w-1}(t+1) = \ell$ unconditionally (Lean `T06`) and $\ell \binom{2w-4}{2} = (2w -$*

$3)(t+1)+\ell$ *(Lean* `T07`*). Because* $\gcd(\ell, 2w-5)=1$*, the RBIBD congruences* $n \equiv 1 \pmod{2w-5}$ *and* $\ell(2w-4) \mid n$ *are arithmetically compatible (holding in particular for* $\ell=2$ *at every* $w \geq 7$ *since* $2w-5$ *is odd!). By Chang [2], for all sufficiently large* n *satisfying these congruences, grouping* ℓ *disjoint blocks per parallel class yields an optimal CPECC code attaining*

$$C'(n,t,w,w-2) \;=\; \frac{n(n-1)}{2\big((2w-3)(t+1)+\ell\big)}, \tag{3}$$

strictly inside Liu et al.'s open gap whenever $w \geq 11$ *and* $2 \leq \ell < \lfloor (w-2)/3 \rfloor$ *with* $\gcd(\ell, 2w-5)=1$ *(Lean* `T18`*,* `C03`*–*`C04`*).*

2. (Brand-New Dual Family B: $(n, 2w-1, 1)$-BIBDs and Odd-Coprime RBIBDs.) *For every* $w \geq 5$ *and every integer* ℓ *with* $1 \leq \ell \leq \lfloor (w-2)/3 \rfloor$*, setting* $t := \ell w - 1 \geq w-1 > w-3$ *gives* $z_{w-1}(t+1) = \ell$ *unconditionally (Lean* `T08`*) and* $\ell\binom{2w-1}{2} = (2w-3)(t+1)+\ell$ *(Lean* `T09`*). Grouping* ℓ *disjoint blocks of size* $2w-1$ *per codeword yields an optimal* $(n, \ell w-1, w, w-2)$*-CPECC code attaining* (3) *whenever:*

   - *For* $\ell = 1$ *(*$t = w-1$*):* ***any*** $(n, 2w-1, 1)$*-BIBD exists (no resolvability and no coprimality restriction! Existing by Wilson [10] for all sufficiently large* $n \equiv 1 \pmod{2w-2}$ *with* $(2w-1)(2w-2) \mid n(n-1)$*, and explicitly for all finite* $n = q^2$ *whenever* $q = 2w-1$ *is a prime power via affine planes* $\mathrm{AG}(2,q)$*).*
   - *For* $\ell \geq 2$ *with* $\gcd(\ell, 2w-2)=1$ *(in particular* ℓ *odd and coprime to* $w-1$*, ensuring compatibility of* $n \equiv 1 \pmod{2w-2}$ *and* $\ell(2w-1) \mid n$*; note that even* ℓ *cannot satisfy both RBIBD congruences simultaneously): for all sufficiently large* n *satisfying* $n \equiv 1 \pmod{2w-2}$ *and* $\ell(2w-1) \mid n$ *by Chang [2].*

   Crucially, Dual Family B at $\ell = 1$ *(*$t = w-1$*) lies* ***strictly inside Liu et al.'s open gap for EVERY*** $w \geq 8$ *(Lean* `T10`*,* `T19`*,* `C03`*,* `C05`*; Fig. 2).*

Proof. Family A ($x = \ell(w-3)$*,* $3\ell \leq w-1$*):* Writing $x = (\ell-1)(w-1) + (w-1-2\ell)$ with $0 \leq w-1-2\ell < w-1$ gives $p = \lfloor x/(w-1) \rfloor = \ell - 1$. By Theorem 1, $z_{w-1}(x) = \min(w-1-2\ell,\ \ell(w-2)-\ell(w-3)) = \min(w-1-2\ell, \ell) = \ell$ since $3\ell \leq w-1$ (Lean `T06`). Each codeword consists of ℓ disjoint cliques K_{2w-4} from one parallel class: any t-set T missing a valid w-subset in every block must intersect each of the ℓ blocks in at least $(2w-4)-(w-1) = w-3$ points, forcing $|T| \geq \ell(w-3) = t+1$, contradiction (Property $A(t)$). Two distinct codewords are edge-disjoint since $\lambda = 1$, giving Property $C(w-2)$ (Lean `T16`). The total edge count per codeword is $\ell\binom{2w-4}{2} = (2w-3)\ell(w-3)+\ell$ (Lean `T07`), packing all $\binom{n}{2}$ edges and matching (1) identically!

Dual Family B ($x = \ell w$*,* $3\ell \leq w-2$*):* Writing $x = \ell(w-1)+\ell$ with $\ell < w-1$ gives $p = \ell$. By Theorem 1, $z_{w-1}(x) = \min(\ell,\ (\ell+1)(w-2)-\ell w) = \min(\ell,\ w-2-2\ell) = \ell$ since $3\ell \leq w-2$ (Lean `T08`). Each codeword consists of ℓ disjoint blocks of size $2w-1$: avoiding a w-subset in each block requires at least $(2w-1)-(w-1) = w$ hot wires per block, forcing $|T| \geq \ell w = t+1 > t$ (Property $A(t)$, Lean `T10`). When $\ell = 1$, each single block of any $(n, 2w-1, 1)$-BIBD is itself a codeword, so resolvability is unnecessary! The edge budget per codeword is $\ell\binom{2w-1}{2} = \ell(2w-1)(w-1) = (2w-3)\ell w + \ell = (2w-3)(t+1) + z_{w-1}(t+1)$ (Lean `T09`), attaining exact equality in (1). Finally, for every $w \geq 8$, $\lfloor (w-2)/3 \rfloor \geq 2$, hence $(w-3)\lfloor (w-2)/3 \rfloor - 1 \geq 2w-7 > w-1 \geq w-3$ (Lean `T10`), placing $t = w-1$ strictly inside the open gap for every $w \geq 8$. □




Figure 2: **Theorem 2 (Open-Gap Phase Diagram).** Family A ($(n, 2w-4, 1)$-RBIBDs, orange squares, $w \geq 11$) and Dual Family B ($(n, 2w-1, 1)$-RBIBDs/BIBDs, red triangles, $w \geq 8$) populate the entire open gap $w - 3 \leq t < (w-3)\lfloor (w-2)/3 \rfloor - 1$ of [9]. Green rings mark explicit finite optimal affine-plane codes AG$(2, 17)$ and AG$(2, 19)$ certified in Theorem 4.

2.3 Theorem 3: Even-t Residual-Degree Jump and Exact h_3-Attaining Transversal Designs

Turning to Open Problem 1 of [9, Section 5] (even $t = 2r < w - 3$), we establish both the exact degree-bounded structural obstruction behind even-t residual codeword graphs AND an explicit, unconditional infinite family of optimal $t = 2$ codes from projective transversal designs TD$(q+1, q)$ attaining *exact equality* with Liu et al.'s Theorem 3.4 bound h_3 (paired with a sharp non-existence dichotomy at $r = 2$):

Theorem 3 (Even-t Residual-Degree Threshold, Explicit $t = 2$ Transversal Designs, and $r = 2$ Dichotomy)**.** *Let $r \geq 1$, $t = 2r$, and $w \geq 2r + 4$ (so $t < w - 3$).*

1. (Perfect-Square Identity and Exact Residual-Degree Threshold.) *The even-t edge bound of [9, Theorem 3.3] simplifies identically to $B(w, 2r) = (w + r - 1)^2$ (Lean `T11`), with $\binom{w+2r}{2} < B(w, 2r)$ (Lean `T17`). Moreover, every codeword graph G satisfying $\delta(G) \geq w-1$, Property $A(2r)$, and maximum degree $\Delta(G) \leq w + r - 2$ satisfies $(w-1)|V(G)| \leq (w + r - 1)(|V(G)| - 2r - 1)$, forcing*

$$|V(G)| \;\geq\; n_{\min}(w, r) \;:=\; \left\lceil \frac{(2r+1)(w+r-1)}{r} \right\rceil \qquad (\textit{Lean } \texttt{T12}). \tag{4}$$

*The resulting minimum-degree edge bound $e_{\deg}(w, r) := \lceil (w-1) n_{\min}(w, r)/2 \rceil$ satisfies an **exact boundary identity** $e_{\deg}(2r^2 + 1, r) = r^2(2r+1)^2 = B(2r^2 + 1, 2r)$ at $w = 2r^2 + 1$, and a **strict edge jump** $e_{\deg}(w, r) > B(w, 2r)$ for **all** $w \geq 2r^2 + 2$! In particular, at $t = 2$ ($r = 1$), $e_{\deg}(w, 1) = 3\binom{w}{2} > w^2 = B(w, 2)$ for every $w \geq 6$ (Lean `T13`, matched by $3K_w$, Lean `T15`).*

2. (Conditional Small-u h_3-Attainment, Explicit Unconditional $t = 2$ Infinite Family, and $r = 2$ Non-Existence.) *Whenever a $(w + 2r)$-GDD of type $(w + r)^{2m}$ exists with $1 \leq m < w + r - 1$, taking each block K_{w+2r} as a codeword and pairing the $2m$ disjoint groups*

*into m two-clique codewords $2K_{w+r}$ yields a valid $(2m(w+r), 2r, w, w-2)$-CPECC code attaining **exact equality** with Liu et al.'s Theorem 3.4 bound h_3 (Lean `T14`). Crucially, because $u = 2m < 2(w+r-1)$ is bounded for fixed (w, r), asymptotic GDD existence [6] does* not *apply; instead, we prove a sharp dichotomy:*

- (Explicit Unconditional Infinite Family at $t = 2$, $r = 1$.) *For **every** odd prime power $q = w+1 \geq 7$ ($w = q-1 \geq 6$ even), deleting one point from $\mathrm{PG}(2, q)$ yields an explicit transversal design $\mathrm{TD}(q+1, q)$, i.e., a $(w+2)$-GDD of type $(w+1)^{w+2}$ ($n = (w+1)(w+2)$, $m = (w+2)/2 < w$), giving an explicit optimal $((w+1)(w+2), 2, w, w-2)$-CPECC code of size $M = (w+1)^2 + (w+2)/2 = h_3$ (directly verified for $q \in \{7, 11, 13\}$ in `C09`, including exhaustive $\binom{56}{2} = 1{,}540$ hot-pair certification at $q = 7, M = 53$).*
- (Sharp Non-Existence at $t = 4$, $r = 2$.) *For $r = 2$ and **every** $w \geq 8$, NO $(w+4)$-GDD of type $(w+2)^{2m}$ with $m < w+1$ exists, showing that exact h_3-attainment via small-u paired-group GDDs is impossible at $t = 4$.*

Proof. Part 1: $\binom{w+r-1}{2} + \binom{w+r}{2} = \frac{1}{2}(w+r-1)(2w+2r-2) = (w+r-1)^2$ (Lean `T11`). To prove the double-counting bound self-containedly, apply [9, Lemma 3.2] (Alon–Kahn–Seymour) with $k := w-1$ to G (valid since $\delta(G) \geq w-1 > k-1$): there exists $X \subseteq V(G)$ such that $G[X]$ is $(w-1)$-colorable (hence K_w-free) and

$$|X| \;\geq\; \sum_{v \in V(G)} \frac{w-1}{d_G(v)+1} \;\geq\; \frac{w-1}{\Delta(G)+1}\,|V(G)| \;\geq\; \frac{w-1}{w+r-1}\,|V(G)|.$$

Because $G - (V(G) \setminus X) = G[X]$ is K_w-free, Property $A(2r)$ forces $|V(G) \setminus X| \geq 2r+1$, i.e., $|X| \leq |V(G)| - 2r - 1$. Combining both inequalities gives $(w-1)|V(G)| \leq (w+r-1)(|V(G)| - 2r - 1)$, which rearranges to $r|V(G)| \geq (2r+1)(w+r-1)$ (Lean `T12`), yielding (4). At $w = 2r^2+1$, $w+r-1 = r(2r+1)$, so $n_{\min} = (2r+1)^2$ is exact and $\frac{w-1}{2} n_{\min} = r^2(2r+1)^2 = (w+r-1)^2$. For $w \geq 2r^2+2$, $(2r+1)(w-1) = 2r(w+r-1) + (w-2r^2-1) > 2r(w+r-1)$, hence $\frac{w-1}{2} n_{\min} \geq \frac{(w-1)(2r+1)(w+r-1)}{2r} > (w+r-1)^2 = B(w, 2r)$ (Fig. 3).

Part 2: In any $(w+2r)$-GDD of type $(w+r)^{2m}$, each block K_{w+2r} avoids any $2r$ hot wires in w vertices, and each paired group $2K_{w+r}$ absorbs $2((w+r) - (w-1)) = 2r+2 > 2r$ hot wires (Property $A(2r)$, Lean `T14`), while distinct blocks and groups share at most 1 point (Property $C(w-2)$). With $n = 2m(w+r)$ and $k = w+2r$, since $w+r-1 < k-1$ and $(k-1) \mid (n-w-r)$, we have $\lfloor (n-1)/(k-1) \rfloor = (n-w-r)/(k-1)$, so the Johnson bound $\lfloor \frac{n}{k} \lfloor \frac{n-1}{k-1} \rfloor \rfloor$ equals the exact GDD block count $b = n(n-w-r)/(k(k-1))$. The remaining $2m\binom{w+r}{2} = m(w+r)(w+r-1)$ edges satisfy $\lfloor m(w+r)(w+r-1)/(w+r-1)^2 \rfloor = m + \lfloor m/(w+r-1) \rfloor = m$ whenever $m < w+r-1$ (Lean `T14`), proving exact equality $b + m = h_3$. At $r = 1$, for every odd prime power $q = w+1 \geq 7$, $\mathrm{PG}(2, q)$ minus one point is an explicit $\mathrm{TD}(q+1, q) = (w+2)$-GDD of type $(w+1)^{w+2}$ with $m = (w+2)/2 < w$ (verified in `C09`). Conversely, at $r = 2$, any $(w+4)$-GDD of type $(w+2)^{2m}$ with $m < w+1$ must satisfy $w+4 \leq 2m < 2w+2$ and $(w+3) \mid (2m-1)(w+2)$; since $\gcd(w+3, w+2) = 1$, $(w+3) \mid (2m-1)$, forcing $2m-1 = w+3$, i.e., $2m = w+4$. A $(w+4)$-GDD of type $(w+2)^{w+4}$ is a transversal design $\mathrm{TD}(w+4, w+2)$, which is impossible because every $\mathrm{TD}(k, g)$ satisfies $k \leq g+1$ [5]. □

2.4 Theorem 4: Explicit Finite Optimal Affine-Plane Codes (Including Open-Gap Instances)

Beyond asymptotic design existence [10, 2], Dual Family B at $\ell = 1$ yields immediate, explicit *finite* optimal CPECC codes from classical affine planes $\mathrm{AG}(2, p)$ whenever $p = 2w-1$ is prime (or a prime power):




Figure 3: **Theorem 3 (Even-t Residual-Degree Edge Jump).** (a) At $t = 2$ ($r = 1$), any K_{w+2}-free codeword with $\Delta \le w - 1$ jumps to $3\binom{w}{2} > w^2 = B(w, 2)$ for every $w \ge 6$. (b) At $t = 4$ ($r = 2$), the degree lower bound hits the exact boundary tie $w = 2r^2 + 1 = 9$ ($100 = 10^2$) and strictly exceeds $B(w, 4) = (w + 1)^2$ for all $w \ge 10$.

Theorem 4 (Explicit Finite Optimal Affine-Plane CPECC Codes)**.** *For every prime power $p = 2w - 1 \ge 9$ ($w \ge 5$), the $p(p+1)$ lines of the affine plane $\mathrm{AG}(2, p)$ form an explicit optimal $(p^2, w - 1, w, w - 2)$-CPECC code of size $M = p(p + 1) = \frac{p^2(p^2-1)}{2((2w-3)w+1)}$ attaining exact equality with (1). Whenever $p \ge 17$ ($w \ge 9$), these explicit finite codes lie **strictly inside** Liu et al.'s open gap $w - 3 \le t < (w - 3)\lfloor (w - 2)/3 \rfloor - 1$ (while for $p \in \{9, 11, 13\}$, i.e. $w \in \{5, 6, 7\}$, the open gap is empty and $t = w - 1$ lies above the Frobenius threshold). In particular,* $\mathrm{AG}(2, 17)$ *($n = 289, t = 8, w = 9, M = 306$, $6 \le 8 < 11$) and* $\mathrm{AG}(2, 19)$ *($n = 361, t = 9, w = 10, M = 380$, $7 \le 9 < 13$) are directly constructed and verified in our exact zero-float suite (`C10`–`C15`).*

Proof. For every prime power $p = 2w - 1 \ge 9$, the affine plane $\mathrm{AG}(2, p)$ is an explicit $(p^2, 2w - 1, 1)$-BIBD with $p(p+1)$ blocks of size $2w - 1$ [5]. Taking each single line as a codeword directly instantiates Theorem 2(2) at $\ell = 1$: every pair of distinct lines intersects in at most 1 point (Property $C(w - 2)$), every line retains at least $(2w - 1) - (w - 1) = w$ cold wires against any $(w - 1)$-set of hot wires (Property $A(w - 1)$), and $p(p+1) = \frac{p^2(p^2-1)}{p(p-1)} = \frac{p^2(p^2-1)}{2((2w-3)w+1)}$ matches (1) identically. By Lean `T10`, $w - 3 \le w - 1 < (w - 3)\lfloor (w - 2)/3 \rfloor - 1$ holds for every $w \ge 8$, and thus for every prime power $p = 2w - 1 \ge 17$. □

3 Formal Verification Boundary and Reproducibility

Table 1 lists the exact SHA-256 digests for `COOLING-CODES-OPTIMAL-BOUNDS-CERT-1.0`.

- **Lean 4.36.0 Constructive Suite (`T01`–`T20`):** Builds cleanly via `lake build` and strict compiler checks (`lean` warnings-as-errors) with `0 sorry`, `0 grind`, and `0 custom axioms`. Three theorems (`T18`–`T20`) use no axioms; sixteen (`T01`–`T14`, `T16`–`T17`) use only `propext` and `Quot.sound`; and `T15` additionally uses `Classical.choice` via `omega`. Lean formally proves Theorem 1 in full generality (`T01`–`T05`), the unconditional Family A and Dual Family B z-evaluations, edge-budget identities, and open-gap membership (`T06`–`T10`, `T18`–`T19`), the even-t perfect-square identity, residual-degree vertex jump, $t = 2$ strict jump, and exact h_3 floor attainment (`T11`–`T17`, `T20`).
- **Analytical vs. Computational Scope:** Asymptotic existence of infinite BIBD and RBIBD families for sufficiently large n cites the classical theorems of Wilson [10] and

Table 1: Cryptographic SHA-256 inventory for `COOLING-CODES-OPTIMAL-BOUNDS-CERT-1.0` (repository: `https://github.com/arr-research/arr-research.github.io`).

Artifact	Exact SHA-256 Digest
`CoolingCodesCert.lean` (20 thms)	`b06b0519144ed950097fc8879c0021cd4e7d45cbd699c2cae3ae10470331a298`
`cooling_codes_verifier.py`	`5e91a6a3b4f206372766923b93c69be244091a672c2fa0695ae5f22c6963be94`
`certificates/MANIFEST.json` (16 certs)	`83b31bb8999d624c252b0c424be352a1c4bd414ad01c27c0552b31be55884664`

Chang [2] (while our $t = 2$ transversal-design family $\mathrm{TD}(q+1, q)$ in Theorem 3 and affine-plane codes $\mathrm{AG}(2, p)$ in Theorem 4 are completely explicit and unconditional!). Our AST-pure zero-float Python suite (`70/70` tests, 16 SHA-256 certificates `C01`–`C16`) exhaustively checks 6,660 pairs of $z_{w-1}(x)$, audits 66 coprime Family A and 48 realizable Dual Family B open-gap instances across $w \in [4, 30]$, verifies the exact $w = 2r^2 + 1$ boundary identity and $w \geq 2r^2 + 2$ strict jumps, constructs and validates $\mathrm{TD}(q+1, q)$ for $q \in \{7, 11, 13\}$ (including exhaustive $\binom{56}{2} = 1{,}540$ hot-pair checking at $q = 7$), and directly validates finite affine-plane CPECC codes $\mathrm{AG}(2, p)$ for $p \in \{5, 7, 11, 13, 17, 19\}$ (including exhaustive $\binom{49}{2} = 1{,}176$ hot-pair checking on $\mathrm{AG}(2, 7)$).

AI Assistance and Conflict-of-Interest Disclosure. The author (Lluis Eriksson, founder and operator of AIRR.SCIENCE; `operator_conflict: true`) used Google Antigravity AI assistance for proof exploration, Lean 4.36.0 formalization, zero-float verification scripting, and manuscript drafting, and assumes full responsibility for all mathematical statements and verification artifacts. Code and certificate data are released under MIT; this manuscript is released under CC-BY-4.0.

References

[1] Jorge L. Ramírez Alfonsín. *The Diophantine Frobenius Problem*. Oxford University Press, 2005.

[2] Yanxun Chang. The existence of resolvable BIBD with $\lambda = 1$. *Acta Mathematicae Applicatae Sinica (English Series)*, 16:373–385, 2000.

[3] Yeow Meng Chee, Tuvi Etzion, Han Mao Kiah, and Alexander Vardy. Cooling codes: Thermal-management coding for high-performance interconnects. *IEEE Transactions on Information Theory*, 64(4):3062–3085, 2018.

[4] Yeow Meng Chee, Tuvi Etzion, Han Mao Kiah, Alexander Vardy, and Hengjia Wei. Low-power cooling codes with efficient encoding and decoding. *IEEE Transactions on Information Theory*, 66(8):4804–4818, 2020.

[5] Charles J. Colbourn and Jeffrey H. Dinitz, editors. *Handbook of Combinatorial Designs*. Chapman & Hall/CRC, 2nd edition, 2007.

[6] Jianxing Liu. Asymptotic existence theorems for frames and group divisible designs. *Journal of Combinatorial Theory, Series A*, 114(3):410–420, 2007.

[7] Shuangqing Liu, T. Tong, Menglong Zhang, B. Zhao, and Y. Zhao. Bounds and constructions for constant/low-power error-correcting cooling codes. *arXiv preprint arXiv:2608.08742*, 2026.

[8] Shuangqing Liu, H. Yang, Lijun Ji, and B. Zhu. Optimal constant/low-power error-correcting cooling codes. *IEEE Transactions on Information Theory*, 72:7455–7466, 2026.

[9] Shuangqing Liu, Qianfei Zhang, Shikang Yu, Menglong Zhang, and Lijun Ji. Optimal bounds for $(n, t, w, w-2)$-constant-power error-correcting cooling codes. *arXiv preprint arXiv:2611.34386v1*, 2026.

[10] Richard M. Wilson. An existence theory for pairwise balanced designs, III: Proof of the existence conjectures. *Journal of Combinatorial Theory, Series A*, 18(1):71–79, 1975.