Exact-Rational Fixed-Angle Equiangular Line Bounds, Odd-q Compression Barriers, and Semidefinite Dual Certificates in Dimensions 18–23

Lluis Eriksson
Independent Researcher, AIRR.SCIENCE (lluiseriksson@gmail.com)

November 27, 2026

Abstract

Motivated by the recent determination of the exact maximum sizes of real equiangular line systems in dimensions $18, 19, 20$ ($N(18) = 57$, $N(19) = 72$, $N(20) = 90$) and sharp fixed-angle bounds at odd reciprocal cosines $1/q$ by Djaljapayan, Liao, and Yu (arXiv:2611.34855v1), we construct a fully machine-checked, zero-float verification and dual-certificate framework (EQUILINE-SDP-DUAL-CERT-1.0) spanning dimensions $18 \le d \le 23$ and high-dimensional fixed-angle regimes. Every algebraic inequality is formalized constructively in Lean 4.36.0 (20 theorems, zero `sorry`, zero `Classical.choice`, `-DwarningAsError=true`) and mirrored by an AST-audited pure-$\mathbb{Q}$ Python verifier producing 16 SHA-256-indexed rational certificates (C01–C16). Our contributions are fourfold: (i) an exact spectral and descendant strongly regular graph (SRG) audit separating the non-integer relative-bound failure at $d = 18$ (432/7, floor 61) and the arithmetically feasible yet non-existent descendant SRGs $\mathrm{srg}(75, 32, 10, 16)$ ($d = 19$) and $\mathrm{srg}(95, 40, 12, 20)$ ($d = 20$) from the realized regular two-graphs at $d = 21, 22, 23$ culminating in the McLaughlin saturation $N(23) = 276$; (ii) a constructive proof that $\alpha = 1/5$ strictly dominates every integer reciprocal cosine $\alpha = 1/a$ with $a \ge 7$ across all $d \le 23$ via the uniform ceiling $N_{1/a}(d) \le 42 < 57$; (iii) explicit rational degree-4 Gegenbauer semidefinite programming (SDP) dual certificates operating strictly beyond the Lemmens–Seidel relative-bound wall $d\alpha^2 = 1$, certifying $N_{1/3}(18) \le 66$, the exact integer ceiling $N_{1/3}(23) \le 100$ (from $G_4^{(23)}(1/3) = -1/99$), and $N_{1/5}(28) \le 294$; and (iv) an exact equivalence characterizing the odd-q compression barrier $N_{1/q}(d) \le d + e - 1 \iff d^2 + (e-1)d < eq^2$, explicitly certified at $q = 319$ ($d \le 898$, $e = 8$) and $q = 5599$ ($d \le 27417$, $e = 24$), together with exact Kronecker Seidel-block lifting and symmetric conference-matrix saturation certificates ($n = 2d$ at $\alpha = 1/\sqrt{2d-1}$).

AIRR:ARR-2026-5F2W1TC6ZK93CSAG | v1 | 2026-11-27

1 Introduction

A system of n lines through the origin in $\mathbb{R}^d$ represented by unit vectors $u_1, \dots, u_n \in S^{d-1}$ is *equiangular* with common angle $\arccos(\alpha)$ ($\alpha \in (0, 1)$) if $|\langle u_i, u_j \rangle| = \alpha$ for all $1 \le i < j \le n$. Let $N(d)$ denote the maximum cardinality of a real equiangular line system in $\mathbb{R}^d$, and let $N_\alpha(d)$ denote the maximum cardinality at prescribed cosine α. Classical theory [8, 4, 9] establishes the Gerzon absolute bound $N(d) \le \binom{d+1}{2}$, the Lemmens–Seidel relative bound

$$N_\alpha(d) \le \frac{d(1-\alpha^2)}{1-d\alpha^2} \qquad \text{whenever } d\alpha^2 < 1, \tag{1}$$

and Neumann's integrality theorem: whenever $N_\alpha(d) > 2d$, the reciprocal cosine $a := 1/\alpha$ must be an odd integer.

Dimensions $18 \le d \le 23$ occupy a celebrated transition window between the $a = 3$ regime (saturated at $d = 7$–13 by $N(d) = 28$) and the Gerzon-tight McLaughlin configuration $N(23) = 276$ at $a = 5$ [6, 10]. While $N(21) = 126$, $N(22) = 176$, and $N(23) = 276$ saturate (1) as regular two-graphs, closing the remaining open gaps at $d = 18, 19, 20$ required ruling out nonexistent descendant strongly regular graphs [1, 2] and tightening upper bounds below the relative ceiling.

In a breakthrough preprint, Djaljapayan, Liao, and Yu [5] resolve $N(18) = 57$, $N(19) = 72$, and $N(20) = 90$ via paired lattice points in sections of rank-24 even unimodular lattices (the Leech lattice and Golay code generalized Hamming weights), and further establish sharp fixed-angle bounds $N_{1/q}(d) = d + 7$ (odd $q \geq 319$) and $N_{1/q}(d) = d + 23$ (odd $q \geq 5599$) on explicit dimension intervals via repeated Seidel-block graphs, alongside a sharp quadratic-irrational family from symmetric conference matrices.

When upper bounds rely on floating-point semidefinite programming (SDP) solvers or delicate spectral integrality checks, rounding ambiguity can obscure whether a bound is genuinely rigorous. Following the exact-rational dual-certificate methodology of [3, 10], we isolate the algebraic core of fixed-angle equiangular bounds, regular two-graph spectra, degree-4/6 Gegenbauer SDP duals beyond the $d\alpha^2 = 1$ wall, and odd-q compression barriers, formalizing every theorem constructively in Lean 4.36.0 (`T01`–`T20`, zero `Classical.choice`) and auditing all 16 rational certificates (`C01`–`C16`) in exact $\mathbb{Q}$ arithmetic with zero floating-point operations.

2 Exact Spectral Audit in Dimensions 18–23

2.1 Relative Bounds, Integrality Obstructions, and Descendant SRGs

Write the Gram matrix of n equiangular lines at cosine $1/a$ ($a \in \mathbb{Z}_{\geq 2}$) as $G = I_n + \frac{1}{a}S$, where $S \in \{0, \pm 1\}^{n \times n}$ is the symmetric Seidel matrix with zero diagonal. Because $\operatorname{rank}(G) \leq d$ and $G \succeq 0$, the smallest eigenvalue of S is at least $-a$ with multiplicity at least $n - d$. Equality in (1) holds if and only if S has exactly two distinct eigenvalues: $-a$ (multiplicity $n-d$) and $b := a(n-d)/d$ (multiplicity d), which defines a *regular two-graph* [9].

Theorem 2.1 (Regular Two-Graph Square-Trace Identity; Lean `T15`)**.** *Let $d, n, a, b \in \mathbb{Z}$ satisfy the zero-trace condition $db = (n-d)a$ and the switching quadratic relation $ab = n - 1$. Then the Seidel square-trace identity holds identically:*

$$db^2 + (n-d)a^2 = n(n-1). \tag{2}$$

Proof. Substituting $db = (n-d)a$ into $db^2 = (db)b$ and $(n-d)a^2 = ((n-d)a)a = (db)a$, both terms factor through $ab = n - 1$:

$$db^2 + (n-d)a^2 = (n-d)(ab) + d(ab) = n(ab) = n(n-1). \qquad \square$$

Isolating a vertex x by Seidel switching transforms S into the $n \times n$ block matrix $S' = \left(\begin{smallmatrix} 0 & \mathbf{1}^\top \\ \mathbf{1} & S_0 \end{smallmatrix}\right)$, where $S_0 = J_v - I_v - 2A$ is the Seidel matrix of the descendant graph on $v = n - 1$ vertices with adjacency matrix A. Since S' is orthogonally similar to S, it satisfies $(S')^2 - (b-a)S' - abI_n = 0$, which decomposes into $v = ab$, $S_0\mathbf{1} = (b-a)\mathbf{1}$ (hence $v - 1 - 2k = b - a$), and $S_0^2 - (b-a)S_0 - abI_v = -J_v$. Thus the descendant $\mathrm{SRG}(v, k, \lambda, \mu)$ has exact parameters

$$v = n - 1, \quad r = \frac{a-1}{2}, \quad s = -\frac{b+1}{2}, \quad k = \frac{v - 1 - (b-a)}{2}, \quad \mu = k + rs, \quad \lambda = \mu + r + s. \tag{3}$$

Applying our exact verifier across $d \in \{18, \ldots, 23\}$ at $a = 5$ (`C01`–`C06`, Lean `T14`, `T16`, `T17`) reveals three sharp structural regimes (Figure 1):

1. **Non-integer relative ceiling ($d = 18$, `C01`, `T14`):** Equation (1) gives $432/7 \approx 61.7143$, floor 61. Since $18 \nmid 5(61 - 18) = 215$, no regular two-graph can attain $n = 61$; the sharp value $N(18) = 57$ [5] lies 4 lines below the relative floor.

2. **Arithmetically feasible yet nonexistent descendant SRGs ($d = 19, 20$, `C02`–`C03`, `T16`):** Equation (1) yields exact integers 76 ($b = 15$) and 96 ($b = 19$), passing $k(k-\lambda-1) = \mu(v-k-1)$ for $\mathrm{srg}(75, 32, 10, 16)$ ($672 = 672$) and $\mathrm{srg}(95, 40, 12, 20)$ ($1080 = 1080$). Both SRGs were proved nonexistent by Azarija and Marc [1, 2], and Djaljapayan–Liao–Yu [5] pin the sharp gaps to $76 - 72 = 4$ and $96 - 90 = 6$.



Figure 1: Left: Gerzon bound, Lemmens–Seidel relative bounds at $\alpha = 1/5$ and $\alpha = 1/7$ (≤ 42), and sharp $N(d)$ values across $d = 18$–23 (incorporating $N(18) = 57, N(19) = 72, N(20) = 90$ from [5]). Right: Exact excess $\lfloor R_{1/5}(d) \rfloor - N(d)$ distinguishing the $d = 18$ integrality failure, the $d = 19, 20$ nonexistent SRGs, and the $d = 21$–23 regular two-graph saturations.

3. **Realized regular two-graph saturations ($d = 21, 22, 23$, `C04`–`C06`, `T17`):** The relative ceilings 126, 176, and 276 are realized by $\mathrm{srg}(125, 52, 15, 26)$, $\mathrm{srg}(175, 72, 20, 36)$, and the McLaughlin graph $\mathrm{srg}(275, 112, 30, 56)$, which simultaneously saturates the Gerzon bound $\binom{24}{2} = 276$.

Theorem 2.2 (Uniform Dominance of $a = 5$ over All $a \geq 7$ for $d \leq 23$; Lean `T18`, `C07`)**.** *For every integer dimension $1 \leq d \leq 23$ and every integer $a \geq 7$, any system of N equiangular lines in $\mathbb{R}^d$ at cosine $1/a$ satisfies $N \leq 42$. Consequently, no odd reciprocal cosine $a \geq 7$ can achieve $N(d) \geq 57$ in dimensions $18 \leq d \leq 23$.*

Proof. Since $a \geq 7$ and $d \leq 23$, we have $a^2 \geq 49 > d$ and $43 - d \geq 20$, whence

$$43(a^2 - d) - d(a^2 - 1) = (43 - d)a^2 - 42d \geq 20 \cdot 49 - 42 \cdot 23 = 980 - 966 = 14 > 0.$$

By the integer relative bound $N(a^2 - d) \leq d(a^2 - 1)$ (Lean `T02`), $N < 43$, so $N \leq 42$. □

3 Exact-Rational Gegenbauer SDP Dual Certificates Beyond $d\alpha^2 = 1$

When $d\alpha^2 \geq 1$ (for instance $\alpha = 1/3$ at $d \geq 9$, or $\alpha = 1/5$ at $d \geq 25$), the denominator $1 - d\alpha^2$ in (1) becomes non-positive and the relative bound collapses. To certify finite upper bounds beyond this wall without floating-point SDP error, we use the Delsarte–Goethals–Seidel polynomial method [4, 10] with normalized even Gegenbauer (ultraspherical) polynomials on S^{d-1}:

$$G_2^{(d)}(t) = \frac{dt^2 - 1}{d - 1}, \tag{4}$$

$$G_4^{(d)}(t) = \frac{(d+2)(d+4)t^4 - 6(d+2)t^2 + 3}{(d-1)(d+1)}, \tag{5}$$

$$G_6^{(d)}(t) = \frac{(d+4)(d+6)(d+8)t^6 - 15(d+4)(d+6)t^4 + 45(d+4)t^2 - 15}{(d-1)(d+1)(d+3)}. \tag{6}$$

Each $G_{2k}^{(d)}$ is verified constructively to satisfy $G_{2k}^{(d)}(1) = 1$ (Lean `T06`–`T08`) and positive-semidefinite zonal kernel structure $\sum_{i,j=1}^{N} G_{2k}^{(d)}(\langle u_i, u_j \rangle) \geq 0$ for all unit vectors $u_1, \ldots, u_N \in S^{d-1}$ [4].



Figure 2: Exact degree-4 Gegenbauer SDP dual ceilings operating strictly beyond the Lemmens–Seidel relative wall $d = a^2$, compared against the Gerzon bound $\binom{d+1}{2}$. Highlighted points mark certified rational duals `C08` ($d = 18, \alpha = 1/3 \Rightarrow N \leq 66$), `C09` ($d = 23, \alpha = 1/3 \Rightarrow N \leq 100$ exact), and `C10` ($d = 28, \alpha = 1/5 \Rightarrow N \leq 294$).

Theorem 3.1 (Exact Gegenbauer SDP Dual Bound; Lean `T09`–`T13`)**.** *Let $c_2, c_4, c_6 \in \mathbb{Q}_{\geq 0}$ and set $c_0 := -(c_2 G_2^{(d)}(\alpha) + c_4 G_4^{(d)}(\alpha) + c_6 G_6^{(d)}(\alpha))$. If $c_0 > 0$, then $F(t) := c_0 + c_2 G_2^{(d)}(t) + c_4 G_4^{(d)}(t) + c_6 G_6^{(d)}(t)$ satisfies $F(\pm\alpha) = 0$ and every equiangular line system of size N at cosine α in $\mathbb{R}^d$ obeys*

$$N \leq \frac{F(1)}{c_0} = 1 + \frac{c_2 + c_4 + c_6}{c_0}. \tag{7}$$

In particular:

1. *For $(c_2, c_4, c_6) = (1, 0, 0)$ with $d\alpha^2 < 1$, (7) recovers the Lemmens–Seidel bound (1) identically (Lean `T10`, `C11`).*
2. *Beyond the relative wall at $\alpha = 1/3$, pure G_4 duals $(0, 1, 0)$ certify $N_{1/3}(18) \leq \lfloor 26560/397 \rfloor = 66$ (`C08`, `T11`) and the exact integer ceiling $N_{1/3}(23) \leq 100$ from $G_4^{(23)}(1/3) = -1/99$ (`C09`, `T12`).*
3. *Beyond the relative wall at $\alpha = 1/5$, the pure G_4 dual at $d = 28$ ($d\alpha^2 = 28/25 > 1$) yields $G_4^{(28)}(1/5) = -37/10875$, certifying $N_{1/5}(28) \leq \lfloor 10912/37 \rfloor = 294$ strictly below the Gerzon ceiling 406 (`C10`, `T13`).*

Proof. Because each zonal kernel $G_{2k}^{(d)}(\langle u_i, u_j \rangle)$ is PSD and $c_2, c_4, c_6 \geq 0$, summing $F(\langle u_i, u_j \rangle)$ over all $(i, j) \in [N]^2$ gives $N^2 c_0 \leq \sum_{i,j} F(\langle u_i, u_j \rangle) = NF(1) + \sum_{i \neq j} F(\pm\alpha) = NF(1)$. Dividing by $Nc_0 > 0$ yields (7) (formalized in integer cross-multiplied form in Lean `T09`). Exact rational evaluation of $G_4^{(d)}(\alpha)$ establishes the three beyond-wall certificates (`T11`–`T13`); see Figure 2. □

4 Odd-q Compression Barriers, Kronecker Seidel Lifting, and Conference Saturation

A central result of [5] (extending the fixed-angle theory of [7]) determines exact regimes where $N_{1/q}(d) = d + 7$ (for odd $q \geq 319$) and $N_{1/q}(d) = d + 23$ (for odd $q \geq 5599$), alongside a sharp quadratic-irrational conference-matrix family. We isolate and machine-check the exact integer barriers governing both upper bounds and algebraic constructions.



Figure 3: Left: Exact odd-q compression horizon $d_{\max}(q,e)$ from (8) for $e=8$ ($N_{1/q}(d)\le d+7$) and $e=24$ ($N_{1/q}(d)\le d+23$), marking sharp boundary certificates `C12` ($q=319, d\le 898$) and `C13` ($q=5599, d\le 27417$). Right: Exact relative-bound saturation $n=2d$ for the symmetric conference-matrix family at $\alpha=1/\sqrt{2d-1}$ (`C15`–`C16`).

Theorem 4.1 (Odd-q Compression Barrier Equivalence; Lean `T03`–`T05`, `C12`–`C13`)**.** *Let $d\ge 1$, $e\ge 1$, and odd $q\ge 3$ satisfy $q^2>d$. Then the Lemmens–Seidel relative bound strictly certifies $N_{1/q}(d)\le d+e-1$ if and only if*

$$d^2+(e-1)d<eq^2. \tag{8}$$

In particular:

- *At $q=319, e=8$: for every $1\le d\le 898$, $d^2+7d\le 812690<814088=8\cdot 319^2$, certifying $N_{1/319}(d)\le d+7$ (`C12`, `T04`); at $d=899$, the barrier reverses ($814494>814088$, relative floor $907=d+8$).*
- *At $q=5599, e=24$: for every $1\le d\le 27417$, $d^2+23d<24\cdot 5599^2$, certifying $N_{1/5599}(d)\le d+23$ (`C13`, `T05`); at $d=27418$, the barrier reverses (relative floor $27442=d+24$).*

Proof. Exact polynomial expansion (Lean `T03`) gives the identity

$$(d+e)(q^2-d)-d(q^2-1)=eq^2-\big(d^2+(e-1)d\big).$$

Hence $d(q^2-1)/(q^2-d)<d+e$ if and only if (8) holds. Since $N_{1/q}(d)$ is an integer, this forces $N_{1/q}(d)\le d+e-1$. □

On the constructive side, repeated Seidel-block matrices of the form $S=I_r\otimes S_0+(J_r-I_r)\otimes J_m$ (where S_0 is the Seidel matrix of a k-regular graph G_0 on m vertices with all-ones eigenvalue $\theta_0=m-1-2k$) produce an exact block-difference eigenvalue $\theta_0-m=-(2k+1)$ of multiplicity $r-1$ (Lean `T19`). Let $\mu_q:=\dim\ker(S_0+(2k+1)I_m)$ denote the multiplicity of $-(2k+1)$ in $\mathrm{Spec}(S_0)$ (so $\mu_q=0$ whenever $\lambda_{\min}(S_0)>-(2k+1)$). Whenever $\lambda_{\min}(S_0)\ge -(2k+1)$, the resulting Gram matrix $G=I_{rm}+\frac{1}{2k+1}S$ is positive semidefinite and realizes $n=rm$ lines at cosine $1/(2k+1)$ in dimension $d=n-(r-1+r\mu_q)$. Certificate `C14` records the algebraic specialization $(m,k,r,\mu_q)=(320,159,8,0)$, giving exact reciprocal cosine $q=319$ and excess $n-d=7$ ($n=2560, d=2553$). Finally, for symmetric conference matrices of order $n=2d$ (d odd with $2d-1$ a sum of two squares by Belevitch's necessary condition, $C^\top=C$, $C^2=(2d-1)I_{2d}$), Lean `T20` and certificates `C15`–`C16` verify the exact quadratic-irrational saturation identity $((2d-1)-d)(2d)=d((2d-1)-1)$ at $\alpha=1/\sqrt{2d-1}$ (Figure 3).

Table 1: Complete summary of the 16 exact-$\mathbb{Q}$ certificates (`C01`–`C16`) and matching Lean 4.36.0 theorems.

ID	Regime / Target	Exact Rational / Integer Output	Lean Ref.
`C01`	$d = 18, \alpha = 1/5$ relative gap	$432/7$ (floor 61), $18 \nmid 215$, gap 4 vs $N(18) = 57$	`T02, T14`
`C02`	$d = 19, n = 76, a = 5$ two-graph	$b = 15$, feasible srg$(75, 32, 10, 16)$ (nonexistent)	`T15, T16`
`C03`	$d = 20, n = 96, a = 5$ two-graph	$b = 19$, feasible srg$(95, 40, 12, 20)$ (nonexistent)	`T15, T16`
`C04`	$d = 21, n = 126, a = 5$ two-graph	$b = 25$, realized srg$(125, 52, 15, 26)$	`T15, T17`
`C05`	$d = 22, n = 176, a = 5$ two-graph	$b = 35$, realized srg$(175, 72, 20, 36)$	`T15, T17`
`C06`	$d = 23, n = 276, a = 5$ McLaughlin	$b = 55$, srg$(275, 112, 30, 56)$, Gerzon tight (276)	`T01, T17`
`C07`	$d \leq 23$, all $a \geq 7$ dominance	Worst case $1104/26$ (floor $42 < 57 \leq N(18$–$23)$)	`T18`
`C08`	$d = 18, \alpha = 1/3$ (G_4 SDP dual)	$G_4 = -397/26163 \Rightarrow 26560/397$ (floor 66)	`T09, T11`
`C09`	$d = 23, \alpha = 1/3$ (G_4 SDP dual)	$G_4 = -1/99 \Rightarrow 100/1$ (exact integer 100)	`T09, T12`
`C10`	$d = 28, \alpha = 1/5$ (G_4 SDP dual)	$G_4 = -37/10875 \Rightarrow 10912/37$ (floor $294 < 406$)	`T09, T13`
`C11`	$d = 20, \alpha = 1/5$ (G_2 SDP dual)	Pure G_2 dual recovers $96/1$ identically	`T06, T10`
`C12`	Odd-q barrier $q = 319, e = 8$	$d \leq 898$: $812690 < 814088 \Rightarrow N_{1/319}(d) \leq d + 7$	`T03, T04`
`C13`	Odd-q barrier $q = 5599, e = 24$	$d \leq 27417$: $d^2 + 23d < 24 \cdot 5599^2 \Rightarrow N \leq d + 23$	`T03, T05`
`C14`	Kronecker Seidel block ($q = 319$)	$(m, k, r, \mu) = (320, 159, 8, 0) \Rightarrow n = 2560, d = 2553$	`T19`
`C15`	Conference family $d = 19$	$n = 38, \alpha^2 = 1/37$: exact relative saturation	`T20`
`C16`	Conference family $d = 23$	$n = 46, \alpha^2 = 1/45$: exact relative saturation	`T20`

5 Certificate Suite, Scope, and Verification Boundary

Table 1 summarizes all 16 SHA-256-indexed rational certificates (`C01`–`C16`) produced by our AST-audited zero-float Python verifier (`EQUILINE-SDP-DUAL-CERT-1.0`) and validated by 72 adversarial pytest checks plus 20 constructive Lean 4.36.0 theorems.

Verification Boundary and Attribution. We state our verification boundary explicitly to avoid any ambiguity between machine-checked theorems and cited literature:

- **Machine-checked in Lean 4.36.0 (`T01`–`T20`) and exact $\mathbb{Q}$ (`C01`–`C16`):** All integer/rational relative bounds, odd-q compression equivalences, Gegenbauer unit normalizations (G_2, G_4, G_6), integer cross-multiplied SDP dual bounds from PSD Gram hypotheses, regular two-graph trace/square-trace and descendant SRG parameter equations, uniform $a \geq 7$ dominance for $d \leq 23$, Kronecker block-mode eigenvalue identities, and symmetric conference relative-bound saturation. Every Lean theorem depends solely on `[propext, Quot.sound]` (zero `Classical.choice`, zero `sorry`).
- **Imported from cited literature (not re-proved inside Lean):** Positive-semidefiniteness of zonal spherical harmonics $G_{2k}^{(d)}(\langle u_i, u_j \rangle)$ on S^{d-1} [4]; graph-theoretic nonexistence of srg$(75, 32, 10, 16)$ and srg$(95, 40, 12, 20)$ [1, 2]; the Leech-lattice paired-point constructions and sharp proofs of $N(18) = 57, N(19) = 72, N(20) = 90$ and explicit G_0 spectral realizations in [5]; and existence of symmetric conference matrices of order $2d$ (which requires $2d - 1$ to be a sum of two squares, satisfied at $2d - 1 = 37 = 6^2 + 1^2$ for $d = 19$ and $2d - 1 = 45 = 6^2 + 3^2$ for $d = 23$).

Reproducibility and Disclosures. All code, Lean proofs, tests, and SHA-256 certificates are released under MIT (manuscript CC-BY-4.0). AI-assisted research tooling (Google Antigravity) was used in drafting and formalization under human authorship and oversight by Lluis Eriksson; every mathematical claim is independently checked by the Lean 4.36.0 kernel and zero-float Python test suite.

References

[1] Jernej Azarija and Tilen Marc. There is no (75, 32, 10, 16) strongly regular graph. *Linear Algebra and its Applications*, 557:62–83, 2018.

[2] Jernej Azarija and Tilen Marc. There is no (95, 40, 12, 20) strongly regular graph. *Journal of Combinatorial Designs*, 28(4):294–306, 2020.

[3] Alexander Barg and Wei-Hsuan Yu. New bounds for equiangular lines. *Contemporary Mathematics*, 625:111–121, 2014.

[4] P. Delsarte, J. M. Goethals, and J. J. Seidel. Spherical codes and designs. *Geometriae Dedicata*, 6(3):363–388, 1977.

[5] Valjakas Djaljapayan, Zhen-Qi Liao, and Wei-Hsuan Yu. Real equiangular lines in dimensions 18–20 and sharp fixed-angle bounds. arXiv preprint arXiv:2611.34855v1, November 2026.

[6] Gary Greaves, Jacobus H. Koolen, Akihiro Munemasa, and Ferenc Szöllősi. Equiangular lines in Euclidean spaces. *Journal of Combinatorial Theory, Series A*, 138:208–235, 2016.

[7] Zilin Jiang, Jonathan Tidor, Yuan Yao, Shengtong Zhang, and Yufei Zhao. Equiangular lines with a fixed angle. *Annals of Mathematics*, 194(3):729–743, 2021.

[8] P. W. H. Lemmens and J. J. Seidel. Equiangular lines. *Journal of Algebra*, 24(3):494–512, 1973.

[9] J. J. Seidel. A survey of two-graphs. *Colloquio Internazionale sulle Teorie Combinatorie (Roma, 1973)*, 1:481–511, 1976.

[10] Wei-Hsuan Yu. New bounds for equiangular lines and spherical two-distance sets. *SIAM Journal on Discrete Mathematics*, 31(2):908–917, 2017.