EXACT ALL-COUPLING NON-PERTURBATIVE BOUNDS FOR LATTICE YANG–MILLS THEORY WITH TOPOLOGICAL θ-TERM: EXACT 2D θ-VACUA, DASHEN TRANSITIONS, AND LARGE-N 4D SUSCEPTIBILITY CONTROLS

LLUÍS ERIKSSON

Abstract. We establish four cluster-expansion-free, rigorous non-perturbative theorems for lattice gauge theories coupled to a topological θ-term, spanning both exactly solvable two-dimensional U(1) θ-vacua and four-dimensional SU(N) Yang–Mills theory ($N \geq 2$). First, we record the general Vafa–Witten free-energy dominance $|Z_\Lambda(\beta,\theta)| \leq Z_\Lambda(\beta,0)$ and imaginary-θ convexity (strict whenever Q is non-constant), and derive the exact finite-volume and thermodynamic solution of the 2D U(1) Villain θ-model on $\mathbb{T}^2_L$ ($L \geq 2$), exhibiting exact topological susceptibility $\chi_t^{\mathrm{V}}(\beta) = (4\pi^2\beta)^{-1}$, an exact first-order Dashen CP-breaking phase transition at $\theta = \pi$ with latent jump $\Delta q(\beta) = (2\pi\beta)^{-1}$, and explicit complex-$\theta$ Yang–Lee zero-free strips. Second, for the 2D U(1) Wilson θ-model with exact integer plaquette winding charge $Q(U) = \frac{1}{2\pi}\sum_p \phi_p \in \mathbb{Z}$, we prove exact zero-quadrature modified-Bessel series representations for the transfer eigenvalues $c_n^{\mathrm{W}}(\beta,\theta)$ and the exact topological susceptibility $\chi_t^{\mathrm{W}}(\beta) = \frac{1}{12} + (\pi^2 I_0(\beta))^{-1}\sum_{k=1}^{\infty}(-1)^k I_k(\beta)/k^2$, proving exact Dashen degeneracy and sector dominance $|c_n^{\mathrm{W}}(\beta,\pi)| < c_0^{\mathrm{W}}(\beta,\pi) = c_1^{\mathrm{W}}(\beta,\pi) < I_0(\beta)$ ($n \notin \{0,1\}$) and exact weak-coupling convergence $4\pi^2\beta\,\chi_t^{\mathrm{W}}(\beta) \to 1$. Third, for 4D SU($N$) lattice Yang–Mills theory on $\mathbb{T}^4_L$ ($L \geq 2$, $N \geq 2$) with the standard parity-odd four-leaf clover topological charge $Q_{\mathrm{clov}}(U)$ ($Q_{\mathrm{clov}}(P_0 U) = -Q_{\mathrm{clov}}(U)$), we prove the explicit uniform link-gradient bound $\|\nabla_e Q_{\mathrm{clov}}(U)\|_{\mathrm{HS}} \leq 3\sqrt{N}/(2\pi^2)$ and combine it with the Bakry–Émery log-Sobolev inequality on the planar strong-coupling window $\gamma = \beta/N^2 < 1/48$ to prove volume- and N-uniform bounds $\chi_t^{\mathrm{clov}}(\gamma,N) \leq 9/(\pi^4\kappa_0(4,\gamma))$ and $\psi_\Lambda(\gamma N^2,\theta_I) - \psi_\Lambda(\gamma N^2,0) \leq \frac{9}{2\pi^4\kappa_0(4,\gamma)}\theta_I^2$ ($\theta = i\theta_I \in i\mathbb{R}$), matching the $O(1)$ Witten–Veneziano large-N scaling uniformly across all $N \geq 2$ and $|\Lambda|$. Fourth, on Lüscher's admissible ensemble $\mathcal{U}_\varepsilon$ ($\|I_N - U_p\|_{\mathrm{op}} \leq \varepsilon$ for any $\varepsilon > 0$, evaluated within $\varepsilon < 1/30$), we prove all-coupling pointwise clover density bounds $|q_{\mathrm{clov}}(x)| \leq 3N\varepsilon^2/(4\pi^2)$ and imaginary-θ free-energy shift bounds $|\psi_{\Lambda,\varepsilon}(\beta,\theta_I) - \psi_{\Lambda,\varepsilon}(\beta,0)| \leq \frac{3N\varepsilon^2}{4\pi^2}|\theta_I|$ for all $\beta \geq 0$. All 25 quantitative benchmarks are certified in 140-digit outward-rounded interval arithmetic (zero black-box special functions, zero quadrature), cross-checked against 40-digit quadrature and direct 2^4 SU(2)/SU(3) lattice simulations, and their integer-scaled (10^{12}) inequalities are mechanically verified in a standalone zero-import, zero-axiom Lean 4.36.0 kernel.

1. Introduction

The vacuum angle θ of Yang–Mills theory sits at the intersection of non-perturbative gauge topology, the strong CP problem, spontaneous CP violation at $\theta = \pi$ (the Dashen phenomenon [4]), and the Witten–Veneziano resolution of the U(1)$_A$ problem [13, 9, 14, 8, 10]. On the lattice, real $\theta \neq 0$ introduces a severe complex-action sign problem, motivating exact dual analyses in 2D U(1) models [11, 6, 3] and imaginary-θ analytic continuation ($\theta = i\theta_I$) together with geometric admissibility [7] in 4D SU(N) gauge theory [10].

In this paper we establish four exact, cluster-expansion-free non-perturbative theorems governing θ-dependence across both 2D U(1) and 4D SU(N) lattice gauge theories:

(1) **Vafa–Witten dominance, imaginary-θ convexity, and exact 2D Villain θ-vacuum (Theorem 2.1):** On any finite lattice Λ with any parity-odd topological charge Q, $|Z_\Lambda(\beta,\theta)| \leq Z_\Lambda(\beta,0)$ for all $\theta \in \mathbb{R}$, while $\psi_\Lambda(\beta,\theta_I) = |\Lambda|^{-1}\ln Z_\Lambda(\beta,i\theta_I)$ is even and convex

Date: November 29, 2026.

in θ_I. For the 2D U(1) Villain model on $\mathbb{T}^2_L$ ($L \geq 2$), exact Poisson summation yields $\chi_t^{\mathrm{V}}(\beta) = (4\pi^2\beta)^{-1}$, an exact Dashen cusp at $\theta = \pi$, and explicit complex-θ Yang–Lee zero-free strips.

(2) **Exact Bessel series for the 2D Wilson U(1) θ-model (Theorem 3.1):** Using the exact integer plaquette winding charge $Q(U) = \frac{1}{2\pi}\sum_{p\in P}\phi_p \in \mathbb{Z}$ on $\mathbb{T}^2_L$, we prove exact modified-Bessel series for every transfer eigenvalue $c_n^{\mathrm{W}}(\beta,\theta)$ and for $\chi_t^{\mathrm{W}}(\beta)$, proving exact Dashen degeneracy and sector dominance $|c_n^{\mathrm{W}}(\beta,\pi)| < c_0^{\mathrm{W}}(\beta,\pi) = c_1^{\mathrm{W}}(\beta,\pi) < I_0(\beta)$ ($n \notin \{0,1\}$) and $\lim_{\beta\to\infty} 4\pi^2\beta\,\chi_t^{\mathrm{W}}(\beta) = 1$.

(3) **Volume- and N-uniform 4D SU(N) clover susceptibility and imaginary-θ Herbst bounds (Theorem 4.1):** Defining the standard 4-leaf clover charge $Q_{\mathrm{clov}}(U)$ on $\mathbb{T}^4_L$ ($L \geq 2$, $N \geq 2$), we prove exact pointwise parity oddness $Q_{\mathrm{clov}}(P_0U) = -Q_{\mathrm{clov}}(U)$ and the explicit uniform link-gradient bound $\|\nabla_e Q_{\mathrm{clov}}(U)\|_{\mathrm{HS}} \leq 3\sqrt{N}/(2\pi^2)$. Applying Bakry–Émery Poincaré and Herbst inequalities [1, 2] on the planar strong-coupling window $\gamma = \beta/N^2 < 1/48$ yields volume- and N-uniform bounds $\chi_t^{\mathrm{clov}}(\gamma,N) \leq 9/(\pi^4\kappa_0(4,\gamma))$ and $\psi_\Lambda(\gamma N^2,\theta_I) - \psi_\Lambda(\gamma N^2,0) \leq \frac{9}{2\pi^4\kappa_0(4,\gamma)}\theta_I^2$, exhibiting exact $O(1)$ Witten–Veneziano large-N scaling.

(4) **All-coupling 4D SU(N) admissible clover bounds (Theorem 5.1):** On Lüscher's admissible set $\mathcal{U}_\varepsilon$ ($\|I_N - U_p\|_{\mathrm{op}} \leq \varepsilon$, any $\varepsilon > 0$), we prove pointwise $|q_{\mathrm{clov}}(x)| \leq 3N\varepsilon^2/(4\pi^2)$ and all-coupling imaginary-θ free-energy bounds $|\psi_{\Lambda,\varepsilon}(\beta,\theta_I) - \psi_{\Lambda,\varepsilon}(\beta,0)| \leq \frac{3N\varepsilon^2}{4\pi^2}|\theta_I|$ for every $\beta \geq 0$.

All 25 benchmarks across Tables 1–3 are certified at 140-digit outward-rounded interval precision (`mpmath.iv`, `mp.mp.dps = 140`, `mp.iv.dps = 140`, zero black-box special functions, zero quadrature), cross-checked against 40-digit quadrature and direct 2^4 SU(2)/SU(3) lattice tests, and their integer-scaled (10^{12}) inequalities are mechanically verified in a standalone zero-import, zero-axiom Lean 4.36.0 kernel (`ThetaVacuumCerts.lean`) [5].

2. Vafa–Witten Dominance and Exact 2D U(1) Villain θ-Vacuum

Let Λ be a finite periodic hypercubic torus $\mathbb{T}^d_L$ ($L \geq 2$, $d \geq 2$) with link gauge variables $U \in G^{|E|}$ ($G = \mathrm{U}(1)$ or $\mathrm{SU}(N)$), bounded measurable parity-even lattice gauge action $S(U) \in \mathbb{R}$, and measurable real-valued parity-odd topological charge $Q(U)$ satisfying $Q(P_0U) = -Q(U)$ almost everywhere under a reflection P_0 that preserves Haar measure dU and $S(U)$, with $\int e^{-\beta S(U) - \theta_I Q(U)}\,dU < \infty$ for all $\theta_I \in \mathbb{R}$ (automatically satisfied when Q is bounded, or for the Villain charge Q^{V} whenever $\beta > 0$). For real $\theta \in \mathbb{R}$ and imaginary $\theta = i\theta_I$ ($\theta_I \in \mathbb{R}$), define

$$Z_\Lambda(\beta,\theta) := \int_{G^{|E|}} e^{-\beta S(U) + i\theta Q(U)}\,dU, \tag{1}$$

$$\psi_\Lambda(\beta,\theta_I) := \frac{1}{|\Lambda|}\ln Z_\Lambda(\beta, i\theta_I) = \frac{1}{|\Lambda|}\ln \int_{G^{|E|}} e^{-\beta S(U) - \theta_I Q(U)}\,dU. \tag{2}$$

For the 2D U(1) Villain θ-model on $\mathbb{T}^2_L$ ($V = |P| = L^2$), each plaquette carries angle $\phi_p \in (-\pi,\pi]$ (subject to the torus Bianchi constraint $\prod_{p\in P} e^{i\phi_p} = 1$, i.e. $\sum_p \phi_p \in 2\pi\mathbb{Z}$) and an auxiliary Villain integer $m_p \in \mathbb{Z}$, with unfolded field $\tilde{\phi}_p := \phi_p + 2\pi m_p \in \mathbb{R}$ and exact integer topological charge $Q^{\mathrm{V}} := \frac{1}{2\pi}\sum_{p\in P}\tilde{\phi}_p \in \mathbb{Z}$:

$$Z^{\mathrm{V}}_{\mathbb{T}^2_L}(\beta,\theta) := \int_{(-\pi,\pi]^V} \delta_{\mathrm{U}(1)}\left(\prod_{p\in P} e^{i\phi_p}\right) \prod_{p\in P}\left(\sum_{m_p\in\mathbb{Z}} e^{-\frac{\beta}{2}(\phi_p+2\pi m_p)^2 + i\frac{\theta}{2\pi}(\phi_p+2\pi m_p)}\,\frac{d\phi_p}{2\pi}\right). \tag{3}$$

Theorem 2.1 (Vafa–Witten dominance, imaginary-θ convexity, and exact 2D Villain θ-vacuum). (1) ***Vafa–Witten dominance and imaginary-θ convexity:*** *For every finite lattice Λ, every*

$\beta \geq 0$, every $\theta \in \mathbb{R}$, and every $\theta_I \in \mathbb{R}$, $Z_\Lambda(\beta,\theta) \in \mathbb{R}$ satisfies $|Z_\Lambda(\beta,\theta)| \leq Z_\Lambda(\beta,0)$ (so whenever $Z_\Lambda(\beta,\theta) > 0$, the vacuum energy density $E_{\mathrm{vac},\Lambda}(\beta,\theta) := -|\Lambda|^{-1}\ln[Z_\Lambda(\beta,\theta)/Z_\Lambda(\beta,0)]$ is non-negative and minimized at $\theta = 0$). Furthermore, $\psi_\Lambda(\beta,\theta_I)$ is smooth, even, and convex in θ_I, with $\psi''_\Lambda(\beta,0) = |\Lambda|^{-1}\langle Q^2\rangle_{\beta,0} = \chi_{t,\Lambda}(\beta) \geq 0$.

(2) ***Exact 2D Villain partition function and free energy:*** *For every $L \geq 2$ ($V = L^2$), every $\beta > 0$, and every $z \in \mathbb{C}$,*

$$Z^{\mathrm{V}}_{\mathbb{T}^2_L}(\beta,z) = (2\pi\beta)^{-V/2}\sum_{n\in\mathbb{Z}}\exp\left(-\frac{V}{2\beta}\left(n-\frac{z}{2\pi}\right)^2\right). \tag{4}$$

For real $\theta \in \mathbb{R}$, $Z^{\mathrm{V}}_{\mathbb{T}^2_L}(\beta,\theta) > 0$ for all V, and the thermodynamic vacuum energy density $E^{\mathrm{V}}_{\mathrm{vac}}(\beta,\theta) := \lim_{L\to\infty} E^{\mathrm{V}}_{\mathrm{vac},\mathbb{T}^2_L}(\beta,\theta)$ and topological susceptibility $\chi^{\mathrm{V}}_t(\beta)$ are

$$E^{\mathrm{V}}_{\mathrm{vac}}(\beta,\theta) = \frac{1}{2\beta}\min_{n\in\mathbb{Z}}\left(n-\frac{\theta}{2\pi}\right)^2, \qquad \chi^{\mathrm{V}}_t(\beta) = \frac{1}{4\pi^2\beta}, \qquad E^{\mathrm{V}}_{\mathrm{vac}}(\beta,\pi) = \frac{1}{8\beta}. \tag{5}$$

At $\theta = \pi$, the sectors $n = 0$ and $n = 1$ become degenerate, producing a first-order Dashen CP-breaking cusp with one-sided derivative jump $\partial_\theta E^{\mathrm{V}}_{\mathrm{vac}}(\beta,\pi^-) - \partial_\theta E^{\mathrm{V}}_{\mathrm{vac}}(\beta,\pi^+) = (2\pi\beta)^{-1}$.

(3) ***Exact complex-θ Yang–Lee zero-free strip:*** *For every $L \geq 2$ and $\beta > 0$, $Z^{\mathrm{V}}_{\mathbb{T}^2_L}(\beta,z) \neq 0$ for every complex $z = \theta_R + i\theta_I$ satisfying $|\theta_R| < \pi$ and $\sum_{n\neq 0}\exp\left(-\frac{V}{2\beta}[n^2 - n\theta_R/\pi]\right) < 1$ (in particular, on the entire vertical strip $|\theta_R| < \pi - \frac{2\pi\beta}{V}\ln 3$ for all $\theta_I \in \mathbb{R}$ when $V > 2\beta\ln 3$).*

Proof. Part (1): Parity P_0 maps $e^{i\theta Q(U)} \mapsto e^{-i\theta Q(U)}$, so $Z_\Lambda(\beta,\theta) = \int e^{-\beta S}\cos(\theta Q)\,dU \in \mathbb{R}$, and $|Z_\Lambda(\beta,\theta)| \leq \int e^{-\beta S}|\cos(\theta Q)|\,dU \leq Z_\Lambda(\beta,0)$ [8]. Similarly, $Z_\Lambda(\beta,-i\theta_I) = Z_\Lambda(\beta,i\theta_I) > 0$, and differentiating $\psi_\Lambda(\beta,\theta_I)$ twice gives $\psi''_\Lambda(\beta,\theta_I) = |\Lambda|^{-1}\,\mathrm{Var}_{\beta,\theta_I}(Q) \geq 0$.

Part (2): Expanding $\delta_{\mathrm{U}(1)}(\prod_p e^{i\phi_p}) = \sum_{n\in\mathbb{Z}}\prod_p e^{-in\phi_p}$ and noting $e^{-in\phi_p} = e^{-in(\phi_p+2\pi m_p)} = e^{-in\tilde{\phi}_p}$ combines the sum over $m_p \in \mathbb{Z}$ and integral over $(-\pi,\pi]$ into a single Gaussian integral over $\tilde{\phi}_p \in \mathbb{R}$:

$$c^{\mathrm{V}}_n(\beta,z) = \int_{-\infty}^{\infty}\frac{d\tilde{\phi}}{2\pi}\,e^{-\frac{\beta}{2}\tilde{\phi}^2 + i\left(\frac{z}{2\pi}-n\right)\tilde{\phi}} = \frac{1}{\sqrt{2\pi\beta}}\exp\left(-\frac{(n-z/(2\pi))^2}{2\beta}\right),$$

proving (4). Taking $-\lim_{V\to\infty} V^{-1}\ln(Z^{\mathrm{V}}(\beta,\theta)/Z^{\mathrm{V}}(\beta,0))$ yields (5) and the cusp jump at $\theta = \pi$ directly.

Part (3): Writing $z = \theta_R + i\theta_I$, $|c^{\mathrm{V}}_n(\beta,z)/c^{\mathrm{V}}_0(\beta,z)|^V = \exp(-\frac{V}{2\beta}(n^2 - n\theta_R/\pi))$, strictly independent of θ_I. By the triangle inequality, $Z^{\mathrm{V}}_{\mathbb{T}^2_L}(\beta,z) \neq 0$ whenever $\sum_{n\neq 0}\exp(-\frac{V}{2\beta}(n^2 - n\theta_R/\pi)) < 1$. For $|\theta_R| \leq \pi(1-\eta)$ with $\eta \in (0,1)$, $n^2 - n\theta_R/\pi \geq |n|(|n| - 1 + \eta) \geq |n|\eta$, so $\sum_{n\neq 0} e^{-V|n|\eta/(2\beta)} = \frac{2e^{-V\eta/(2\beta)}}{1-e^{-V\eta/(2\beta)}} < 1 \iff e^{-V\eta/(2\beta)} < 1/3 \iff \eta > \frac{2\beta}{V}\ln 3$. (Note that since (4) is a Jacobi ϑ_3 function, its exact complex zeros lie solely on the vertical lines $\theta_R \in \pi + 2\pi\mathbb{Z}$.) □

3. Exact Bessel Series for the 2D Wilson U(1) θ-Model

For the 2D U(1) Wilson gauge action [12] on $\mathbb{T}^2_L$ ($L \geq 2$, $V = L^2 \geq 4$) with exact integer plaquette topological charge $Q(U) := \frac{1}{2\pi}\sum_{p\in P}\phi_p \in \mathbb{Z}$ ($\phi_p \in (-\pi,\pi]$, parity-odd off the measure-zero set $\bigcup_{p\in P}\{\phi_p = \pi\}$) [11, 3], the V-fold circular convolution of $h(\phi) := e^{\beta\cos\phi + i\frac{\theta}{2\pi}\phi}$ is continuous at 0 with ℓ^1 Fourier coefficients $(c^{\mathrm{W}}_n(\beta,\theta))^V = O(|n|^{-V})$ (for $V \geq 4$), yielding the exact partition function

$$Z^{\mathrm{W}}_{\mathbb{T}^2_L}(\beta,\theta) := \int_{(-\pi,\pi]^V}\delta_{\mathrm{U}(1)}\left(\prod_{p\in P}e^{i\phi_p}\right)\prod_{p\in P}e^{\beta\cos\phi_p + i\frac{\theta}{2\pi}\phi_p}\,\frac{d\phi_p}{2\pi} = \sum_{n\in\mathbb{Z}}\left(c^{\mathrm{W}}_n(\beta,\theta)\right)^V, \tag{6}$$

where $c_n^{\mathrm{W}}(\beta,\theta) := \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} e^{\beta\cos\phi}\cos((\frac{\theta}{2\pi}-n)\phi)$, and thermodynamic vacuum energy density $E_{\mathrm{vac}}^{\mathrm{W}}(\beta,\theta) := -\lim_{L\to\infty} V^{-1}\ln[Z_{\mathbb{T}_L^2}^{\mathrm{W}}(\beta,\theta)/Z_{\mathbb{T}_L^2}^{\mathrm{W}}(\beta,0)]$ whenever $Z_{\mathbb{T}_L^2}^{\mathrm{W}}(\beta,\theta)>0$ for large L.

Theorem 3.1 (Exact Bessel series, Dashen transition, and weak-coupling limit in 2D Wilson U(1))**.** *For every $\beta\geq 0$, every $\theta\in\mathbb{R}$, and every $n\in\mathbb{Z}$ (with $\alpha_n := \frac{\theta}{2\pi}-n$):*

(1) ***Exact zero-quadrature Bessel series and $\theta=\pi$ sector dominance:*** *Whenever $\alpha_n\notin\mathbb{Z}$ (and by limit $I_{|\alpha_n|}(\beta)$ when $\alpha_n\in\mathbb{Z}$),*

$$c_n^{\mathrm{W}}(\beta,\theta) = \frac{\sin(\pi\alpha_n)}{\pi}\sum_{k\in\mathbb{Z}}\frac{(-1)^k I_{|k|}(\beta)}{\alpha_n+k}. \tag{7}$$

At $\theta=\pi$, $c_0^{\mathrm{W}}(\beta,\pi)=c_1^{\mathrm{W}}(\beta,\pi)$ satisfies the exact alternating series

$$c_0^{\mathrm{W}}(\beta,\pi) = \frac{2}{\pi}I_0(\beta) - \frac{4}{\pi}\sum_{k=1}^{\infty}\frac{(-1)^k I_k(\beta)}{4k^2-1} \;<\; I_0(\beta) = c_0^{\mathrm{W}}(\beta,0), \tag{8}$$

together with strict dominance $|c_n^{\mathrm{W}}(\beta,\pi)|<c_0^{\mathrm{W}}(\beta,\pi)$ for every $n\notin\{0,1\}$ and all $\beta\geq 0$. Consequently, $Z_{\mathbb{T}_L^2}^{\mathrm{W}}(\beta,\pi)>0$ for all sufficiently large L, and the exact Dashen vacuum energy is $E_{\mathrm{vac}}^{\mathrm{W}}(\beta,\pi)=\ln(I_0(\beta)/c_0^{\mathrm{W}}(\beta,\pi))>0$.

(2) ***Exact topological susceptibility and weak-coupling asymptotics:*** *The thermodynamic topological susceptibility $\chi_t^{\mathrm{W}}(\beta) := \lim_{L\to\infty}\frac{1}{L^2}\langle Q^2\rangle_{\beta,0}^{\mathrm{W}}$ admits the exact Bessel series*

$$\chi_t^{\mathrm{W}}(\beta) = \frac{1}{12} + \frac{1}{\pi^2 I_0(\beta)}\sum_{k=1}^{\infty}\frac{(-1)^k I_k(\beta)}{k^2}, \tag{9}$$

satisfying $\chi_t^{\mathrm{W}}(0)=\frac{1}{12}$ and $\lim_{\beta\to\infty}4\pi^2\beta\,\chi_t^{\mathrm{W}}(\beta)=1$.

Proof. For (7), substitute the uniformly convergent Jacobi–Anger expansion $e^{\beta\cos\phi}=\sum_{k\in\mathbb{Z}}I_{|k|}(\beta)e^{ik\phi}$ into $c_n^{\mathrm{W}}(\beta,\theta)=\int_{-\pi}^{\pi}\frac{d\phi}{2\pi}e^{\beta\cos\phi+i\alpha_n\phi}$ and integrate term-by-term using

$$\int_{-\pi}^{\pi}\frac{d\phi}{2\pi}\,e^{i(\alpha_n+k)\phi} = \frac{\sin(\pi(\alpha_n+k))}{\pi(\alpha_n+k)} = \frac{(-1)^k\sin(\pi\alpha_n)}{\pi(\alpha_n+k)}.$$

At $\theta=\pi$, $\alpha_0=\frac{1}{2}$ and $\alpha_1=-\frac{1}{2}$, so $\cos(\alpha_0\phi)=\cos(\alpha_1\phi)=\cos(\phi/2)>0$ on $(-\pi,\pi)$, yielding $0<c_0^{\mathrm{W}}(\beta,\pi)=c_1^{\mathrm{W}}(\beta,\pi)<\int_{-\pi}^{\pi}\frac{d\phi}{2\pi}e^{\beta\cos\phi}=I_0(\beta)$. Pairing $+k$ and $-k$ in (7) at $\alpha_0=\frac{1}{2}$ via $\frac{1}{1/2+k}+\frac{1}{1/2-k}=\frac{-4}{4k^2-1}$ proves (8).

To prove strict sector dominance $|c_n^{\mathrm{W}}(\beta,\pi)|<c_0^{\mathrm{W}}(\beta,\pi)$ for every $n\notin\{0,1\}$, let $2m+1 := |2n-1|\geq 3$ ($m\geq 1$) and $w(\phi) := e^{\beta\cos\phi}$ (even, strictly positive, with $-w'(\phi)=\beta\sin\phi\,w(\phi)\geq 0$ on $[0,\pi]$). Integrating by parts on $[0,\pi]$ gives

$$c_n^{\mathrm{W}}(\beta,\pi) = \frac{1}{\pi}\int_0^{\pi} w(\phi)\cos\!\left(\frac{2m+1}{2}\phi\right)d\phi = \frac{2(-1)^m e^{-\beta}}{(2m+1)\pi} + \frac{2}{(2m+1)\pi}\int_0^{\pi}(-w'(\phi))\sin\!\left(\frac{2m+1}{2}\phi\right)d\phi,$$

whereas for $m=0$ ($n\in\{0,1\}$), $c_0^{\mathrm{W}}(\beta,\pi)=\frac{2e^{-\beta}}{\pi}+\frac{2}{\pi}\int_0^{\pi}(-w'(\phi))\sin(\frac{\phi}{2})\,d\phi$. Because $|\sin(\frac{2m+1}{2}\phi)|\leq(2m+1)\sin(\frac{\phi}{2})$ on $[0,\pi]$ and $-w'(\phi)\geq 0$, the integral term of $|c_n^{\mathrm{W}}(\beta,\pi)|$ is bounded by that of $c_0^{\mathrm{W}}(\beta,\pi)$, while the boundary term obeys $\frac{2e^{-\beta}}{(2m+1)\pi}<\frac{2e^{-\beta}}{\pi}$ strictly (since $2m+1\geq 3$ and $e^{-\beta}>0$). Thus $|c_n^{\mathrm{W}}(\beta,\pi)|<c_0^{\mathrm{W}}(\beta,\pi)$ for all $n\notin\{0,1\}$, and moreover $|c_n^{\mathrm{W}}(\beta,\pi)|\leq\frac{2e^{\beta}}{(2m+1)\pi}=O(1/|n|)$. Hence $\rho := \sup_{n\notin\{0,1\}}|c_n^{\mathrm{W}}(\beta,\pi)/c_0^{\mathrm{W}}(\beta,\pi)|<1$, and for $V\geq 4$,

$$\sum_{n\notin\{0,1\}}\left|\frac{c_n^{\mathrm{W}}(\beta,\pi)}{c_0^{\mathrm{W}}(\beta,\pi)}\right|^V \leq \rho^{V-4}\sum_{n\notin\{0,1\}}\left|\frac{c_n^{\mathrm{W}}(\beta,\pi)}{c_0^{\mathrm{W}}(\beta,\pi)}\right|^4 \xrightarrow[V\to\infty]{} 0,$$

so $Z^{\mathrm{W}}_{\mathbb{T}^2_L}(\beta,\pi) = 2(c_0^{\mathrm{W}}(\beta,\pi))^V(1+o(1)) > 0$ for large L and $E^{\mathrm{W}}_{\mathrm{vac}}(\beta,\pi) = \ln(I_0(\beta)/c_0^{\mathrm{W}}(\beta,\pi)) > 0$.

For (9), integration by parts on $c_n^{\mathrm{W}}(\beta,\theta)$ and its first two θ-derivatives shows $c_n^{\mathrm{W}}, \partial_\theta c_n^{\mathrm{W}}, \partial_\theta^2 c_n^{\mathrm{W}} = O(1/|n|)$ uniformly for $|\theta| \le \pi$. For $V = L^2 \ge 4$, $\sum_{n\in\mathbb{Z}}(c_n^{\mathrm{W}}(\beta,\theta))^V$ can therefore be differentiated term-by-term twice at $\theta = 0$ (noting $\partial_\theta c_0^{\mathrm{W}}(\beta,0) = 0$ and $\partial_\theta c_{-n}^{\mathrm{W}}(\beta,0) = -\partial_\theta c_n^{\mathrm{W}}(\beta,0)$ cancels in the first derivative), giving

$$\frac{1}{V}\langle Q^2\rangle^{\mathrm{W}}_{\beta,0} = -\frac{\sum_{n\in\mathbb{Z}}\big[(V-1)(c_n^{\mathrm{W}}(\beta,0))^{V-2}(\partial_\theta c_n^{\mathrm{W}}(\beta,0))^2 + (c_n^{\mathrm{W}}(\beta,0))^{V-1}\partial_\theta^2 c_n^{\mathrm{W}}(\beta,0)\big]}{\sum_{n\in\mathbb{Z}}(c_n^{\mathrm{W}}(\beta,0))^V}.$$

Because $c_0^{\mathrm{W}}(\beta,0) = I_0(\beta) > I_{|n|}(\beta) = c_n^{\mathrm{W}}(\beta,0) \ge 0$ for all $n \ne 0$ (as $I_0(\beta) - I_n(\beta) = \int_{-\pi}^{\pi}\frac{d\phi}{2\pi}e^{\beta\cos\phi}(1-\cos(n\phi)) > 0$) and $\partial_\theta c_0^{\mathrm{W}}(\beta,0) = 0$, dividing numerator and denominator by $I_0(\beta)^V$ leaves the $n = 0$ term $-I_0(\beta)^{-1}\partial_\theta^2 c_0^{\mathrm{W}}(\beta,0)$ while every $n \ne 0$ contribution (both $(V-1)(I_{|n|}/I_0)^{V-2}(\partial_\theta c_n^{\mathrm{W}}/I_0)^2$ and $(I_{|n|}/I_0)^{V-1}\partial_\theta^2 c_n^{\mathrm{W}}/I_0$, with convergent $\sum_{n\ne0}(I_{|n|}/I_0)^2((\partial_\theta c_n^{\mathrm{W}})^2 + |\partial_\theta^2 c_n^{\mathrm{W}}|)$ for $V \ge 4$) vanishes exponentially as $V \to \infty$ since $I_{|n|}(\beta) \le I_1(\beta) < I_0(\beta)$, leaving

$$\chi_t^{\mathrm{W}}(\beta) = -\frac{1}{I_0(\beta)}\frac{\partial^2 c_0^{\mathrm{W}}(\beta,\theta)}{\partial\theta^2}\bigg|_{\theta=0} = \frac{1}{I_0(\beta)}\int_{-\pi}^{\pi}\frac{d\phi}{2\pi}\,e^{\beta\cos\phi}\left(\frac{\phi}{2\pi}\right)^2.$$

Inserting $e^{\beta\cos\phi} = I_0(\beta) + 2\sum_{k=1}^{\infty} I_k(\beta)\cos(k\phi)$ and integrating term-by-term via $\int_{-\pi}^{\pi}\frac{d\phi}{2\pi}(\frac{\phi}{2\pi})^2 = \frac{1}{12}$ and $\int_{-\pi}^{\pi}\frac{d\phi}{2\pi}(\frac{\phi}{2\pi})^2\cos(k\phi) = \frac{(-1)^k}{2\pi^2k^2}$ $(k \ge 1)$ proves (9). At $\beta = 0$, $I_k(0) = \delta_{k,0}$ gives $\chi_t^{\mathrm{W}}(0) = \frac{1}{12}$. As $\beta \to \infty$, Laplace's method on $\int_{-\pi}^{\pi} e^{-\beta(1-\cos\phi)}\phi^2\,d\phi / \int_{-\pi}^{\pi} e^{-\beta(1-\cos\phi)}\,d\phi = \frac{1}{\beta}(1+O(\beta^{-1}))$ yields $4\pi^2\beta\,\chi_t^{\mathrm{W}}(\beta) \to 1$. □

4. Volume- and N-Uniform 4D SU(N) Clover Susceptibility and Herbst Bounds

Now let $\Lambda = \mathbb{T}^4_L$ $(L \ge 2)$ be a 4D periodic hypercubic torus with gauge group SU(N) $(N \ge 2)$ and bi-invariant Hilbert–Schmidt inner product $\langle X, Y\rangle_{\mathrm{HS}} = \sum_{e\in E}\mathrm{Tr}(X_e^\dagger Y_e)$ on $\mathfrak{su}(N)^{|E|}$. For each site $x \in \mathbb{T}^4_L$ and each ordered plane (μ,ν) $(1 \le \mu < \nu \le 4)$, let $U_{\mu\nu}^{(++)}(x), U_{\mu\nu}^{(-+)}(x), U_{\mu\nu}^{(--)}(x), U_{\mu\nu}^{(+-)}(x)$ be the four standard oriented elementary plaquettes incident to x in the (μ,ν)-plane (all sharing the $(+\mu,+\nu)$ orientation), and define the Hermitian 4-leaf clover field strength and clover topological charge by

$$F_{\mu\nu}^{\mathrm{clov}}(x) := \frac{C_{\mu\nu}(x) - C_{\mu\nu}(x)^\dagger}{2i}, \qquad C_{\mu\nu}(x) := \frac{1}{4}\big(U_{\mu\nu}^{(++)} + U_{\mu\nu}^{(-+)} + U_{\mu\nu}^{(--)} + U_{\mu\nu}^{(+-)}\big)(x), \tag{10}$$

$$q_{\mathrm{clov}}(x) := \frac{1}{4\pi^2}\Big(\mathrm{Tr}\big(F_{12}^{\mathrm{clov}}F_{34}^{\mathrm{clov}}\big) - \mathrm{Tr}\big(F_{13}^{\mathrm{clov}}F_{24}^{\mathrm{clov}}\big) + \mathrm{Tr}\big(F_{14}^{\mathrm{clov}}F_{23}^{\mathrm{clov}}\big)\Big)(x), \qquad Q_{\mathrm{clov}}(U) := \sum_{x\in\mathbb{T}^4_L} q_{\mathrm{clov}}(x). \tag{11}$$

Consider the planar-scaled 4D Wilson measure $d\mu_\beta(U) := Z_\Lambda(\beta)^{-1}e^{-\gamma N^2 S_W(U)}\,dU$ $(\beta = \gamma N^2$, $S_W(U) = \sum_{p\in P}(1 - \frac{1}{N}\,\mathrm{Re}\,\mathrm{Tr}\,U_p))$.

Theorem 4.1 (Volume- and N-uniform 4D SU(N) clover susceptibility and imaginary-θ excess)**.** *Let $\Lambda = \mathbb{T}^4_L$ with $L \ge 2$ and $N \ge 2$.*

(1) ***Exact parity oddness and explicit uniform link-gradient bound:*** *Under spatial reflection P_0 ($x_1 \mapsto -x_1 \bmod L$, $U_{(x,1)} \mapsto U^\dagger_{(P_0x-\hat{1},1)}$, $U_{(x,\nu)} \mapsto U_{(P_0x,\nu)}$ for $\nu \ge 2$), $Q_{\mathrm{clov}}(P_0U) = -Q_{\mathrm{clov}}(U)$ identically for all $U \in \mathrm{SU}(N)^{|E|}$ (hence $\mathbb{E}_{\mu_\beta}[Q_{\mathrm{clov}}] = 0$). Moreover, for every link $e \in E$ and every $U \in \mathrm{SU}(N)^{|E|}$,*

$$\big\|\nabla_e Q_{\mathrm{clov}}(U)\big\|_{\mathrm{HS}} \le \frac{3\sqrt{N}}{2\pi^2}, \qquad \big\|\nabla Q_{\mathrm{clov}}(U)\big\|^2_{\mathrm{HS}} \le \frac{9N|\Lambda|}{\pi^4}. \tag{12}$$

(2) ***Volume- and N-uniform susceptibility and Herbst bounds ($O(1)$ Witten–Veneziano scaling):*** *For every $\gamma \in [0, 1/48)$, set $\kappa_0(4,\gamma) := \frac{1}{2} - 24\gamma > 0$. Then uniformly in $L \geq 2$ and $N \geq 2$ (with $\beta = \gamma N^2$),*

$$\chi_t^{\mathrm{clov}}(\gamma, N) := \frac{1}{|\Lambda|}\mathbb{E}_{\mu_\beta}\left[Q_{\mathrm{clov}}(U)^2\right] \;\leq\; \frac{9}{\pi^4\,\kappa_0(4,\gamma)}, \tag{13}$$

$$0 \;\leq\; \psi_\Lambda(\gamma N^2, \theta_I) - \psi_\Lambda(\gamma N^2, 0) \;\leq\; \frac{9}{2\pi^4\,\kappa_0(4,\gamma)}\,\theta_I^2 \qquad \forall \theta_I \in \mathbb{R}. \tag{14}$$

Proof. For part (1), under P_0 the four clover leaves in any $(1,\nu)$-plane ($\nu \in \{2,3,4\}$) permute with orientation reversal ($C_{1\nu}(x) \mapsto C_{1\nu}(P_0x)^\dagger$), so $F_{1\nu}^{\mathrm{clov}}(x) \mapsto -F_{1\nu}^{\mathrm{clov}}(P_0x)$, whereas in any (ρ,σ)-plane with $\rho,\sigma \in \{2,3,4\}$, $F_{\rho\sigma}^{\mathrm{clov}}(x) \mapsto +F_{\rho\sigma}^{\mathrm{clov}}(P_0x)$ at the exact same site P_0x. Hence $q_{\mathrm{clov}}(P_0U; P_0x) = -q_{\mathrm{clov}}(U;x)$, so $Q_{\mathrm{clov}}(P_0U) = -Q_{\mathrm{clov}}(U)$ identically.

To prove (12), fix a link $e \in E$ and a unit tangent vector $X_e \in \mathfrak{su}(N)$ ($\|X_e\|_{\mathrm{HS}} = 1$), varying $U_e(s) = e^{sX_e}U_e$. Because $L \geq 2$, link e belongs to exactly 6 distinct plaquettes in 4D (two in each of the 3 planes containing the direction of e), and each such plaquette appears at its 4 corners as one of the 4 leaves (each carrying factor $1/4$) in $F_{\mu\nu}^{\mathrm{clov}}(x)$, never appearing in the orthogonal partner $F_{\rho\sigma}^{\mathrm{clov}}(x)$ (since $\{\mu,\nu\} \cap \{\rho,\sigma\} = \emptyset$). Thus link e contributes to $6 \times 4 = 24$ leaf-site terms, each with prefactor $\frac{1}{4\pi^2} \cdot \frac{1}{4}$. For a single plaquette leaf U_p containing e, $\delta_e U_p = Y_e U_p$ with $Y_e^\dagger = -Y_e$ and $\|Y_e\|_{\mathrm{HS}} = 1$, so $\delta_e((U_p - U_p^\dagger)/(2i)) = \frac{1}{2i}(Y_eU_p + U_p^\dagger Y_e)$. Since each unitary leaf has $\|F_p\|_{\mathrm{op}} \leq 1$, $\|F_{\rho\sigma}^{\mathrm{clov}}(x)\|_{\mathrm{HS}} \leq \frac{1}{4}\sum_{j=1}^4 \|F_{\rho\sigma}^{(j)}\|_{\mathrm{HS}} \leq \sqrt{N}$, and Cauchy–Schwarz gives

$$\left|\mathrm{Tr}\left(\frac{Y_eU_p + U_p^\dagger Y_e}{2i}\,F_{\rho\sigma}^{\mathrm{clov}}\right)\right| \leq \frac{1}{2}\Big(\|Y_e\|_{\mathrm{HS}}\|U_pF_{\rho\sigma}^{\mathrm{clov}}\|_{\mathrm{HS}} + \|Y_e\|_{\mathrm{HS}}\|F_{\rho\sigma}^{\mathrm{clov}}U_p^\dagger\|_{\mathrm{HS}}\Big) \leq \sqrt{N}.$$

Summing all 24 leaf-site contributions yields $|\langle \nabla_e Q_{\mathrm{clov}}, X_e\rangle_{\mathrm{HS}}| \leq \frac{24}{4\pi^2\cdot 4}\sqrt{N} = \frac{3\sqrt{N}}{2\pi^2}$, proving the first inequality of (12); summing $\|\nabla_e Q_{\mathrm{clov}}\|_{\mathrm{HS}}^2$ over all $|E| = 4|\Lambda|$ links yields the second.

For part (2), on $\mathrm{SU}(N)^{|E|}$ the bi-invariant Ricci curvature is $\mathrm{Ric} = \frac{N}{2}I$ and the Wilson Hessian obeys $\|N\,\mathrm{Hess}\,S_W(U)\|_{\mathrm{op}} \leq 8(d-1) = 24$, so $\mathrm{Ric} + \gamma N^2\,\mathrm{Hess}\,S_W \geq N(\frac{1}{2} - 24\gamma)I = N\kappa_0(4,\gamma)I > 0$. By the Bakry–Émery theorem [1, 2], μ_β satisfies $\mathrm{LSI}(\rho_0)$ with $\rho_0 := N\kappa_0(4,\gamma)$. Applying the implied Poincaré inequality to the mean-zero observable Q_{clov} gives

$$\chi_t^{\mathrm{clov}}(\gamma, N) = \frac{1}{|\Lambda|}\,\mathrm{Var}_{\mu_\beta}(Q_{\mathrm{clov}}) \leq \frac{1}{|\Lambda|\,N\kappa_0(4,\gamma)}\int \|\nabla Q_{\mathrm{clov}}\|_{\mathrm{HS}}^2\,d\mu_\beta \leq \frac{9}{\pi^4\kappa_0(4,\gamma)},$$

where both $|\Lambda|$ and N cancel identically! Likewise, Herbst's sub-Gaussian lemma for $\mathrm{LSI}(\rho_0)$ applied to Q_{clov} with global Lipschitz constant $L_Q := \sup_U \|\nabla Q_{\mathrm{clov}}\|_{\mathrm{HS}} \leq \frac{3\sqrt{N|\Lambda|}}{\pi^2}$ yields $\ln \mathbb{E}_{\mu_\beta}[e^{-\theta_I Q_{\mathrm{clov}}}] \leq \frac{L_Q^2}{2\rho_0}\theta_I^2 \leq \frac{9|\Lambda|}{2\pi^4\kappa_0(4,\gamma)}\theta_I^2$; dividing by $|\Lambda|$ and noting $\psi_\Lambda(\gamma,\theta_I) \geq \psi_\Lambda(\gamma,0)$ by Theorem 2.1(1) proves (14). □

5. All-Coupling 4D SU(N) Bounds under Lüscher Admissibility

On Lüscher's admissible set $\mathcal{U}_\varepsilon := \{U \in \mathrm{SU}(N)^{|E|} : \|I_N - U_p\|_{\mathrm{op}} \leq \varepsilon\ \forall p \in P\}$ ($\varepsilon > 0$) [7], define the admissible imaginary-θ free energy $\psi_{\Lambda,\varepsilon}(\beta,\theta_I) := |\Lambda|^{-1}\ln\int_{\mathcal{U}_\varepsilon} e^{-\beta S_W(U) - \theta_I Q_{\mathrm{clov}}(U)}\,dU$.

Theorem 5.1 (All-coupling admissible clover density and imaginary-θ shift bounds)**.** *For every $L \geq 2$, every $N \geq 2$, every $\varepsilon > 0$, every $\beta \geq 0$, and every $U \in \mathcal{U}_\varepsilon$:*

$$|q_{\mathrm{clov}}(x)| \;\leq\; \frac{3N\varepsilon^2}{4\pi^2} \quad (\forall x \in \mathbb{T}_L^4), \qquad 0 \;\leq\; \psi_{\Lambda,\varepsilon}(\beta,\theta_I) - \psi_{\Lambda,\varepsilon}(\beta,0) \;\leq\; \frac{3N\varepsilon^2}{4\pi^2}\,|\theta_I| \quad (\forall \theta_I \in \mathbb{R}). \tag{15}$$

Proof. For every plaquette U_p with $\|I_N - U_p\|_{\mathrm{op}} \leq \varepsilon$, the Hermitian field strength $F_p = \frac{1}{2i}((U_p - I_N) - (U_p^\dagger - I_N))$ satisfies $\|F_p\|_{\mathrm{op}} \leq \varepsilon$. By the triangle inequality over the four clover leaves (10),

$\|F^{\mathrm{clov}}_{\mu\nu}(x)\|_{\mathrm{op}} \le \varepsilon$. Since $|\operatorname{Tr}(AB)| \le N\|A\|_{\mathrm{op}}\|B\|_{\mathrm{op}}$ for $N \times N$ matrices, summing the three orthogonal plane pairs in (11) gives $|q_{\mathrm{clov}}(x)| \le \frac{3N\varepsilon^2}{4\pi^2}$. Summing over $x \in \mathbb{T}^4_L$ yields $|Q_{\mathrm{clov}}(U)| \le \frac{3N\varepsilon^2}{4\pi^2}|\Lambda|$ pointwise on $\mathcal{U}_\varepsilon$, so $e^{-\theta_I Q_{\mathrm{clov}}(U)} \le \exp(\frac{3N\varepsilon^2}{4\pi^2}|\theta_I|\,|\Lambda|)$, proving the right-hand inequality in (15). Since $\mathcal{U}_\varepsilon$ is P_0-invariant ($\|I_N - U_p^\dagger\|_{\mathrm{op}} = \|I_N - U_p\|_{\mathrm{op}}$) and has strictly positive Haar measure for every $\varepsilon > 0$ (containing the open neighborhood $\{\|I_N - U_e\|_{\mathrm{op}} < \varepsilon/4\}$ of I_N), parity oddness of Q_{clov} and Jensen's inequality (Theorem 2.1(1)) yield the left-hand inequality. □

6. Certified 140-Digit Benchmarks and Independent Verification

All 25 benchmarks in Tables 1–3 were computed by `certify_theta_vacuum.py` in `mpmath.iv` at 140-digit working and interval precision (`mp.mp.dps = 140`, `mp.iv.dps = 140`), using self-contained series implementations of π, ln, and $I_k(\beta)$ with analytic remainder bounds (zero black-box special functions, zero quadrature), outward-rounded directly via `mp.floor`/`mp.ceil` at scale 10^{12}.

Table 1. Certified 140-digit benchmarks for the exact 2D U(1) Wilson vs. Villain θ-vacuum (Theorems 2.1 and 3.1). Every column is conservatively floored to 6 decimals from its outward-rounded 140-digit binary interval (so exact rationals such as $1/(8\beta)$ at $\beta = 5, 10, 20, 40$ appear as $0.024999, 0.012499, \ldots$).

β	$\chi_t^{\mathrm{W}}(\beta)$	$4\pi^2\beta\,\chi_t^{\mathrm{W}}(\beta)$	$E_{\mathrm{vac}}^{\mathrm{W}}(\beta,\pi)$	$E_{\mathrm{vac}}^{\mathrm{V}}(\beta,\pi) = \frac{1}{8\beta}$
0	0.083333	—	0.451582	—
1/2	0.059495	1.174401	0.305057	0.250000
1	0.040636	1.604254	0.201420	0.125000
2	0.019364	1.528923	0.094194	0.062500
3	0.011060	1.309988	0.054084	0.041666
5	0.005755	1.136150	0.028347	0.024999
10	0.002676	1.056550	0.013203	0.012499
20	0.001300	1.026476	0.006415	0.006249
40	0.000641	1.012852	0.003165	0.003124

Table 2. Certified 140-digit volume- and N-uniform 4D SU(N) Bakry–Émery clover topological susceptibility bounds (13) and imaginary-θ excess bounds (14) valid for all $N \ge 2$ and $L \ge 2$ (Theorem 4.1; lower bounds floored, upper bounds ceiled at 10^{-6}).

$\gamma = \beta/N^2$	θ_I	$\kappa_0(4,\gamma) \ge$	$\chi_t^{\mathrm{clov}}(\gamma,N) \le$	$\psi_\Lambda(\gamma N^2,\theta_I) - \psi_\Lambda(\gamma N^2,0) \le$
0	1	0.500000	0.184788	0.092394
0	3	0.500000	0.184788	0.831545
1/200	1	0.379999	0.243142	0.121571
1/100	1	0.260000	0.355361	0.177681
1/100	3	0.260000	0.355361	1.599125
1/80	1	0.199999	0.461970	0.230985
1/60	1	0.099999	0.923939	0.461970
1/50	1	0.019999	4.619693	2.309847

Independent Verification and Lean 4.36.0 Benchmark Kernel. Our repository provides a four-part test suite (`test_theta_vacuum.py`, 4/4 passing) and a standalone 92-line Lean 4.36.0 kernel (`ThetaVacuumCerts.lean`):

TABLE 3. Certified 140-digit all-coupling 4D SU(N) Lüscher-admissible clover topological density bounds and imaginary-θ free-energy shifts (15) valid for all $\beta \geq 0$ (Theorem 5.1; upper bounds ceiled at 10^{-6}).

Group	ε	θ_I	$\lvert q_{\mathrm{clov}}(x)\rvert \leq$	$\psi_{\Lambda,\varepsilon}(\beta,\theta_I) - \psi_{\Lambda,\varepsilon}(\beta,0) \leq$
SU(2)	1/100	1	0.000016	0.000016
SU(2)	1/50	1	0.000061	0.000061
SU(2)	1/50	3	0.000061	0.000183
SU(2)	1/35	1	0.000125	0.000125
SU(3)	1/100	1	0.000023	0.000023
SU(3)	1/50	1	0.000092	0.000092
SU(3)	1/50	3	0.000092	0.000274
SU(3)	1/35	1	0.000187	0.000187



(a) Exact 2D U(1) susceptibility ratio **(b) 4D $O(1)$ large-N susceptibility bound**

FIGURE 1. Certified non-perturbative θ-vacuum profiles. (a) Exact 2D U(1) Wilson susceptibility ratio $4\pi^2\beta\,\chi_t^{\mathrm{W}}(\beta)$ approaching the exact Villain limit 1 (Theorem 3.1). (b) Volume- and N-uniform 4D SU(N) Bakry–Émery clover susceptibility bound exhibiting exact $O(1)$ Witten–Veneziano scaling (Theorem 4.1).

- *Direct* 40*-digit quadrature of* 2*D Wilson* θ*-model:* Confirms (7)–(9) and exact Dashen degeneracy $c_0^{\mathrm{W}}(\beta,\pi) = c_1^{\mathrm{W}}(\beta,\pi) < I_0(\beta)$ to 10^{-30}.
- *Direct* 2^4 4*D* SU(2) *and* SU(3) *lattice verification:* Confirms exact pointwise parity oddness $Q_{\mathrm{clov}}(P_0U) = -Q_{\mathrm{clov}}(U)$ to 10^{-12}, the link-gradient bound (12) by central finite differences, and the admissible clover density bound (15).
- *Independent* 80*-digit cross-check and parity:* All 25 rows in Tables 1–3 are independently recomputed at 80-digit precision and matched verbatim across JSON, Lean 4.36.0, and LaTeX.
- *Zero-axiom Lean 4.36.0 benchmark kernel:* The file `ThetaVacuumCerts.lean` compiles cleanly with `-DwarningAsError=true`, 0 warnings, 0 `sorry`, and `#print axioms` reporting *zero axioms* across all five theorems certifying the 25 integer-scaled (10^{12}) inequalities and clover/Bakry–Émery combinatorics (the analytical proofs are given in Sections 2–5).

7. Scope and Limitations

(1) *Clover charge vs. integer topological charge in 4D:* The lattice clover charge $Q_{\mathrm{clov}}(U)$ is smooth and parity-odd on all of $\mathrm{SU}(N)^{|E|}$, making Bakry–Émery Poincaré/Herbst inequalities directly applicable, but unlike the 2D winding charge $Q(U) \in \mathbb{Z}$ or Lüscher's geometric/overlap charge $Q_{\mathrm{ov}} \in \mathbb{Z}$ it is not quantized to integers at finite lattice spacing.

(2) *Strong-coupling window in Theorem 4.1:* The volume- and N-uniform Bakry–Émery bounds (13)–(14) hold on $\gamma = \beta/N^2 < 1/48$. Establishing $O(1)$ continuum topological susceptibility and proving a 4D Dashen transition at $\theta = \pi$ in the continuum limit of $\mathrm{SU}(N)$ Yang–Mills theory remain open problems.

(3) *Real-θ sign problem in 4D:* Theorem 5.1 controls the imaginary-θ free-energy shift across all $\beta \geq 0$ on $\mathcal{U}_\varepsilon$; analytic continuation to real $\theta = \pi$ in 4D requires controlling complex Yang–Lee zeros beyond the exactly solvable 2D models of Sections 2–3.

Data and Code Availability

All verification scripts (`certify_theta_vacuum.py`, `generate_lean.py`), certified 140-digit data (`theta_vacuum_certs.json`), independent tests (`test_theta_vacuum.py`), and the standalone zero-axiom Lean 4.36.0 benchmark kernel (`ThetaVacuumCerts.lean`) are archived online at:

https://airr.science/papers/ARR-2026-5K1DT0HY6G8CEVAH/

References

1. Dominique Bakry and Michel Émery, *Diffusions hypercontractives*, **1123** (1985), 177–206.
2. Dominique Bakry, Ivan Gentil, and Michel Ledoux, *Analysis and geometry of Markov diffusion operators*, Grundlehren der mathematischen Wissenschaften, vol. 348, Springer, 2014.
3. Claudio Bonati and Massimo D'Elia, *Topological critical slowing down: Variations on a toy model*, Physical Review E **98** (2018), no. 1, 013308.
4. Roger Dashen, *Some features of chiral symmetry breaking*, Physical Review D **3** (1971), no. 8, 1879–1889.
5. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Computer Science, vol. 12699, Springer, 2021, pp. 625–635.
6. Christof Gattringer, Thomas Kloiber, and Vasily Sazonov, *Solving the sign problems of the massless lattice Schwinger model with a dual formulation*, Nuclear Physics B **897** (2015), 732–748.
7. Martin Lüscher, *Topology of lattice gauge fields*, Communications in Mathematical Physics **85** (1982), no. 1, 39–48.
8. Cumrun Vafa and Edward Witten, *Parity conservation in quantum chromodynamics*, Physical Review Letters **53** (1984), no. 6, 535–536.
9. Gabriele Veneziano, *$U(1)$ without instantons*, Nuclear Physics B **159** (1979), no. 1–2, 213–224.
10. Ettore Vicari and Haralambos Panagopoulos, *θ dependence of $SU(N)$ gauge theories in the presence of a topological term*, Physics Reports **470** (2009), no. 3–4, 93–150.
11. Uwe-Jens Wiese, *Numerical simulation of lattice θ-vacua: The 2-d $U(1)$ gauge theory as a test case*, Nuclear Physics B **318** (1989), no. 1, 153–175.
12. Kenneth G. Wilson, *Confinement of quarks*, Physical Review D **10** (1974), no. 8, 2445–2459.
13. Edward Witten, *Current algebra theorems for the $U(1)$ "Goldstone boson"*, Nuclear Physics B **156** (1979), no. 2, 269–283.
14. ———, *Large N chiral dynamics*, Annals of Physics **128** (1980), no. 2, 363–375.

Independent Researcher, Östersund, Sweden
Email address: https://airr.science/papers/ARR-2026-5K1DT0HY6G8CEVAH/