Minimax Constant Collapse, Centered Diagonal Sandwiches, and Sharp Binary Lind–Singmaster Counting for Repeated Entries in Pascal's Triangle

Lluis Eriksson*

November 26, 2026

Abstract

We sharpen all three main quantitative pillars of the recent analysis of repeated entries in Pascal's triangle by Leshem and Sayag [1] (`arXiv:2611.34736v1`), where $E_H = \{a \geq 3 : N(a) > 2H\}$ and $A_H(x) = \#(E_H \cap [1, x])$.

First (Theorem 1), we collapse the exponent constant in Leshem and Sayag's global bound $A_H(x) \leq A_2(x) \leq \exp(2\log x/(\log\log x)^{1/4})$ (Proposition 3.7 of [1]) uniformly for *all* $x \geq 3$ and $H \geq 2$ from 2 down to $\frac{113}{100} = 1.13$, and asymptotically to $(\ln 5)^{1/4} = 1.1263375\ldots$ (a 43.68% reduction in the exponent), using only their own ingredients: $J(2,3) = 3$, $J(2,4) = 1$, Rémond's 5^{s^4-64} bound, and exact enumeration of $E_2 \cap [1, 10^6]$. Moreover, we prove the exact minimax theorem $\lim_{x\to\infty} \frac{(\log\log x)^{1/4}}{\log x} \min_{D\geq 4} \log U_{\mathrm{Rem}}(x; D) = (\ln 5)^{1/4}$, showing that $(\ln 5)^{1/4}$ is the unimprovable limit of the Rémond-versus-diagonal trade-off (any $c < (\ln 5)^{1/4}$ causes super-exponential blowup). Independently, excluding the four effectively solved collision pairs $\{(2,3),(2,4),(3,4),(2,5)\}$ halves their effective Bombieri–Pila exponent (Proposition 3.3) for $A_2(x)$ from 1/6 to 1/12 (and improves $A_3(x)$ from 1/12 to 1/15).

Second (Theorem 2), we prove the universal Stirling-centered falling-factorial sandwich $\max\big(m^k(1 - \frac{k(k^2-1)}{24m^2}),\ (m - \frac{3(k^2-1)}{64m})^k\big) \leq (n)_k < m^k$ $(m = n - \frac{k-1}{2})$ for all $k \geq 2$ and $n \geq 2k$. This yields the zero-unknown-constant counting bracket $\lfloor X_k - \frac{3k-1}{2} \rfloor \leq \#(T_k \cap [1, x]) \leq \lfloor X_k - \frac{3k-1}{2} + \frac{3(k^2-1)}{64X_k} \rfloor$ $(X_k = (k!x)^{1/k})$ and eliminates the systematic $-\frac{3}{4}(K-1)(K+2)$ bias from the expansion of $A_1(x)$ in Theorem 1.2 of [1] (cutting $K = 4$ truncated-union errors from 15.33–17.91 down to 0.16–0.41 at the decades $\{10^3, 10^4, 10^5, 10^6\}$, and keeping the full-interval error on $[10^3, 10^6]$ inside $[-1.44, +1.52]$, strictly within Theorem 2(c)'s envelope).

Third (Theorem 3), for the Lind–Singmaster Fibonacci family $\mathcal{F} = \{b_i\}_{i\geq 1}$, we prove the exact Lucas identities $5n_i = L_{4i+5} - 1$, $5k_i = L_{4i+3} - 4$, $5(n_i - k_i) = L_{4i+4} + 3$, collapse Leshem and Sayag's entropy constant $c = \frac{\varphi^5}{5} h(\varphi^{-2})$ in $\mathbb{Q}(\sqrt{5})$ to $c = \frac{\varphi^4}{\sqrt{5}} \ln\varphi \approx 1.4750334$, derive the three-term Robbins–Stirling law $\log b_i = c\varphi^{4i} - 2i\ln\varphi - \frac{1}{2}\ln\frac{2\pi\varphi^4}{5} + O(\varphi^{-4i})$, and sharpen Proposition A.1's unspecified $O(1)$ error to the exact binary law $\#(\mathcal{F}\cap[1,x]) - \lfloor \frac{\log\log x - \log c}{4\log\varphi} \rfloor \in \{0, 1\}$ for *every* real $x \geq 3003$, with 100% asymptotic density of exact floor equality on the $\log\log x$ scale. Core algebraic theorems are constructively formalized in Lean 4.35.0 (17/17 theorems, 0 `sorry`, 0 `Classical.choice`, 0 warnings) and cross-checked by an AST-audited zero-float verifier (16/16 SHA-256 certificates, 56/56 pytest tests).

1 Introduction and Summary of Sharp Improvements

Let $N(a) = \#\{(n,k) \in \mathbb{N}^2 : 1 \leq k < n,\ \binom{n}{k} = a\}$ be the multiplicity of an integer $a > 1$ in Pascal's triangle, and let $M(a) = \#\{(n,k) \in \mathbb{N}^2 : 2 \leq k \leq n/2,\ \binom{n}{k} = a\}$ count its nontrivial

*Independent Researcher. Manuscript and figures licensed under CC-BY-4.0; Lean 4 formalization, exact zero-float rational verification scripts, and SHA-256 certificates licensed under MIT. Source repository: `https://github.com/lluiseriksson/pascal-repeated-entries-sharp-density`.

Table 1: Quantitative comparison between Leshem & Sayag [1] (`arXiv:2611.34736v1`) and our sharpened bounds. All rational certificates are verified with zero floating-point arithmetic under protocol `PASCAL-SHARP-DENSITY-1.0`.

Target in [1]	**Leshem–Sayag Bound**	**Our Sharpened Bound**	**Ref. / Cert.**
Prop. 3.7 (A_2, $\forall x \geq 3$)	$\exp(2\log x/(\log\log x)^{1/4})$	$\exp(\frac{113}{100}\log x/(\log\log x)^{1/4})$ (43.5% cut)	Thm. 1(a), `C02`–`C04`
Prop. 3.7 asymptotic	Constant 2	Exact minimax $(\ln 5)^{1/4} \approx 1.1263$ (43.7% cut)	Thm. 1(b), `C05`
Prop. 3.3 ($H=2$)	$A_2(x) \ll_\varepsilon x^{1/6+\varepsilon}$	$A_2(x) \ll_\varepsilon x^{1/12+\varepsilon}$ (2× sharper)	Thm. 1(c), `C06`
Prop. 3.3 ($H=3$)	$A_3(x) \ll_\varepsilon x^{1/12+\varepsilon}$	$A_3(x) \ll_\varepsilon x^{1/15+\varepsilon}$	Thm. 1(c), `C06`
Prop. 4.1 ($\#(T_k)$)	$(k!x)^{1/k} + O_k(1)$	$[\lfloor X_k - \frac{3k-1}{2}\rfloor, \lfloor X_k - \frac{3k-1}{2} + \frac{3(k^2-1)}{64X_k}\rfloor]$	Thm. 2(b), `C07`–`C08`
Thm. 1.2 ($K=4$ union)	Bias 15.3–17.9 on $[10^3, 10^6]$	Centered $+\Delta_4$: $[-1.44, +1.52]$ (< 0.42 at 10^k)	Thm. 2(c), `C09`–`C10`
Prop. A.1 constant c	$\frac{\varphi^5}{5}h(\varphi^{-2})$	$\frac{\varphi^4}{\sqrt{5}}\ln\varphi = (\frac{3}{2}+\frac{7}{10}\sqrt{5})\ln\varphi$	Thm. 3(b), `C12`
Prop. A.1 ($\#(\mathcal{F})$)	$\frac{\log\log x}{4\log\varphi} + O(1)$	$\lfloor\frac{\log\log x - \log c}{4\log\varphi}\rfloor + \{0,1\}$ ($\forall x \geq 3003$)	Thm. 3(c), `C13`–`C15`

left-half representations. Singmaster's conjecture [14, 2, 9, 11] asserts that $N(a) = O(1)$, while the largest known multiplicity is $N(3003) = 8$ and the infinite Fibonacci family of Lind and Singmaster [10, 15] exhibits infinitely many values with $N(a) \geq 6$.

Recently, Leshem and Sayag [1] studied the exceptional sets $E_H = \{a \geq 3 : N(a) > 2H\}$, their counting functions $A_H(x) = \#(E_H \cap [1, x])$, and their Dirichlet series $D_H(s) = \sum_{a \in E_H} a^{-s}$, proving three flagship quantitative results:

1. **Higher exceptional sets ($H \geq 2$):** Using Bombieri–Pila [5], they obtain $A_H(x) \ll_{H,\varepsilon} x^{1/(H(H+1))+\varepsilon}$ (Proposition 3.3), and using Rémond's effective bound for rational points on curves of genus ≥ 2 [13, 4] with cutoff $D = \lfloor(\frac{\log\log x}{2\log 5})^{1/4}\rfloor$, they prove $A_H(x) \leq A_2(x) \leq \exp(\frac{2\log x}{(\log\log x)^{1/4}})$ for all $x \geq 3$ (Proposition 3.7), establishing $\lambda(E_H) = 0$.

2. **First exceptional set ($H = 1$):** Using the crude diagonal enclosure $(n-k+1)^k/k! \leq \binom{n}{k} \leq n^k/k!$, they prove $\#(T_k \cap [1, x]) = (k!x)^{1/k} + O_k(1)$ and deduce $A_1(x) = \sum_{k=2}^{K}(k!x)^{1/k} + O_K(x^{1/(K+1)})$ for every fixed $K \geq 2$ (Theorem 1.2 / Proposition 4.1).

3. **Lind–Singmaster lower bound:** For $\mathcal{F} = \{b_i\}_{i\geq 1}$ with $b_i = \binom{F_{2i+2}F_{2i+3}}{F_{2i}F_{2i+3}}$, they prove $\#(\mathcal{F} \cap [1, x]) = \frac{\log\log x}{4\log\varphi} + O(1)$ with constant $c = \frac{\varphi^5}{5}h(\varphi^{-2})$ (Appendix A, Proposition A.1).

In this paper we sharpen every single one of these three pillars to its exact analytic or minimax limit, summarized in Table 1.

2 Theorem 1: Global Constant Collapse and Minimax Optimality for $A_H(x)$

Recall from [1] that for $K \geq 2$, $P_K(x) := \#\{(n, k) \in \mathbb{N}^2 : K \leq k \leq n/2,\ \binom{n}{k} \leq x\} \leq \sum_{k=K}^{\lfloor\log_2 x\rfloor} kx^{1/k}$, and for $2 \leq r < s$, $J(r, s) := \#\{(u, v) \in \mathbb{N}^2 : u \geq 2r,\ v \geq 2s,\ \binom{u}{r} = \binom{v}{s}\}$. For $(r, s) \notin \{(2, 3), (2, 4)\}$, the curve $F_{r,s}(U, V) = s!(U)_r - r!(V)_s = 0$ is absolutely irreducible of degree s and genus ≥ 2 with coefficients bounded by $M = r!s!$ [4, 1], so Rémond's bound [13, Prop. 10.1] gives

$$J(r, s) \leq \exp\Big(5^{s^4-64}(\log M)\log(\log M + 2)\Big) \leq \exp\big(4s^3\, 5^{s^4-64}\big) \qquad (s \geq 4), \tag{1}$$

since $\log M < 2\log(s!) \leq 2s\log s < 2s^2$ and $\log(\log M + 2) \leq 2\log(2s) \leq 2s$. In Proposition 3.7 of [1], two sources of slack inflated the exponent constant from $(\ln 5)^{1/4} \approx 1.1263$ to 2: (i) discarding the factor $5^{-64} \approx 5.42 \times 10^{-45}$ in (1), and (ii) forcing $5^{D^4} \leq \sqrt{\log x}$ via $D = \lfloor(\frac{\ell}{2\ln 5})^{1/4}\rfloor$, which sacrifices an entire factor of $2^{1/4}$ inside the exponent. Retaining both yields an all-$x \geq 3$ constant within 0.33% of the exact minimax limit.

Theorem 1 (Global constant collapse, minimax optimality, and Bombieri–Pila sharpening)**.**

(a) ***(Uniform all-$x \geq 3$ constant collapse)*** *For every integer $H \geq 2$ and every real $x \geq 3$,*

$$A_H(x) \leq A_2(x) \leq \exp\left(\frac{113}{100}\,\frac{\log x}{(\log\log x)^{1/4}}\right), \tag{2}$$

and asymptotically

$$\limsup_{x\to\infty} \frac{(\log\log x)^{1/4}}{\log x} \log A_2(x) \leq (\ln 5)^{1/4} = 1.1263375343\ldots \tag{3}$$

(a 43.68% reduction relative to the constant 2 of [1, Prop. 3.7]).

(b) ***(Exact minimax optimality of $(\ln 5)^{1/4}$)*** *Define the Rémond-versus-diagonal upper-bound functional $U_{\mathrm{Rem}}(x; D) := \exp(5^{D^4-64}) + x^{1/(D+1)}$ $(D \in \mathbb{Z}_{\geq 4})$. Then*

$$\lim_{x\to\infty} \frac{(\log\log x)^{1/4}}{\log x} \min_{D\geq 4} \log U_{\mathrm{Rem}}(x; D) = (\ln 5)^{1/4}. \tag{4}$$

In particular, any cutoff sequence $D(x)$ achieving $x^{1/(D(x)+1)} \leq \exp(c\log x/(\log\log x)^{1/4})$ with fixed $c < (\ln 5)^{1/4}$ forces $\log U_{\mathrm{Rem}}(x; D(x)) \geq 5^{-64}(\log x)^{1+\delta/2} \gg \log x$ for $\delta = (\ln 5)/c^4 - 1 > 0$ (core algebraic scaling step in Lean 4: `minimax_remond_exceeds_pos`).

(c) ***(Effective Bombieri–Pila exponent sharpening)*** *Incorporating the four completely solved pairs $(2,3), (2,4), (3,4), (2,5)$ [3, 7, 8, 12, 6] improves the effective Bombieri–Pila bound of [1, Prop. 3.3] (without Rémond's theorem) to*

$$A_2(x) \ll_\varepsilon x^{1/12+\varepsilon} \qquad \text{and} \qquad A_3(x) \ll_\varepsilon x^{1/15+\varepsilon} \tag{5}$$

(cutting the A_2 exponent by a factor of 2 from 1/6 to 1/12, and A_3 from 1/12 to 1/15; Lean 4: `bombieri_pila_H2_exponent_halved`, `bombieri_pila_H3_exponent_improved`).

Proof. Write $L := \log x$, $\ell := \log L = \log\log x$, and $T := L/\ell^{1/4}$. Since $E_H \subseteq E_2$ for all $H \geq 2$, it suffices to bound $A_2(x)$ across three exhaustive regimes.

Regime I ($3 \leq x \leq 10^6$). Direct exhaustive enumeration of all left-half binomial coefficients $\binom{n}{k} \leq 10^6$ $(2 \leq k \leq n/2$; Certificate `C02`, with the first five binomial identities also verified in Lean 4 by `first_e2_collisions_exact`) establishes

$$E_2 \cap [1, 10^6] = \{120,\ 210,\ 1540,\ 3003,\ 7140,\ 11628,\ 24310\}.$$

For $3 \leq x < 120$, $A_2(x) = 0$ so (2) is trivial. On $[120, 10^6]$, $L \geq \ln 120 > 4.78$, and $L \mapsto (\log L)^{1/4}/L$ is strictly decreasing on $[e^{1/4}, \infty)$, while $A_2(x)$ is piecewise constant with jumps only at elements of E_2. Hence $\max_{120\leq x\leq 10^6} \frac{\ell^{1/4}\log\max(1, A_2(x))}{L}$ is attained at one of the 7 jump points; Certificate `C02` verifies that its maximum is $< 0.238 < \frac{113}{100}$ (attained at $a = 24310$, $A_2 = 7$).

Regime II ($10^6 \leq x \leq \exp(e^{400})$, so $\ell \in [\ln\ln 10^6, 400] \subset [2.62, 400]$). Every $a \in E_2$ has $M(a) \geq 2$. Either both smallest left-half indices lie in $\{2, 3\}$ (at most $J(2,3) = 3$ values by Avanesov [3]), or a has an index $k \geq 4$. Hence $A_2(x) \leq 3 + P_4(x)$. Separating $k = 4$ from $k \geq 5$ in Lemma 2.1 of [1] gives

$$3 + P_4(x) \leq 3 + 4x^{1/4} + (\log_2 x)^2 x^{1/5} \leq \left(7 + (L/\ln 2)^2 e^{-L/20}\right) e^{L/4}.$$

Because $L \mapsto L^2 e^{-L/20}$ attains its unique global maximum on $(0, \infty)$ at $L = 40$ with value $1600e^{-2}$, the prefactor is uniformly bounded by $7 + (40/\ln 2)^2 e^{-2} < 460$ for all $L > 0$, and for $\ell \geq 10$ $(L > 22026)$ by 8 (since $(L/20)^4/24 > (L/\ln 2)^2$). Partitioning $[2.62, 400]$ into 8 contiguous

subintervals $[a_j, b_j]$ and bounding $\ell^{1/4} \le b_j^{1/4}$ and $L \ge e^{a_j}$ on each subinterval (Certificate `C03`) proves $\frac{\ell^{1/4}\log(3+P_4(x))}{L} < \frac{113}{100}$ throughout $[2.62, 400]$ (reaching $400^{1/4}/4 + o(1) \approx 1.1180$ at $\ell = 400$).

Regime III ($\ell \ge 400$). For any integer $D \ge 4$, using $J(2,3) = 3$, $J(2,4) = 1$, and (1) across the $\binom{D-1}{2} - 2 < D^2/2$ remaining pairs yields $A_2(x) \le C_D + P_{D+1}(x)$ with

$$C_D \le 4 + \frac{D^2}{2}\exp(4D^3\, 5^{D^4-64}), \qquad P_{D+1}(x) \le (\log_2 x)^2 \exp\!\left(\frac{L}{D+1}\right). \tag{6}$$

For $\ell \in [400, 105400]$, Certificate `C04` verifies 12 contiguous handoff intervals $[\ell_D, \ell'_D]$ for $D = 4, 5, \ldots, 15$. Because $\ell \mapsto \ell - \frac{1}{4}\ln\ell - \frac{1}{2}$ and $\ell \mapsto \frac{\ell^{1/4}}{D+1}$ are both strictly increasing on $[400, \infty)$, certifying $(D^4-64)\ln 5 + \ln(4D^3) \le \ell_D - \frac{1}{4}\ln\ell_D - \frac{1}{2}$ at the lower endpoint ℓ_D (giving $4D^3\, 5^{D^4-64} \le e^{-1/2}T < 0.61T$) and $\frac{(\ell'_D)^{1/4}}{D+1} + 10^{-4} < \frac{113}{100}$ at the upper endpoint ℓ'_D (with $\frac{\ell_D^{1/4}}{D+1} > 0.89 > e^{-1/2}$) certifies the entire interval $[\ell_D, \ell'_D]$; since $\log(4 + D^2/2 + (L/\ln 2)^2) < 10^{-150}T$ for $\ell \ge 400$, we obtain $A_2(x) \le \exp(\frac{113}{100}T)$.

For $\ell \ge 105400$, choose the optimal cutoff

$$D(\ell) := \left\lfloor \left(\frac{\ell - \ln\ell + 60\ln 5}{\ln 5}\right)^{1/4} \right\rfloor \ge 16. \tag{7}$$

Since $60\ln 5 < 97 < 105400 \le \ell$, we have $\ell - \ln\ell + 60\ln 5 < 2\ell$, so $D^4 < 2\ell/\ln 5$ and $\ln(4D^3) < \ln 4 + \frac{3}{4}\ln(2\ell/\ln 5)$. Combining with $D^4 \ln 5 \le \ell - \ln\ell + 60\ln 5$ yields

$$(D^4 - 64)\ln 5 + \ln(4D^3) < \ell - \frac{1}{4}\ln\ell + \left(\ln 4 - 4\ln 5 + \frac{3}{4}\ln\frac{2}{\ln 5}\right) < \ell - \frac{1}{4}\ln\ell - 4, \tag{8}$$

because $\ln 4 - 4\ln 5 + \frac{3}{4}\ln(2/\ln 5) < -4.888 < -4$ (Certificate `C04`). Hence $4D^3\, 5^{D^4-64} < e^{-4}T < 0.02T < \frac{\ell^{1/4}}{D+1}T$, while $\log(4 + D^2/2 + (L/\ln 2)^2) < 10^{-150}T$. Meanwhile $D + 1 > (\frac{\ell - \ln\ell + 60\ln 5}{\ln 5})^{1/4}$, so

$$\frac{\ell^{1/4}}{D+1} < (\ln 5)^{1/4}\left(\frac{\ell}{\ell - \ln\ell + 60\ln 5}\right)^{1/4}. \tag{9}$$

For $105400 \le \ell \le e^{60\ln 5}$ we have $\ln\ell \le 60\ln 5$ and hence $\frac{\ell^{1/4}}{D+1} < (\ln 5)^{1/4}$; for $\ell > e^{60\ln 5} > 10^{41}$ we have $\ln\ell/\ell < 10^{-39}$, so $\frac{\ell^{1/4}}{D+1} < (\ln 5)^{1/4}(1 + 10^{-39}) < 1.12634 < \frac{113}{100}$. As $\ell \to \infty$ the right-hand side converges to $(\ln 5)^{1/4}$, proving both (2) and (3).

For (b), the upper bound $\limsup \le (\ln 5)^{1/4}$ follows from (7). Conversely, fix any $c < (\ln 5)^{1/4}$ and put $\delta := (\ln 5)/c^4 - 1 > 0$. If $D \ge 4$ satisfies $\log(x^{1/(D+1)}) \le cT$, then $D + 1 \ge \ell^{1/4}/c$, so $D \ge (\ell^{1/4}/c)(1 - c\ell^{-1/4})$. For all sufficiently large ℓ, $(1 - c\ell^{-1/4})^4(1 + \delta) \ge 1 + \delta/2$, whence $D^4 \ln 5 \ge (1 + \delta/2)\ell$ (scaling algebra verified in Lean 4 by `minimax_remond_excess_pos`) and therefore $\log U_{\mathrm{Rem}}(x; D) \ge 5^{D^4-64} \ge 5^{-64}e^{(1+\delta/2)\ell} = 5^{-64}L^{1+\delta/2} \gg T$. Thus $\liminf \ge (\ln 5)^{1/4}$, proving (4).

For (c), any $a \in E_H \cap [1, x]$ has H distinct left-half indices $2 \le k_1 < \cdots < k_H$ with $k_{H-1} \ge H$ and $k_H \ge H + 1$. Set $D_2 = 11$ for $H = 2$ and $D_3 = 14$ for $H = 3$. If $k_H > D_H$, then a is counted by $P_{D_H+1}(x) \ll x^{1/(D_H+1)}$ ($x^{1/12}$ for $H = 2$, $x^{1/15}$ for $H = 3$). Otherwise (k_{H-1}, k_H) is a pair $H \le r < s \le D_H$. Removing the four completely solved finite pairs $\{(2,3), (2,4), (3,4), (2,5)\}$ (contributing $O(1)$ elements in total), every remaining pair satisfies $rs \ge 12$ for $H = 2$ and $rs \ge 15$ for $H = 3$ (Lean 4: `bombieri_pila_H2_exponent_halved`, `bombieri_pila_H3_exponent_improved`), so Proposition 3.2 of [1] yields (5). □

Figure 1 displays the global collapse $2 \to 1.13$ across $\ell = \log\log x$ and the razor-sharp minimax cliff at $(\ln 5)^{1/4}$.



Figure 1: Theorem 1. (a) Normalized exponent $\ell^{1/4}\log U(x;D^*)/\log x$ across $\ell = \log\log x$: Leshem–Sayag's constant 2 vs. our optimal cutoff $D^*(\ell)$, staying uniformly below 1.13 for all $x \geq 3$ and converging to $(\ln 5)^{1/4} \approx 1.1263$ (Certificates `C02`–`C04`). (b) Exact minimax phase cliff at $c = (\ln 5)^{1/4}$ for $\ell = 10^4$: any $c < (\ln 5)^{1/4}$ triggers super-exponential blowup of C_D^{Rem} (Certificate `C05`).

3 Theorem 2: Stirling-Centered Falling-Factorial Sandwich and Bias-Free $A_1(x)$ Expansion

Leshem and Sayag's Proposition 4.1 estimates $\#(T_k \cap [1,x]) = (k!x)^{1/k} + O_k(1)$ via $(n-k+1)^k \leq (n)_k \leq n^k$, which leaves an $O(k)$ width per diagonal and accumulates an unmodeled $-\frac{3}{4}(K-1)(K+2)$ systematic bias when summed over $2 \leq k \leq K$. Centering the falling factorial $(n)_k = n(n-1)\cdots(n-k+1)$ at its exact midpoint $m := n - \frac{k-1}{2}$ collapses the $O(k)$ uncertainty to an explicit $O(k^2/X_k)$ bracket valid for *all* $k \geq 2$ and $n \geq 2k$.

Theorem 2 (Stirling-centered falling-factorial sandwich and bias-free $A_1(x)$ expansion)**.** *Let $k \geq 2$ be an integer, and for $n \geq 2k$ set $m := n - \frac{k-1}{2} \geq \frac{3k+1}{2}$.*

(a) ***(Universal two-sided centered sandwich)*** *For every $k \geq 2$ and every integer $n \geq 2k$,*

$$\max\left(m^k\left(1 - \frac{k(k^2-1)}{24m^2}\right),\ \left(m - \frac{3(k^2-1)}{64m}\right)^k\right) \leq (n)_k < m^k. \tag{10}$$

(The strict upper bound is formalized constructively in Lean 4 as `fallFact_strict_upper_centered`, together with the variance identity `three_mul_sumCenteredSq_exact` and scaled Weierstrass bound `weierstrass_prod_lower_int`.)

(b) ***(Zero-unknown-constant diagonal counting bracket)*** *For every $k \geq 2$ and every real $x \geq \binom{2k}{k}$, letting $X_k := (k!x)^{1/k}$,*

$$\left\lfloor X_k - \frac{3k-1}{2} \right\rfloor \leq \#(T_k \cap [1,x]) \leq \left\lfloor X_k - \frac{3k-1}{2} + \frac{3(k^2-1)}{64X_k} \right\rfloor. \tag{11}$$

(c) ***(Bias-free truncated expansion of $A_1(x)$)*** *For $K \geq 2$, let $\Delta_K(x) := \sum_{a \leq x}(m_K(a)-1)_+$ be the exact multiplicity overcount among $T_2, \ldots, T_K$ (where $m_K(a) = \#\{2 \leq k \leq K : a \in T_k\}$), and set $\widehat{A}_{1,K}(x) := \sum_{k=2}^{K}(k!x)^{1/k} - \frac{3}{4}(K-1)(K+2) - \Delta_K(x)$. Then for all $x \geq \binom{2K}{K}$,*

$$-\frac{K-1}{2} \leq \#\left(\bigcup_{k=2}^{K} T_k \cap [1,x]\right) - \widehat{A}_{1,K}(x) \leq \frac{K-1}{2} + \sum_{k=2}^{K}\frac{3(k^2-1)}{64(k!x)^{1/k}}, \tag{12}$$

and the remainder $R_K(x) := A_1(x) - \sum_{k=2}^{K}(k!x)^{1/k}$ *in [1, Thm. 1.2] satisfies* $R_K(x) = ((K+1)!x)^{1/(K+1)} + O_K(x^{1/(K+2)})$ *(so the exponent* $1/(K+1)$ *and leading coefficient* $((K+1)!)^{1/(K+1)}$ *are sharp).*

Proof. For (a), pair symmetric factors in $(n)_k$ around $m = n - \frac{k-1}{2}$: for each $0 \le j < \lfloor k/2 \rfloor$, putting $\delta_j := \frac{k-1}{2} - j > 0$, we have $(n-j)(n-k+1+j) = (m+\delta_j)(m-\delta_j) = m^2 - \delta_j^2 < m^2$ (Lean 4: `four_mul_lt_sq_sum`), while if k is odd the middle factor equals m. Hence

$$(n)_k = m^k \prod_{j=0}^{\lfloor k/2 \rfloor - 1} \left(1 - \frac{\delta_j^2}{m^2}\right) < m^k, \tag{13}$$

where the inequality is strict since $k \ge 2$ ensures at least one factor with $\delta_0 = \frac{k-1}{2} > 0$ (Lean 4: `fallFact_strict_upper_centered`). Next, summing squared offsets gives

$$\sum_{j=0}^{\lfloor k/2 \rfloor - 1} \delta_j^2 = \frac{1}{2} \sum_{j=0}^{k-1} \left(\frac{k-1}{2} - j\right)^2 = \frac{k(k^2-1)}{24} \tag{14}$$

(Lean 4: `three_mul_sumCenteredSq_exact`). Applying Weierstrass's inequality $\prod(1-u_j) \ge 1 - \sum u_j$ for $u_j = \delta_j^2/m^2 \in (0,1)$ (Lean 4: `weierstrass_prod_lower_int`) gives the first lower bound $m^k(1 - \frac{k(k^2-1)}{24m^2}) \le (n)_k$. For the universal $1/k$-th root lower bound, since $n \ge 2k$ implies $2m = 2n - (k-1) \ge 3k+1 > 3(k-1)$, every u_j satisfies $0 < u_j \le (\frac{k-1}{2m})^2 < \frac{1}{9}$. On $[0, 1/9)$, $\ln(1-u) = -\sum_{r=1}^{\infty} \frac{u^r}{r} \ge -\frac{u}{1-u} > -\frac{9}{8}u$. Taking logarithms in (13), dividing by k, and using $e^{-v} \ge 1 - v$ yields

$$\frac{((n)_k)^{1/k}}{m} \ge \exp\left(-\frac{9}{8km^2} \sum_{j=0}^{\lfloor k/2 \rfloor - 1} \delta_j^2\right) = \exp\left(-\frac{3(k^2-1)}{64m^2}\right) \ge 1 - \frac{3(k^2-1)}{64m^2},$$

and multiplying by m and raising to the k-th power proves (10).

For (b), let $n_{\max} := \max\{n \ge 2k : \binom{n}{k} \le x\}$, so $\#(T_k \cap [1,x]) = n_{\max} - 2k + 1$. If $n_* := \lfloor X_k + \frac{k-1}{2} \rfloor \ge 2k$, then $m_* = n_* - \frac{k-1}{2} \le X_k$, so $(n_*)_k < m_*^k \le X_k^k = k!x$ by (a), giving $n_{\max} \ge n_*$ and hence $\#(T_k \cap [1,x]) \ge \lfloor X_k - \frac{3k-1}{2} \rfloor$. If $n_* < 2k$, then $\lfloor X_k - \frac{3k-1}{2} \rfloor \le 0 < 1 \le \#(T_k \cap [1,x])$ since $x \ge \binom{2k}{k}$. For the upper bound, $(n_{\max})_k \le X_k^k$ and (a) give $m_{\max} - \frac{3(k^2-1)}{64 m_{\max}} \le X_k$. If $m_{\max} \le X_k$ the upper bound in (11) is trivial; if $m_{\max} > X_k$, then $1/m_{\max} < 1/X_k$ immediately gives $m_{\max} < X_k + \frac{3(k^2-1)}{64X_k}$, hence $n_{\max} \le \lfloor X_k + \frac{k-1}{2} + \frac{3(k^2-1)}{64X_k} \rfloor$ and (11) follows.

For (c), write $\lfloor y \rfloor = y - \frac{1}{2} + \theta(y)$ with $\theta(y) \in (-\frac{1}{2}, \frac{1}{2}]$. Then $\lfloor X_k - \frac{3k-1}{2} \rfloor = X_k - \frac{3k}{2} + \theta_k$. Summing over $2 \le k \le K$ using $\sum_{k=2}^{K} \frac{3k}{2} = \frac{3}{4}(K-1)(K+2)$ (equivalently $\sum_{k=2}^{K}(3k-1) = \frac{1}{2}(K-1)(3K+4)$, Lean 4 `two_mul_sumOffsetNum`) and subtracting the exact overcount $\Delta_K(x) = \sum_{a \le x}(m_K(a) - 1)_+$ yields (12). The leading asymptotic of $R_K(x)$ follows by applying (11) at $k = K+1$ together with $P_{K+2}(x) \ll_K x^{1/(K+2)}$ and $\Delta_{K+1}(x) \ll_{K,\varepsilon} x^{\varepsilon}$. □

At $K = 4$, the exact finite collision set among T_2, T_3, T_4 is $\{120, 210, 1540, 7140\}$ (since $J(2,3) = 3$, $J(2,4) = 1$, $J(3,4) = 0$; Certificate `C10`), so $\Delta_4(x) = 4$ for all $x \ge 7140$. Certificate `C09` confirms that removing $-\frac{27}{2} - \Delta_4(x)$ cuts the truncated-union error from 15.33–17.91 down to 0.16–0.41 at the decade checkpoints $\{10^3, 10^4, 10^5, 10^6\}$, while across all integers $x \in [10^3, 10^6]$ the signed error ranges over $[-1.4375, +1.5150]$ (extremized at $x = 52325$ and $x = 5995$), strictly inside Theorem 2(c)'s envelope $[-1.5,\ 1.5 + \sum_{k=2}^{4} \frac{3(k^2-1)}{64X_k}]$ (which equals $+1.5487$ at $x = 5995$; Figure 2(b)).



Figure 2: Theorem 2. (a) Exact centered ratio $(n)_k/(n-\frac{k-1}{2})^k$ vs. our strict upper bound 1 and universal lower bound $(1-\frac{3(k^2-1)}{64m^2})^k$ for $k=3,5,8$ (Certificate `C07`). (b) Truncated-union error $\#(\bigcup_{k=2}^{4} T_k \cap [1,x]) - \text{approx}$ on $[10^3, 10^6]$: Leshem–Sayag's bare sum carries a persistent -17.5 bias (for $x \geq 7140$), whereas our centered $+\Delta_4$ expansion stays strictly within Theorem 2(c)'s envelope (shaded, extrema $[-1.44, +1.52]$; Certificate `C09`).

4 Theorem 3: Exact Golden-Ratio Collapse and Sharp $\{0,1\}$ Lind–Singmaster Law

Recall the Lind–Singmaster Fibonacci family $\mathcal{F} = \{b_i : i \geq 1\} \subseteq E_2$ [10, 15], where $b_i := \binom{n_i}{k_i} = \binom{n_i-1}{k_i+1}$ with $n_i := F_{2i+2}F_{2i+3}$, $k_i := F_{2i}F_{2i+3}$, and $m_i := n_i - k_i = F_{2i+1}F_{2i+3}$. Let $L_m = \varphi^m + (-\varphi)^{-m}$ denote the Lucas numbers $(L_0 = 2, L_1 = 1)$ and $\varphi = (1+\sqrt{5})/2$.

Theorem 3 (Exact Lucas identities, constant collapse, and sharp $\{0,1\}$ Lind–Singmaster counting law)**.**

(a) ***(Exact Lucas identities and Cassini binomial equality)*** *For every integer $i \geq 1$,*

$$5n_i = L_{4i+5} - 1, \qquad 5k_i = L_{4i+3} - 4, \qquad 5m_i = L_{4i+4} + 3, \tag{15}$$

and $n_i(k_i+1) = m_i(m_i-1)$, certifying $\binom{n_i}{k_i} = \binom{n_i-1}{k_i+1}$ (Lean 4: `lind_singmaster_three_lucas_identities`, `lind_singmaster_binomial_ratio`).

(b) ***(Exact $\mathbb{Q}(\sqrt{5})$ collapse of the entropy constant)*** *Leshem and Sayag's constant $c = \frac{\varphi^5}{5} h(\varphi^{-2})$ [1, Prop. A.1] (where $h(t) = -t\ln t - (1-t)\ln(1-t)$) simplifies identically to*

$$c = \frac{\varphi^4}{\sqrt{5}} \ln \varphi = \left(\frac{3}{2} + \frac{7}{10}\sqrt{5}\right) \ln \varphi = 1.4750333842\ldots \tag{16}$$

(c) ***(Three-term Robbins–Stirling expansion and sharp $\{0,1\}$ counting law)*** *For every $i \geq 1$,*

$$\log b_i = c\,\varphi^{4i} - 2i \ln \varphi - \frac{1}{2} \ln\left(\frac{2\pi\varphi^4}{5}\right) + O(\varphi^{-4i}). \tag{17}$$

Define $\delta_i := 4i\ln\varphi + \ln c - \ln\ln b_i$. Then $0 < \delta_i < 4\ln\varphi$ holds for every *$i \geq 1$, with $\delta_1 \approx 0.233166$ and $\delta_i = \frac{2i\ln\varphi}{c\varphi^{4i}}(1+O(i^{-1})) \to 0^+$. Consequently, for* every *real $x \geq b_1 = 3003$,*

$$\#(\mathcal{F} \cap [1,x]) - \left\lfloor \frac{\log\log x - \log c}{4\log\varphi} \right\rfloor \in \{0,1\} \tag{18}$$

(Lean 4: `binary_counting_discrepancy_zero_one`). Moreover, the discrepancy equals 1 on a fraction $\mu_i := \frac{\delta_i}{4\ln\varphi} = \frac{i}{2c\varphi^{4i}}(1+O(i^{-1}))$ of the i-th $\log\log x$ period, so exact floor equality $\#(\mathcal{F}\cap[1,x]) = \lfloor \frac{\log\log x - \log c}{4\log\varphi} \rfloor$ holds with 100% asymptotic density in $\log\log x$.

Proof. For (a), by Binet's formula $F_a = (\varphi^a - (-\varphi)^{-a})/\sqrt{5}$, multiplying $5F_{2i+2}F_{2i+3} = (\varphi^{2i+2} - \varphi^{-(2i+2)})(\varphi^{2i+3}+\varphi^{-(2i+3)}) = \varphi^{4i+5} - \varphi^{-(4i+5)} - (\varphi - \varphi^{-1}) = L_{4i+5} - 1$ since $\varphi - \varphi^{-1} = 1$. Similarly $5F_{2i}F_{2i+3} = L_{4i+3} - (\varphi^3 - \varphi^{-3}) = L_{4i+3} - 4$ and $5F_{2i+1}F_{2i+3} = L_{4i+4} + (\varphi^2 + \varphi^{-2}) = L_{4i+4} + 3$ (also proved purely inductively in Lean 4 via `five_fib_product_lucas_joint`). For the binomial equality, $n_i(k_i+1) - m_i(m_i-1) = F_{2i+3}^2(F_{2i+2}F_{2i} - F_{2i+1}^2) + F_{2i+3}(F_{2i+2} + F_{2i+1}) = F_{2i+3}^2(-1) + F_{2i+3}^2 = 0$ by Cassini's identity (Lean 4: `cassini_even`).

For (b), since $\varphi^{-2} + \varphi^{-1} = 1$, we have $1 - \varphi^{-2} = \varphi^{-1}$, whence

$$h(\varphi^{-2}) = -\varphi^{-2}\ln(\varphi^{-2}) - \varphi^{-1}\ln(\varphi^{-1}) = (2\varphi^{-2} + \varphi^{-1})\ln\varphi = (\varphi^{-2}+1)\ln\varphi = \sqrt{5}\,\varphi^{-1}\ln\varphi,$$

using $\varphi^{-2} + 1 = 3 - \varphi = \frac{5-\sqrt{5}}{2} = \sqrt{5}\,\frac{\sqrt{5}-1}{2} = \sqrt{5}\,\varphi^{-1}$. Multiplying by $\varphi^5/5$ gives $c = \frac{\varphi^4}{\sqrt{5}}\ln\varphi = (\frac{3}{2} + \frac{7}{10}\sqrt{5})\ln\varphi$ (Certificate `C12`).

For (c), let $P_i := \frac{\varphi^{4i+5}}{5}$, $Q_i := \frac{\varphi^{4i+3}}{5}$, $R_i := \frac{\varphi^{4i+4}}{5}$, so $P_i = Q_i + R_i$. By (15) and $L_m = \varphi^m + (-1)^m\varphi^{-m}$, we have exact shifts

$$n_i = P_i - \frac{1}{5} - \frac{\varphi^{-(4i+5)}}{5}, \qquad k_i = Q_i - \frac{4}{5} - \frac{\varphi^{-(4i+3)}}{5}, \qquad m_i = R_i + \frac{3}{5} + \frac{\varphi^{-(4i+4)}}{5}.$$

Applying Robbins's two-sided Stirling inequality $\frac{1}{12m+1} < \ln(m!) - (m\ln m - m + \frac{1}{2}\ln(2\pi m)) < \frac{1}{12m}$ to $b_i = n_i!/(k_i!m_i!)$ and expanding $n_i\ln n_i - k_i\ln k_i - m_i\ln m_i$ around (P_i, Q_i, R_i):

- The leading term is $P_i\ln P_i - Q_i\ln Q_i - R_i\ln R_i = P_i h(\varphi^{-2}) = c\,\varphi^{4i}$.
- The $O(1)$ linear shift term $(\Delta n)\ln P_i - (\Delta k)\ln Q_i - (\Delta m)\ln R_i$ has $\Delta n - \Delta k - \Delta m = -\frac{1}{5} + \frac{4}{5} - \frac{3}{5} = 0$ (canceling all $4i\ln\varphi$ and $\ln 5$ terms!) and leaves exact constant $\frac{-5+12-12}{5}\ln\varphi = -\ln\varphi$.
- The Stirling square-root term $-\frac{1}{2}\ln\frac{2\pi k_i m_i}{n_i} = -\frac{1}{2}\ln\frac{2\pi\varphi^{4i+2}}{5} + O(\varphi^{-4i}) = -2i\ln\varphi - \frac{1}{2}\ln\frac{2\pi\varphi^2}{5} + O(\varphi^{-4i})$.

Combining $-\ln\varphi - \frac{1}{2}\ln\frac{2\pi\varphi^2}{5} = -\frac{1}{2}\ln\frac{2\pi\varphi^4}{5}$ establishes (17) (with explicit $O(\varphi^{-4i})$ residual verified to 6 digits for $i = 1,\ldots,5$ in Certificate `C13`).

To verify $0 < \delta_i < 4\ln\varphi$ for *all* $i \geq 1$: first note that $b_i = \binom{k_i+m_i}{k_i}$ is strictly increasing for all $i \geq 1$ since both $k_i = F_{2i}F_{2i+3}$ and $m_i = F_{2i+1}F_{2i+3}$ strictly increase with i. For $1 \leq i \leq 5$, Certificate `C14` rigorously brackets $\delta_1 \in (0.2331, 0.2332)$, $\delta_2 \in (0.0444, 0.0445)$, $\delta_3 \in (0.00838, 0.00839)$, $\delta_4 \in (0.00151, 0.00152)$, and $\delta_5 \in (0.00026, 0.00027)$, all strictly inside $(0, 4\ln\varphi \approx 1.9248)$. For every $i \geq 6$, $n_i \geq n_6 = F_{14}F_{15} = 229970$, $k_i \geq k_6 = 87840$, $m_i \geq m_6 = 142130$, and $|\Delta n_i|, |\Delta k_i|, |\Delta m_i| < 1$. By Taylor's theorem with Lagrange remainder for the strictly convex function $f(t) = t\ln t$ ($f''(t) = 1/t > 0$) together with Robbins's bounds, the magnitude of the $O(\varphi^{-4i})$ residual in (17) is explicitly bounded for all $i \geq 6$ by the sum of four terms: (i) the φ^{-4i} linear-shift term $\leq (\ln\varphi)/\varphi^{29} < 4.2\times10^{-7}$; (ii) the quadratic Taylor remainder $\leq \frac{1}{2n_6} + \frac{1}{2k_6} + \frac{1}{2m_6} < 1.14\times10^{-5}$; (iii) the Stirling square-root shift $\leq \frac{1}{2}(\frac{1}{n_6-1} + \frac{1}{k_6-1} + \frac{1}{m_6-1}) < 1.14\times10^{-5}$; and (iv) the Robbins remainder $\leq \frac{1}{12n_6} + \frac{1}{12k_6} + \frac{1}{12m_6} < 1.9\times10^{-6}$, totaling $< 2.51\times10^{-5} < 10^{-4}$ (certified in exact rational arithmetic in Certificate `C14`). Consequently $c\varphi^{4i} - 2i\ln\varphi - 1.08 < \log b_i < c\varphi^{4i} - 2i\ln\varphi - 1.07 < c\varphi^{4i}$ for all $i \geq 6$. Since $2i\ln\varphi + 1.08 < \frac{1}{2}c\varphi^{4i}$ at $i = 6$ (Certificate `C14`) and $\varphi^4 > 6$, this holds for all $i \geq 6$, giving $\frac{1}{2}c\varphi^{4i} < \log b_i < c\varphi^{4i}$ and hence $0 < \delta_i < \ln 2 < 0.694 < 4\ln\varphi$. Setting $y := \log\log x - \log c$, $y_i := \log\log b_i - \log c = 4i\ln\varphi - \delta_i \in ((i-1)P, iP)$ with $P := 4\ln\varphi$ directly yields (18) via `binary_counting_discrepancy_zero_one`, and $\delta_i = -\ln(1 - \frac{2i\ln\varphi + O(1)}{c\varphi^{4i}}) = \frac{2i\ln\varphi}{c\varphi^{4i}}(1 + O(i^{-1}))$ yields the period-wise failure measure $\mu_i = \frac{\delta_i}{4\ln\varphi} \to 0$. □

Figure 3 illustrates the exact $\{0,1\}$ step-function law and the super-exponential decay $\mu_i \sim \frac{i}{2c\varphi^{4i}} \to 0$.



Figure 3: Theorem 3. (a) Exact Lind–Singmaster count $\#(\mathcal{F} \cap [1, x])$ vs. $\lfloor g(x) \rfloor$ where $g(x) = (\log\log x - \log c)/(4\log\varphi)$: discrepancy is strictly $\{0, 1\}$-valued for all $x \geq 3003$ (Certificate `C14`). (b) Super-exponential collapse of the period-wise failure measure $\mu_i = \delta_i/(4\log\varphi)$ matching $\frac{i}{2c\varphi^{4i}}$ (Certificate `C15`).

5 Formal Verification and Reproducibility

- **Lean 4.35.0 formalization:** `lean/PascalSharpDensity.lean` compiles cleanly via `lake build` in 2.0 s with 0 errors, 0 warnings, 0 `sorry`, 0 custom axioms, and 0 `Classical.choice` across all 17 audited theorems: `four_mul_eq_sq_sub_sq`, `four_mul_lt_sq_sum`, `fallFact_add_two`, `fallFact_strict_upper_centered`, `three_mul_sumShiftSq`, `three_mul_sumCenteredSq_exact`, `weierstrass_prod_lower_int`, `two_mul_sumOffsetNum`, `cassini_even`, `lind_singmaster_binomial_ratio`, `five_fib_product_lucas_joint`, `lind_singmaster_three_lucas_identities`, `bombieri_pila_H2_exponent_halved`, `bombieri_pila_H3_exponent_improved`, `first_e2_collisions_exact`, `minimax_remond_excess_pos`, and `binary_counting_discrepancy_zero_one`.
- **AST-audited zero-float rational verifier:** `verification/verify_paper101.py` computes all logarithms, exponentials, k-th roots, π, φ, and Robbins–Stirling brackets in exact `Fraction` arithmetic, passes a strict AST zero-float hygiene gate, and emits 16 SHA-256 certificates (`C01`–`C16`) under protocol `PASCAL-SHARP-DENSITY-1.0`.
- **Adversarial pytest suite:** `tests/test_verify_paper101.py` passes 56/56 tests in 0.19 s, including 30 adversarial forgery/mutation checks (wrong Rémond cutoffs, sub-minimax constants $c < (\ln 5)^{1/4}$, excluded solved pairs, out-of-range diagonal thresholds, Cassini perturbation, SHA-256 tampering, and AST float-rejection tests) plus an on-disk certificate digest integrity check.

AI Assistance Disclosure. In accordance with transparent scientific disclosure practices, Google Antigravity assisted with Lean 4 proof engineering, zero-float rational interval verification scripting, figure generation, and LaTeX typesetting. All mathematical definitions, theorems, proofs, formalization checks, and final manuscript claims were reviewed and validated by the sole author, Lluis Eriksson.

A Self-Contained Verification Tables and Certificate Ledger

To enable full verification directly from this manuscript without external files, Table 2 tabulates our zero-float Robbins–Stirling verification of Theorem 3 for $i = 1, \ldots, 5$, and Table 3 lists all 16 SHA-256 certificate digests.

Table 2: Zero-float Robbins–Stirling verification of Theorem 3 for $i = 1, \ldots, 5$ (Certificates `C11`, `C13`–`C15`). Exact constant shift $-\frac{1}{2}\ln(2\pi\varphi^4/5) = -1.07664322\ldots$, period $4\ln\varphi = 1.92484730\ldots$.

i	$n_i = F_{2i+2}F_{2i+3}$	$k_i = F_{2i}F_{2i+3}$	$\log b_i$	$r_i = \log b_i - \text{asymp}$	δ_i	$\mu_i = \delta_i/(4\ln\varphi)$
1	15	5	8.007367	−0.063595	0.233166	0.121135
2	104	39	66.284242	−0.009439	0.044423	0.023079
3	714	272	470.990874	−0.001381	0.008384	0.004356
4	4895	1869	3250.471471	−0.000202	0.001514	0.000787
5	33552	12815	22306.941116	−0.000029	0.000264	0.000137

Table 3: Verified SHA-256 certificate ledger under protocol `PASCAL-SHARP-DENSITY-1.0` (manifest hash `16ea5a22fb4a...ed5`).

ID	Verified Claim	SHA-256 Prefix
`C01`	Zero-float rational brackets for $(\ln 5)^{1/4}, \varphi, \ln\varphi, \pi, c_{\mathrm{LS}}$	`a5cc0a5a5ba4e183`
`C02`	Exhaustive $E_2 \cap [1, 10^6]$ (7 elements, max ratio $< 0.24 < 1.13$)	`a418e74ec1331ca7`
`C03`	Thm. 1(a) medium range $\ell \in [2.62, 400]$ via $3 + P_4(x)$	`ff7e25ccb865142d`
`C04`	Thm. 1(a) Rémond handoffs $D = 4\ldots15 + \ell \geq 105400$ lock	`16e1bc86ad54bffe`
`C05`	Thm. 1(b) exact minimax blowup below $(\ln 5)^{1/4} \approx 1.1263375$	`2256472b42ab7bf4`
`C06`	Thm. 1(c) Bombieri–Pila halving $H = 2$ $(1/6 \to 1/12)$, $H = 3$ $(1/12 \to 1/15)$	`34f38a64d13fdc31`
`C07`	Thm. 2(a) centered falling-factorial sandwich $(2 \leq k \leq 12)$	`07db35ad195c45b4`
`C08`	Thm. 2(b) non-asymptotic diagonal bracket (30 (k, x) pairs)	`64e4a8025c31b12e`
`C09`	Thm. 2(c) $K = 4$ union error: < 0.42 at 10^k, $[-1.44, +1.52]$ on $[10^3, 10^6]$	`693449aca99f7b3c`
`C10`	Exact finite collisions $J(2,3) = 3, J(2,4) = 1, J(3,4) = 0, J(2,5) = 2, \Delta_5(10^6) = 6$	`40ad7063d1f2fc7e`
`C11`	Thm. 3(a) exact Lucas identities & binomial equality $(i = 1\ldots20)$	`18199efc65b2b1bd`
`C12`	Thm. 3(b) exact $\mathbb{Q}(\sqrt{5})$ collapse $c = (\frac{3}{2} + \frac{7}{10}\sqrt{5})\ln\varphi$	`ef470372335e0c59`
`C13`	Thm. 3(c) three-term Robbins–Stirling expansion $(i = 1\ldots5)$	`5dba39ed2f6583fe`
`C14`	Thm. 3(c) sharp $\{0, 1\}$ law ($\forall x \geq 3003$, incl. $i \geq 6$ bound)	`cc3ce46af2b5aef7`
`C15`	Thm. 3(c) super-exponential decay of period failure measure $\mu_i \to 0$	`644088d458d34e45`
`C16`	Exact correspondence ledger to Leshem & Sayag (`arXiv:2611.34736v1`)	`5a155a2aad9d9a90`

References

[1] A. Leshem and E. Sayag, *On the density of repeated entries in Pascal's triangle*, arXiv:2611.34736v1, 25 November 2026.

[2] H. L. Abbott, P. Erdős, and D. Hanson, *On the number of times an integer occurs as a binomial coefficient*, Amer. Math. Monthly **81**, 256–261, 1974.

[3] E. T. Avanesov, *Solution of a problem on figurate numbers*, Acta Arith. **12**, 409–420, 1966/67.

[4] F. Beukers, T. N. Shorey, and R. Tijdeman, *Irreducibility of polynomials and arithmetic progressions with equal products of terms*, Number Theory in Progress, Vol. 1, de Gruyter, Berlin, 11–26, 1999.

[5] E. Bombieri and J. Pila, *The number of integral points on arcs and ovals*, Duke Math. J. **59**, 337–357, 1989.

[6] Y. Bugeaud, M. Mignotte, S. Siksek, M. Stoll, and Sz. Tengely, *Integral points on hyperelliptic curves*, Algebra Number Theory **2**, 859–885, 2008.

[7] B. M. M. de Weger, *A binomial Diophantine equation*, Quart. J. Math. Oxford Ser. (2) **47**, 221–231, 1996.

[8] B. M. M. de Weger, *Equal binomial coefficients: Some elementary considerations*, J. Number Theory **63**, 373–386, 1997.

[9] D. M. Kane, *Improved bounds on the number of ways of expressing t as a binomial coefficient*, Integers **7**, Paper A53, 2007.

[10] D. A. Lind, *The quadratic field* $\mathbb{Q}(\sqrt{5})$ *and a certain Diophantine equation*, Fibonacci Quart. **6**, 86–93, 1968.

[11] K. Matomäki, M. Radziwiłł, X. Shao, T. Tao, and J. Teräväinen, *Singmaster's conjecture in the interior of Pascal's triangle*, Q. J. Math. **73**, 1137–1177, 2022.

[12] Á. Pintér, *A note on the Diophantine equation* $\binom{x}{4} = \binom{y}{2}$, Publ. Math. Debrecen **47**, 411–415, 1995.

[13] G. Rémond, *Nombre de points rationnels des courbes*, Proc. London Math. Soc. (3) **101**, 759–794, 2010.

[14] D. Singmaster, *How often does an integer occur as a binomial coefficient?*, Amer. Math. Monthly **78**, 385–386, 1971.

[15] D. Singmaster, *Repeated binomial coefficients and Fibonacci numbers*, Fibonacci Quart. **13**, 295–298, 1975.