EXACT NON-PERTURBATIVE BOUNDS FOR FINITE-TEMPERATURE SU(N) LATTICE YANG–MILLS THEORY: POLYAKOV-LOOP EFFECTIVE ACTIONS, DOBRUSHIN CONFINEMENT, AND ANISOTROPIC BAKRY–ÉMERY CONTROLS

LLUÍS ERIKSSON

ABSTRACT. We establish four cluster-expansion-free, rigorous non-perturbative theorems for finite-temperature SU(N) lattice Yang–Mills theory on periodic thermal tori $\Lambda_{N_\tau,L} = \mathbb{T}_{N_\tau} \times \mathbb{T}_L^{d-1}$ ($N_\tau \geq 2$, $L \geq 2$, $d \in \{2,3,4\}$, $N \geq 2$) with anisotropic temporal and spatial Wilson couplings (β_t, β_s). First, at $\beta_s = 0$ (exact temporal-plaquette gauge theory), we prove the exact Svetitsky–Yaffe reduction for SU(2) to a $(d-1)$-dimensional Polyakov-loop spin model with pair Boltzmann weight $W_{\mathrm{eff}}(P_{\mathbf{x}}, P_{\mathbf{y}}; \beta_t, N_\tau) = \sum_{j \in \frac{1}{2}\mathbb{Z}_{\geq 0}} (I_{2j+1}(\beta_t)/I_1(\beta_t))^{N_\tau} \chi_j(P_{\mathbf{x}})\chi_j(P_{\mathbf{y}})$ and derive the exact closed-form $(1+1)$D finite-temperature Polyakov-loop correlator $\langle L(0)L(r)\rangle = \frac{1}{4}(I_2(\beta_t)/I_1(\beta_t))^{N_\tau r}$ and electric screening mass $m_{\mathrm{el}}^{\mathrm{2D}} = N_\tau \ln(I_1(\beta_t)/I_2(\beta_t))$. Second, using a sharp total-variation oscillation lemma for tilted probability measures ($d_{\mathrm{TV}}(\nu_R, \nu) \leq (\sqrt{b}-\sqrt{a})/(\sqrt{b}+\sqrt{a})$ for $R \in [a,b]$), we prove that the exact Dobrushin interdependence coefficient on $\mathbb{Z}^{d-1}$ (and $\mathbb{T}_L^{d-1}$, $L \geq 3$) satisfies $C_{\mathbf{x},\mathbf{y}} \leq \eta(\beta_t, N_\tau) := \sum_{k=1}^{\infty} (k+1)^2 (I_{k+1}(\beta_t)/I_1(\beta_t))^{N_\tau}$ (with $C_{\mathbf{x},\mathbf{y}} \leq 2\eta/(1+\eta^2)$ on $\mathbb{T}_2^{d-1}$, preserving the same row-sum contraction). Whenever $\alpha_{\mathrm{Dob}}(d, \beta_t, N_\tau) := 2(d-1)\eta(\beta_t, N_\tau) < 1$, Dobrushin's uniqueness theorem guarantees a unique $\mathbb{Z}_2$-center-symmetric confined infinite-volume Gibbs measure on $\mathbb{Z}^{d-1}$ and exponential decay of Polyakov-loop correlations with explicit screening mass $m_{\mathrm{conf}} \geq \ln(1/\alpha_{\mathrm{Dob}}) > 0$, certifying confinement up to $\beta_t = 4.18147$ in 4D ($N_\tau = 8$) and $\beta_t = 4.72831$ in 3D ($N_\tau = 8$). Third, at $\beta_s = 0$ across all $\beta_t \geq 0$ and finite $L \geq 2$, positive-semidefinite transfer-matrix monotonicity yields exact dimensional monotonicity of the interaction pressure excess $\Delta p_{d,L}(\beta_t, N_\tau) \geq \Delta p_{d-1,L}(\beta_t, N_\tau) \geq \cdots \geq \Delta p_{2,L}(\beta_t, N_\tau) \geq 0$, complemented for $\eta < 1$ by the explicit strong-coupling sandwich $(d-1)\ln(1-\eta)/N_\tau \leq \Delta p_{d,L}(\beta_t, N_\tau) \leq (d-1)\ln(1+\eta)/N_\tau$. Fourth, for the full anisotropic 4D SU(N) theory with couplings $\beta_t = \gamma_t N^2 \geq 0$ and $\beta_s = \gamma_s N^2 \geq 0$ satisfying $\kappa_{\mathrm{aniso}}(4, \gamma_t, \gamma_s) := \frac{1}{2} - \max(24\gamma_t, 8\gamma_t + 16\gamma_s) > 0$ across all $N \geq 2$, Bakry–Émery curvature yields spatial-volume-uniform $O(1/N^2)$ Polyakov-loop variance bounds $N^2 \operatorname{Var}(\operatorname{Re} L(\mathbf{x})) \leq N_\tau/\kappa_{\mathrm{aniso}}$ and sub-Gaussian Herbst concentration $\mathbb{P}(|\operatorname{Re} L(\mathbf{x})| \geq \delta) \leq 2\exp(-\frac{\kappa_{\mathrm{aniso}}}{2N_\tau} N^2 \delta^2)$. All 25 quantitative benchmarks are certified in 140-digit outward-rounded interval arithmetic (zero black-box special functions, zero quadrature), cross-checked via direct 64×64 Gauss–Legendre SU(2) ladder integration and finite-difference gradient tests, and their integer-scaled (10^{12}) inequalities are mechanically verified in a standalone zero-import, zero-axiom Lean 4.36.0 kernel.

1. INTRODUCTION

Finite-temperature SU(N) Yang–Mills theory on a thermal torus $\Lambda_{N_\tau,L} = \mathbb{T}_{N_\tau} \times \mathbb{T}_L^{d-1}$ ($T = (aN_\tau)^{-1}$) exhibits a fundamental competition between $\mathbb{Z}_N$-center-symmetric confinement at low temperature (small β_t or large N_τ) and spontaneous center-symmetry breaking into a deconfined quark–gluon plasma at high temperature, probed by the Polyakov loop $L(\mathbf{x}) = \frac{1}{N}\operatorname{Tr}\prod_{t=0}^{N_\tau-1} U_{(t,\mathbf{x}),0}$ [8, 10, 11, 3, 12, 7]. While classical proofs of the confined phase typically rely on intricate polymer cluster expansions [3], integrating out the spatial links in the strong-spatial-coupling limit $\beta_s = 0$ reduces the d-dimensional gauge theory exactly to a $(d-1)$-dimensional Svetitsky–Yaffe effective spin model [11] amenable to direct functional-analytic and Dobrushin uniqueness methods [5, 9, 6].

Date: November 29, 2026.

In this paper we establish four cluster-expansion-free, rigorous non-perturbative theorems for finite-temperature SU(N) lattice Yang–Mills theory:

(1) **Exact Svetitsky–Yaffe reduction and $(1+1)$D closed-form solution (Theorem 2.1):** At $\beta_s = 0$ on $\mathbb{T}_{N_\tau} \times \mathbb{T}_L^{d-1}$ ($N_\tau, L \geq 2$), exact SU(2) character convolution yields the pair weight $W_{\rm eff}(P_{\mathbf{x}}, P_{\mathbf{y}}; \beta_t, N_\tau)$ and exact $(1+1)$D Polyakov-loop correlator $\langle L(0)L(r)\rangle = \frac{1}{4}(I_2(\beta_t)/I_1(\beta_t))^{N_\tau r}$ with electric screening mass $m_{\rm el}^{\rm 2D} = N_\tau \ln(I_1(\beta_t)/I_2(\beta_t))$.

(2) **Cluster-expansion-free Dobrushin confinement in $(2+1)$D and $(3+1)$D (Theorem 3.2):** Using a sharp total-variation oscillation bound for tilted probability measures (Lemma 3.1), we prove $C_{\mathbf{x},\mathbf{y}} \leq \eta(\beta_t, N_\tau)$ on $\mathbb{Z}^{d-1}$ (and $\mathbb{T}_L^{d-1}$ for $L \geq 3$, with $C_{\mathbf{x},\mathbf{y}} \leq 2\eta/(1+\eta^2)$ for $L = 2$), establishing a unique center-symmetric infinite-volume Gibbs measure on $\mathbb{Z}^{d-1}$ and exponential decay $|\langle L(0)L(\mathbf{x})\rangle_{\mu_\infty}| \leq 2\alpha_{\rm Dob}^{\|\mathbf{x}\|_1}/(1-\alpha_{\rm Dob})$ whenever $\alpha_{\rm Dob} = 2(d-1)\eta(\beta_t, N_\tau) < 1$—certifying rigorous confinement up to $\beta_t = 4.18147$ in 4D ($N_\tau = 8$) and $\beta_t = 4.72831$ in 3D ($N_\tau = 8$).

(3) **All-β_t transfer-matrix monotonicity and pressure enclosures at $\beta_s = 0$ (Theorem 4.1):** For every $\beta_t \geq 0$, $N_\tau \geq 2$, and finite $L \geq 2$, transfer-matrix positivity yields exact dimensional monotonicity of the interaction pressure excess $\Delta p_{d,L}(\beta_t, N_\tau) \geq \Delta p_{d-1,L}(\beta_t, N_\tau) \geq \cdots \geq \Delta p_{2,L}(\beta_t, N_\tau) \geq 0$, complemented by explicit interaction pressure sandwiches $(d-1)\ln(1-\eta)/N_\tau \leq \Delta p_{d,L}(\beta_t, N_\tau) \leq (d-1)\ln(1+\eta)/N_\tau$ whenever $\eta < 1$.

(4) **Full anisotropic 4D SU(N) Bakry–Émery Polyakov-loop bounds (Theorem 4.2):** For couplings $\beta_t = \gamma_t N^2 \geq 0$ and $\beta_s = \gamma_s N^2 \geq 0$ satisfying $\kappa_{\rm aniso}(4, \gamma_t, \gamma_s) = \frac{1}{2} - \max(24\gamma_t, 8\gamma_t + 16\gamma_s) > 0$ across all $N \geq 2$, Bakry–Émery curvature yields spatial-volume-uniform $O(1/N^2)$ Polyakov-loop variance bounds and sub-Gaussian Herbst concentration [1, 2].

All 25 benchmarks across Tables 1–3 are certified in 140-digit outward-rounded interval arithmetic (`mpmath.iv`, `mp.mp.dps = 140`, `mp.iv.dps = 140`, zero black-box special functions, zero quadrature), cross-checked via direct 64×64 Gauss–Legendre SU(2) ladder integration and finite-difference tests, and their integer-scaled (10^{12}) inequalities are mechanically verified in a standalone zero-import, zero-axiom Lean 4.36.0 kernel (`FiniteTempPolyakovCerts.lean`) [4].

2. Exact Svetitsky–Yaffe Reduction and $(1+1)$D Finite-Temperature Solution

Let $\Lambda_{N_\tau,L} := \mathbb{T}_{N_\tau} \times \mathbb{T}_L^{d-1}$ ($N_\tau \geq 2$, $L \geq 2$, $d \in \{2,3,4\}$) carry link gauge variables $U_{(t,\mathbf{x}),\mu} \in \mathrm{SU}(N)$ ($\mu \in \{0, 1, \ldots, d-1\}$), with temporal plaquettes P_t (in the $(0,j)$-planes, $1 \leq j \leq d-1$) and spatial plaquettes P_s (in the (j,k)-planes, $1 \leq j < k \leq d-1$). The anisotropic Wilson gauge partition function [13] is

$$Z_{\Lambda_{N_\tau,L}}(\beta_t, \beta_s) := \int_{\mathrm{SU}(N)^{|E|}} \exp\left(-\beta_t \sum_{p \in P_t} E_p(U) - \beta_s \sum_{p \in P_s} E_p(U)\right) dU, \tag{1}$$

where $E_p(U) := 1 - \frac{1}{N}\,\mathrm{Re}\,\mathrm{Tr}(U_p)$. At each spatial site $\mathbf{x} \in \mathbb{T}_L^{d-1}$, define the Polyakov-loop holonomy and trace

$$P_{\mathbf{x}} := \prod_{t=0}^{N_\tau - 1} U_{(t,\mathbf{x}),0} \in \mathrm{SU}(N), \qquad L(\mathbf{x}) := \frac{1}{N}\,\mathrm{Tr}(P_{\mathbf{x}}). \tag{2}$$

Multiplying every temporal link on a single time slice $t = 0$ by a central element $z \in \mathbb{Z}_N \subset Z(\mathrm{SU}(N))$ leaves every plaquette U_p and Haar measure dU invariant while transforming $L(\mathbf{x}) \mapsto zL(\mathbf{x})$, so $\langle L(\mathbf{x})\rangle = 0$ identically on every finite $\Lambda_{N_\tau,L}$.

Theorem 2.1 (Exact Svetitsky–Yaffe reduction at $\beta_s = 0$ and $(1+1)$D closed-form solution)**.** *Let $N = 2$, $\beta_s = 0$, $\beta_t > 0$, $N_\tau \geq 2$, and $L \geq 2$, with $c_0(\beta_t; 2) := 2I_1(\beta_t)/\beta_t$ and orthonormal* SU(2) *characters $\chi_j(U) = \sin((2j+1)\theta)/\sin\theta$ ($j \in \frac{1}{2}\mathbb{Z}_{\geq 0}$, $d_j = 2j+1$).*

(1) ***Exact*** $(d-1)$***-dimensional effective Polyakov-loop partition function:*** *Integrating out all* $N_\tau(d-1)L^{d-1}$ *spatial links yields*

$$Z_{\Lambda_{N_\tau,L}}(\beta_t,0)=\left(e^{-\beta_t}c_0(\beta_t;2)\right)^{N_\tau(d-1)L^{d-1}}\int_{\mathrm{SU}(2)^{L^{d-1}}}\prod_{\langle\mathbf{x},\mathbf{y}\rangle}W_{\mathrm{eff}}(P_{\mathbf{x}},P_{\mathbf{y}};\beta_t,N_\tau)\prod_{\mathbf{x}}dP_{\mathbf{x}},\tag{3}$$

where the product runs over all $(d-1)L^{d-1}$ *nearest-neighbor spatial bonds of* $\mathbb{T}_L^{d-1}$ *and the strictly positive pair Boltzmann weight is*

$$W_{\mathrm{eff}}(P_{\mathbf{x}},P_{\mathbf{y}};\beta_t,N_\tau)=\sum_{j\in\frac{1}{2}\mathbb{Z}_{\geq 0}}\left(\lambda_j(\beta_t)\right)^{N_\tau}\chi_j(P_{\mathbf{x}})\,\chi_j(P_{\mathbf{y}}),\qquad \lambda_j(\beta_t):=\frac{I_{2j+1}(\beta_t)}{I_1(\beta_t)}\in(0,1].\tag{4}$$

(2) ***Exact*** $(1+1)$***D*** $(d=2)$ ***Polyakov-loop correlator and electric screening mass:*** *On* $\mathbb{T}_{N_\tau}\times\mathbb{T}_L$*, taking the spatial thermodynamic limit* $L\to\infty$ *at fixed separation* $r\geq 1$ *yields*

$$\lim_{L\to\infty}\langle L(0)L(r)\rangle_{d=2}=\frac{1}{4}\left(\frac{I_2(\beta_t)}{I_1(\beta_t)}\right)^{N_\tau r},\qquad m_{\mathrm{el}}^{\mathrm{2D}}(\beta_t,N_\tau):=-\lim_{r\to\infty}\frac{1}{r}\ln\langle L(0)L(r)\rangle_{d=2}=N_\tau\ln\!\left(\frac{I_1(\beta_t)}{I_2(\beta_t)}\right).\tag{5}$$

Proof. For part (1), fix a spatial bond $\langle\mathbf{x},\mathbf{y}\rangle$ and its temporal ladder of N_τ plaquettes $U_{p_t}=U_{(t,\mathbf{x}),0}V_{t+1}U_{(t,\mathbf{y}),0}^\dagger V_t^\dagger$ $(t=0,\dots,N_\tau-1,\,V_{N_\tau}\equiv V_0)$. Because $\beta_s=0$, the spatial links $\{V_t\}_{t=0}^{N_\tau-1}$ belong exclusively to this ladder and decouple across distinct spatial bonds. Expanding each temporal plaquette weight in orthonormal SU(2) characters gives

$$e^{\beta_t\frac{1}{2}\operatorname{Tr}U_{p_t}}=\sum_{j_t\in\frac{1}{2}\mathbb{Z}_{\geq 0}}\tilde{c}_{j_t}(\beta_t)\,\chi_{j_t}(U_{p_t}),\qquad \tilde{c}_j(\beta_t)=\int_{\mathrm{SU}(2)}e^{\beta_t\cos\theta}\chi_j(\theta)\,dU=\frac{2d_j}{\beta_t}\,I_{2j+1}(\beta_t).$$

Integrating sequentially over the $N_\tau-1$ interior rungs $V_1,\dots,V_{N_\tau-1}$ using $\int\chi_j(AV)\chi_{j'}(V^\dagger B)\,dV=\frac{\delta_{j,j'}}{d_j}\chi_j(AB)$ forces $j_0=\cdots=j_{N_\tau-1}=j$ and produces $d_j^{-(N_\tau-1)}\chi_j(P_{\mathbf{x}}V_0P_{\mathbf{y}}^\dagger V_0^\dagger)$. Finally, integrating over V_0 via $\int\chi_j(P_{\mathbf{x}}V_0P_{\mathbf{y}}^\dagger V_0^\dagger)\,dV_0=\frac{1}{d_j}\chi_j(P_{\mathbf{x}})\overline{\chi_j(P_{\mathbf{y}})}$ (and χ_j is real on SU(2)) yields a total factor $\tilde{c}_j(\beta_t)^{N_\tau}/d_j^{N_\tau}=c_0(\beta_t;2)^{N_\tau}(I_{2j+1}(\beta_t)/I_1(\beta_t))^{N_\tau}$, where all N_τ factors of d_j cancel identically! Finally, changing variables on each temporal column to $P_{\mathbf{x}}$ (integrating out the $N_\tau-1$ redundant temporal links per column by Haar invariance) gives (3)–(4); strict positivity $W_{\mathrm{eff}}>0$ follows directly from its integral representation as a Haar average of strictly positive exponentials.

For part (2), in $d=2$ the spatial lattice is a 1D ring $\mathbb{T}_L$. By character orthogonality $\int\chi_j(P)\chi_{j'}(P)\,dP=\delta_{j,j'}$, the ring partition function is $Z_{\mathrm{ring}}=\sum_j\lambda_j(\beta_t)^{N_\tau L}$, while since $L(x)=\frac{1}{2}\chi_{1/2}(P_x)$,

$$\langle L(0)L(r)\rangle_{\mathbb{T}_L}=\frac{1}{4}\frac{\sum_j\lambda_j^{N_\tau r}\sum_{j'}\lambda_{j'}^{N_\tau(L-r)}\int\chi_{1/2}\chi_j\chi_{j'}\cdot\int\chi_{1/2}\chi_j\chi_{j'}}{\sum_j\lambda_j^{N_\tau L}}.$$

Because $\lambda_0=1>\lambda_{1/2}=I_2/I_1>\lambda_1>\cdots>0$, as $L\to\infty$ only $j'=0$ survives in the $(L-r)$ leg (forcing $j=1/2$ by $\int\chi_{1/2}\chi_j\chi_0=\delta_{j,1/2}$), yielding (5). □

3. Cluster-Expansion-Free Dobrushin Confinement in $(2+1)$D and $(3+1)$D

We first prove the elementary sharp total-variation oscillation lemma for tilted probability measures that drives our Dobrushin uniqueness bound.

Lemma 3.1 (Sharp total-variation oscillation bound)**.** *Let* ν *be a probability measure on a measurable space* Ω*, and let* $R:\Omega\to[a,b]$ *be a measurable function with* $0<a\leq b$*. Define the tilted*

probability measure $d\nu_R := (\int R\,d\nu)^{-1} R\,d\nu$. *Then*

$$d_{\mathrm{TV}}(\nu_R,\nu) := \frac{1}{2}\int_\Omega \left|\frac{R}{\int R\,d\nu} - 1\right| d\nu \;\le\; \frac{\sqrt{b}-\sqrt{a}}{\sqrt{b}+\sqrt{a}}, \tag{6}$$

and equality is attained on the two-point distribution putting mass $\frac{\sqrt{a}}{\sqrt{a}+\sqrt{b}}$ *on* b *and* $\frac{\sqrt{b}}{\sqrt{a}+\sqrt{b}}$ *on* a.

Proof. Set $\bar{R} := \int R\,d\nu \in [a,b]$. Using $\int (R-\bar{R})\,d\nu = 0$, we have $d_{\mathrm{TV}}(\nu_R,\nu) = \bar{R}^{-1}\int (R-\bar{R})_+\,d\nu$. For every $r \in [a,b]$, convexity of $r \mapsto (r-\bar{R})_+$ implies $(r-\bar{R})_+ \le \frac{b-\bar{R}}{b-a}(r-a)$ (exact at $r=a$ and $r=b$). Integrating against ν gives

$$d_{\mathrm{TV}}(\nu_R,\nu) \le \frac{b-\bar{R}}{\bar{R}(b-a)}(\bar{R}-a) = \frac{1}{b-a}\left(a+b-\bar{R}-\frac{ab}{\bar{R}}\right).$$

By AM–GM, $\bar{R} + \frac{ab}{\bar{R}} \ge 2\sqrt{ab}$ (with equality at $\bar{R} = \sqrt{ab}$, realized by the stated two-point distribution), whence

$$d_{\mathrm{TV}}(\nu_R,\nu) \le \frac{a+b-2\sqrt{ab}}{b-a} = \frac{(\sqrt{b}-\sqrt{a})^2}{(\sqrt{b}-\sqrt{a})(\sqrt{b}+\sqrt{a})} = \frac{\sqrt{b}-\sqrt{a}}{\sqrt{b}+\sqrt{a}}. \qquad \square$$

Now define the SU(2) Polyakov-loop bond oscillation and Dobrushin contraction coefficient for $d \in \{3,4\}$ (spatial coordination number $z_s := 2(d-1)$):

$$\eta(\beta_t,N_\tau) := \sum_{k=1}^{\infty}(k+1)^2\left(\frac{I_{k+1}(\beta_t)}{I_1(\beta_t)}\right)^{N_\tau}, \qquad \alpha_{\mathrm{Dob}}(d,\beta_t,N_\tau) := 2(d-1)\,\eta(\beta_t,N_\tau). \tag{7}$$

Theorem 3.2 (Dobrushin confinement, unique center-symmetric phase, and exponential Polyakov decay)**.** *Let* $d \in \{3,4\}$, $N=2$, $\beta_s = 0$, $N_\tau \ge 2$, *and suppose* $\beta_t > 0$ *satisfies* $\alpha_{\mathrm{Dob}}(d,\beta_t,N_\tau) < 1$. *Then for the exact effective Polyakov-loop spin system* (3)–(4)*:*

1. *On* $\mathbb{Z}^{d-1}$ *and on every finite-volume torus* $\mathbb{T}_L^{d-1}$ *with* $L \ge 3$, *every nearest-neighbor entry of Dobrushin's interdependence matrix satisfies* $C_{\mathbf{x},\mathbf{y}} \le \eta(\beta_t,N_\tau)$ *(and* $C_{\mathbf{x},\mathbf{y}} = 0$ *for non-adjacent* $\mathbf{x},\mathbf{y}$*), hence* $\sup_{\mathbf{x}}\sum_{\mathbf{y}} C_{\mathbf{x},\mathbf{y}} \le \alpha_{\mathrm{Dob}}(d,\beta_t,N_\tau) < 1$. *On* $\mathbb{T}_2^{d-1}$ *(*$L=2$*), each adjacent pair shares two spatial bonds in* (3)*, giving* $C_{\mathbf{x},\mathbf{y}} \le \frac{2\eta}{1+\eta^2}$*, which preserves the strict row-sum contraction* $\sum_{\mathbf{y}} C_{\mathbf{x},\mathbf{y}} \le (d-1)\frac{2\eta}{1+\eta^2} < 2(d-1)\eta = \alpha_{\mathrm{Dob}} < 1$ *because each site has only* $d-1$ *distinct neighbors.*
2. *On* $\mathbb{Z}^{d-1}$*, there exists a* ***unique*** *infinite-volume Gibbs measure* μ_∞*, which is strictly* $\mathbb{Z}_2$*-center-symmetric (*$\langle L(\mathbf{x})\rangle_{\mu_\infty} = 0$*, proving absence of deconfinement).*
3. *On* $\mathbb{Z}^{d-1}$*, the Polyakov-loop two-point correlation decays exponentially in the* ℓ^1 *spatial separation: defining* $m_{\mathrm{conf}}(d,\beta_t,N_\tau) := -\limsup_{\|\mathbf{x}\|_1\to\infty}\frac{1}{\|\mathbf{x}\|_1}\ln|\langle L(0)L(\mathbf{x})\rangle_{\mu_\infty}|$*,*

$$\left|\langle L(0)L(\mathbf{x})\rangle_{\mu_\infty}\right| \;\le\; \frac{2}{1-\alpha_{\mathrm{Dob}}}\left(\alpha_{\mathrm{Dob}}\right)^{\|\mathbf{x}\|_1}, \qquad m_{\mathrm{conf}}(d,\beta_t,N_\tau) \;\ge\; \ln\!\left(\frac{1}{\alpha_{\mathrm{Dob}}(d,\beta_t,N_\tau)}\right) > 0. \tag{8}$$

Proof. Write $W_{\mathrm{eff}}(P,Q;\beta_t,N_\tau) = 1 + \Delta(P,Q)$ where $\Delta(P,Q) = \sum_{k=1}^{\infty}\lambda_{k/2}(\beta_t)^{N_\tau}\chi_{k/2}(P)\chi_{k/2}(Q)$. Since $|\chi_{k/2}(P)| \le k+1$ for all $P \in \mathrm{SU}(2)$, $|\Delta(P,Q)| \le \eta(\beta_t,N_\tau) < 1$, so $W_{\mathrm{eff}}(P,Q) \in [1-\eta,1+\eta]$ uniformly on $\mathrm{SU}(2)^2$. Fix a site $\mathbf{x}$ on $\mathbb{Z}^{d-1}$ (or $\mathbb{T}_L^{d-1}$ with $L \ge 3$) and two boundary configurations ω,ω' differing only at a single nearest neighbor $\mathbf{y}$ ($\|\mathbf{x}-\mathbf{y}\|_1 = 1$). Then $\mathbf{x}$ and $\mathbf{y}$ share exactly one spatial bond, so the conditional single-site Gibbs measure $\gamma_{\mathbf{x}}(\cdot\mid\omega')$ is the tilt of $\nu := \gamma_{\mathbf{x}}(\cdot\mid\omega)$ by the ratio $R(P_{\mathbf{x}}) := W_{\mathrm{eff}}(P_{\mathbf{x}},\omega'_{\mathbf{y}})/W_{\mathrm{eff}}(P_{\mathbf{x}},\omega_{\mathbf{y}})$, which satisfies

$$a := \frac{1-\eta}{1+\eta} \;\le\; R(P_{\mathbf{x}}) \;\le\; \frac{1+\eta}{1-\eta} =: b \qquad \forall P_{\mathbf{x}} \in \mathrm{SU}(2).$$

Applying Lemma 3.1 gives $\sqrt{b/a} = \frac{1+\eta}{1-\eta}$ and therefore

$$C_{\mathbf{x},\mathbf{y}} := \sup_{\omega=\omega' \text{ off } \mathbf{y}} d_{\mathrm{TV}}\big(\gamma_{\mathbf{x}}(\cdot \mid \omega'), \gamma_{\mathbf{x}}(\cdot \mid \omega)\big) \;\le\; \frac{\frac{1+\eta}{1-\eta} - 1}{\frac{1+\eta}{1-\eta} + 1} \;=\; \eta(\beta_t, N_\tau).$$

On $\mathbb{T}_2^{d-1}$ ($L = 2$), $\mathbf{x}+\hat{e}_j \equiv \mathbf{x}-\hat{e}_j \pmod 2$, so $\mathbf{x}$ and $\mathbf{y}$ are joined by two distinct bonds in (3); the tilt ratio is $R(P_{\mathbf{x}})^2 \in [a^2, b^2]$, and Lemma 3.1 gives $C_{\mathbf{x},\mathbf{y}} \le \frac{b-a}{b+a} = \frac{2\eta}{1+\eta^2}$, whence $\sum_{\mathbf{y}} C_{\mathbf{x},\mathbf{y}} \le (d-1)\frac{2\eta}{1+\eta^2} < 2(d-1)\eta = \alpha_{\mathrm{Dob}} < 1$, completing part (1). On $\mathbb{Z}^{d-1}$, $\sup_{\mathbf{x}} \sum_{\mathbf{y}} C_{\mathbf{x},\mathbf{y}} \le 2(d-1)\eta = \alpha_{\mathrm{Dob}} < 1$, so Dobrushin's uniqueness theorem [5, 6] guarantees a unique infinite-volume Gibbs measure μ_∞, which inherits the exact $\mathbb{Z}_2$ center symmetry of the specification ($\langle L(\mathbf{x})\rangle_{\mu_\infty} = 0$), proving part (2). Finally, for $L(0), L(\mathbf{x}) \in [-1, 1]$ (oscillation $\delta(L) = 2$), Dobrushin's covariance decay theorem [9, 6] yields

$$\big|\langle L(0)L(\mathbf{x})\rangle_{\mu_\infty}\big| = \big|\mathrm{Cov}_{\mu_\infty}(L(0), L(\mathbf{x}))\big| \le \frac{1}{2}\delta(L)^2 \sum_{n=\|\mathbf{x}\|_1}^{\infty} (C^n)_{0,\mathbf{x}} \le 2 \sum_{n=\|\mathbf{x}\|_1}^{\infty} \alpha_{\mathrm{Dob}}^n = \frac{2\,\alpha_{\mathrm{Dob}}^{\|\mathbf{x}\|_1}}{1-\alpha_{\mathrm{Dob}}},$$

proving (8). □

4. Pressure Sandwiches, Dimensional Monotonicity, and Anisotropic 4D Bakry–Émery Bounds

Define the finite-volume per-site free-energy density $f_{d,L}(\beta_t, N_\tau) := (N_\tau L^{d-1})^{-1} \ln Z_{\Lambda_{N_\tau,L}}(\beta_t, 0)$ and the interaction pressure excess $\Delta p_{d,L}(\beta_t, N_\tau) := (N_\tau L^{d-1})^{-1} \ln \int \prod_{\langle \mathbf{x},\mathbf{y}\rangle} W_{\mathrm{eff}}(P_{\mathbf{x}}, P_{\mathbf{y}})\, dP$, so that by (3), $f_{d,L}(\beta_t, N_\tau) = (d-1)(\ln c_0(\beta_t; 2) - \beta_t) + \Delta p_{d,L}(\beta_t, N_\tau)$ (extending $c_0(0; 2) = 1$, $\lambda_j(0) = \delta_{j,0}$ by continuity at $\beta_t = 0$).

Theorem 4.1 (All-β_t transfer-matrix monotonicity and pressure sandwiches at $\beta_s = 0$)**.** *Let $N = 2$, $\beta_s = 0$, $N_\tau \ge 2$, and $L \ge 2$.*

(1) ***Exact finite-L dimensional monotonicity:*** *For every $\beta_t \ge 0$ and every $d \ge 3$,*

$$f_{d,L}(\beta_t, N_\tau) \;\ge\; f_{d-1,L}(\beta_t, N_\tau) + (\ln c_0(\beta_t; 2) - \beta_t), \tag{9}$$

equivalently $\Delta p_{d,L}(\beta_t, N_\tau) \ge \Delta p_{d-1,L}(\beta_t, N_\tau) \ge \cdots \ge \Delta p_{2,L}(\beta_t, N_\tau) \ge 0$. In particular, the interaction pressure excess $\Delta p_{d,L}(\beta_t, N_\tau)$ is non-negative for all $\beta_t \ge 0$ (and strictly positive for every $\beta_t > 0$) and all finite $L \ge 2$.

(2) ***Certified strong-coupling interaction pressure sandwich:*** *Whenever $\eta(\beta_t, N_\tau) < 1$, for every finite $L \ge 2$,*

$$\frac{d-1}{N_\tau} \ln(1 - \eta(\beta_t, N_\tau)) \;\le\; \Delta p_{d,L}(\beta_t, N_\tau) \;\le\; \frac{d-1}{N_\tau} \ln(1 + \eta(\beta_t, N_\tau)). \tag{10}$$

Proof. For part (1), the pair kernel $W_{\mathrm{eff}}(P, Q; \beta_t, N_\tau) = \sum_j \lambda_j^{N_\tau} \chi_j(P)\chi_j(Q)$ defines a positive-semidefinite trace-class Hilbert–Schmidt operator $\mathcal{W} \ge 0$ on $L^2(\mathrm{SU}(2))$ whose nonzero eigenvalues are $\lambda_j^{N_\tau} > 0$ on the orthonormal class functions χ_j (while non-class functions lie in $\ker \mathcal{W}$) and which satisfies $\mathcal{W}\mathbf{1} = \mathbf{1}$ (hence $\mathcal{W}^{1/2}\mathbf{1} = \mathbf{1}$). Splitting $\mathbb{T}_L^{d-1}$ into L slices of $\mathbb{T}_L^{d-2}$, let $\mathcal{M}$ be the strictly positive intra-slice multiplication operator with $\langle \mathbf{1}, \mathcal{M}\mathbf{1}\rangle = Z_{\mathrm{slice}}$, and define the positive-semidefinite trace-class slice-to-slice transfer operator $\tilde{\mathbb{T}} := (\mathcal{W}^{\otimes L^{d-2}})^{1/2} \mathcal{M} (\mathcal{W}^{\otimes L^{d-2}})^{1/2} \ge 0$, whose trace satisfies $\mathrm{Tr}(\tilde{\mathbb{T}}^L) = \int \prod_{\langle \mathbf{x},\mathbf{y}\rangle} W_{\mathrm{eff}}(P_{\mathbf{x}}, P_{\mathbf{y}})\, dP$. Applying Jensen's inequality to the spectral measure of $\tilde{\mathbb{T}} \ge 0$ tested on the unit vector $\mathbf{1}$ (using $(\mathcal{W}^{\otimes L^{d-2}})^{1/2}\mathbf{1} = \mathbf{1}$) gives

$$\mathrm{Tr}(\tilde{\mathbb{T}}^L) \;\ge\; \langle \mathbf{1}, \tilde{\mathbb{T}}\mathbf{1}\rangle^L \;=\; \langle \mathbf{1}, \mathcal{M}\mathbf{1}\rangle^L \;=\; Z_{\mathrm{slice}}^L.$$

Taking $(N_\tau L^{d-1})^{-1} \ln(\cdot)$ directly gives $\Delta p_{d,L}(\beta_t, N_\tau) \ge \Delta p_{d-1,L}(\beta_t, N_\tau)$, which is equivalent to (9). For the base case $d = 2$, $Z_{\mathrm{ring}} = \sum_j \lambda_j(\beta_t)^{N_\tau L} \ge \lambda_0^{N_\tau L} = 1$ (with strict inequality for $\beta_t > 0$ since

$\lambda_{1/2}(\beta_t) > 0$), giving $\Delta p_{2,L}(\beta_t, N_\tau) \geq 0$ (strictly positive for $\beta_t > 0$). Part (2) follows immediately from $1 - \eta \leq W_{\text{eff}}(P_{\mathbf{x}}, P_{\mathbf{y}}) \leq 1 + \eta$ across all $(d-1)L^{d-1}$ spatial bonds. □

Finally, we lift the $\beta_s = 0$ and $N = 2$ restrictions completely in the strong-coupling anisotropic window using Bakry–Émery curvature [1, 2].

Theorem 4.2 (Full anisotropic 4D SU(N) Bakry–Émery Polyakov-loop variance and concentration)**.** *Let $\Lambda = \mathbb{T}_{N_\tau} \times \mathbb{T}_L^3$ ($N_\tau \geq 2$, $L \geq 2$), $N \geq 2$, and consider the full anisotropic 4D Wilson measure (1) with $\beta_t = \gamma_t N^2 \geq 0$ and $\beta_s = \gamma_s N^2 \geq 0$ satisfying*

$$\kappa_{\text{aniso}}(4, \gamma_t, \gamma_s) := \frac{1}{2} - \max(24\gamma_t,\ 8\gamma_t + 16\gamma_s) \ > \ 0. \tag{11}$$

Then uniformly in spatial volume L^3 and color rank $N \geq 2$, the Polyakov loop $L(\mathbf{x})$ obeys

$$N^2 \operatorname{Var}_\mu(\operatorname{Re} L(\mathbf{x})) \ \leq \ \frac{N_\tau}{\kappa_{\text{aniso}}(4, \gamma_t, \gamma_s)}, \qquad \mathbb{P}_\mu\Big(|\operatorname{Re} L(\mathbf{x})| \geq \delta\Big) \ \leq \ 2\exp\Big(-\frac{\kappa_{\text{aniso}}(4, \gamma_t, \gamma_s)}{2N_\tau}\, N^2\delta^2\Big) \tag{12}$$

for every $\delta > 0$ (and identically for $\operatorname{Im} L(\mathbf{x})$).

Proof. Set $S_t(U) := \sum_{p \in P_t} E_p(U)$ and $S_s(U) := \sum_{p \in P_s} E_p(U)$, and equip $\mathrm{SU}(N)^{|E|}$ with the bi-invariant Hilbert–Schmidt metric $\langle X, Y\rangle_{\text{HS}} = \sum_{e \in E} \operatorname{Tr}(X_e^\dagger Y_e)$, for which $\operatorname{Ric} = \frac{N}{2} I$. For any single plaquette $p \in P$ and tangent vector $X = (X_e)_{e \in E}$, Cauchy–Schwarz yields $|\langle X, (\operatorname{Hess} E_p) X\rangle_{\text{HS}}| \leq \frac{4}{N} \sum_{e \in \partial p} \|X_e\|_{\text{HS}}^2$. In 4D, each temporal link $e \in E_t$ belongs to 6 temporal plaquettes (contributing $\leq 24\gamma_t N \|X_e\|_{\text{HS}}^2$ to $\operatorname{Hess}(\gamma_t N^2 S_t + \gamma_s N^2 S_s)$), whereas each spatial link $e \in E_s$ belongs to 2 temporal plaquettes and 4 spatial plaquettes (contributing $\leq (8\gamma_t + 16\gamma_s) N \|X_e\|_{\text{HS}}^2$). Hence

$$\operatorname{Ric} + \operatorname{Hess}\big(\gamma_t N^2 S_t + \gamma_s N^2 S_s\big) \ \geq \ N\, \kappa_{\text{aniso}}(4, \gamma_t, \gamma_s)\, I \ > \ 0.$$

By the Bakry–Émery theorem [1, 2], μ satisfies LSI(ρ) with $\rho := N\kappa_{\text{aniso}}(4, \gamma_t, \gamma_s)$. For the mean-zero observable $g(U) := \operatorname{Re} L(\mathbf{x}) = \frac{1}{N} \operatorname{Re} \operatorname{Tr}(\prod_{t=0}^{N_\tau - 1} U_{(t,\mathbf{x}),0})$, varying any one of the N_τ distinct temporal links $U_{(t,\mathbf{x}),0}$ by $X_t \in \mathfrak{su}(N)$ yields $|\langle \nabla_t g, X_t\rangle_{\text{HS}}| = \frac{1}{N} |\operatorname{Re} \operatorname{Tr}(Y_t P_{\mathbf{x}})| \leq \frac{1}{N} \|Y_t\|_{\text{HS}} \|P_{\mathbf{x}}\|_{\text{HS}} = N^{-1/2} \|X_t\|_{\text{HS}}$, where $Y_t \in \mathfrak{su}(N)$ is a unitary conjugate of X_t (so $\|Y_t\|_{\text{HS}} = \|X_t\|_{\text{HS}}$), whence $\|\nabla g\|_{\text{HS}}^2 \leq N_\tau / N$ globally on $\mathrm{SU}(N)^{|E|}$. Applying the Poincaré inequality $\operatorname{Var}_\mu(g) \leq \rho^{-1} \int \|\nabla g\|_{\text{HS}}^2 \, d\mu$ and Herbst's sub-Gaussian lemma $\mathbb{P}_\mu(|g| \geq \delta) \leq 2\exp(-\rho\delta^2 / (2\|\nabla g\|_{\text{HS},\infty}^2))$ immediately proves (12). □

5. Certified 140-Digit Benchmarks and Independent Verification

All 25 benchmarks in Tables 1–3 were certified by `certify_finite_temp_polyakov.py` in `mpmath.iv` at 140-digit precision (`mp.mp.dps = 140`, `mp.iv.dps = 140`, zero black-box special functions, zero quadrature), outward-rounded directly via `mp.floor`/`mp.ceil` at scale 10^{12}. Lower bounds ($\geq$) are conservatively floored and upper bounds ($\leq$) ceiled at 10^{-6}.

Independent Verification and Lean 4.36.0 Benchmark Kernel. Our repository provides a four-part test suite (`test_finite_temp_polyakov.py`, 4/4 passing) and a standalone 94-line Lean 4.36.0 kernel (`FiniteTempPolyakovCerts.lean`):

- *Direct* 64×64 *Gauss–Legendre* SU(2) *ladder integration:* Confirms the exact character expansion (4) at $N_\tau = 2$ to 10^{-12} across $\beta_t \in \{0.8, 2.0\}$ and $\theta \in \{0, 0.9, \pi\}$.
- *Direct gradient and TV-oscillation tests:* Verifies $\|\nabla \operatorname{Re} L(\mathbf{x})\|_{\text{HS}}^2 \leq N_\tau / N$ on SU(2) and SU(3) ($N_\tau \in \{2, 4, 6\}$) by central finite differences, and verifies sharpness of Lemma 3.1 to 10^{-13}.
- *Independent* 80*-digit cross-check and parity:* All 25 rows in Tables 1–3 are independently recomputed at 80-digit precision and matched verbatim across JSON, Lean 4.36.0, and LaTeX.

TABLE 1. Certified 140-digit $(1+1)$D electric screening mass $m_{\mathrm{el}}^{\mathrm{2D}}$ (5) (conservatively floored at 10^{-6}), Dobrushin contraction α_{Dob}, rigorous confinement mass lower bound m_{conf} (8), and strong-coupling interaction pressure sandwich (10) for SU(2) at $\beta_s = 0$.

d	N_τ	β_t	$m_{\mathrm{el}}^{\mathrm{2D}}$	$\alpha_{\mathrm{Dob}} \le$	$m_{\mathrm{conf}} \ge$	$\Delta p_{d,L} \in$
3	2	1/2	4.179502	0.248712	1.391462	$[-0.064195, 0.060322]$
3	4	3/2	4.266780	0.226084	1.486851	$[-0.029091, 0.027491]$
3	4	2	3.346893	0.574682	0.553938	$[-0.077550, 0.067122]$
3	8	4	3.347828	0.569314	0.563324	$[-0.038384, 0.033268]$
4	2	1/2	4.179502	0.373068	0.985997	$[-0.096293, 0.090483]$
4	4	3/2	4.266780	0.339126	1.081386	$[-0.043636, 0.041236]$
4	4	2	3.346893	0.862023	0.148473	$[-0.116325, 0.100683]$
4	6	3	3.394609	0.816373	0.202884	$[-0.073128, 0.063784]$
4	8	4	3.347828	0.853970	0.157859	$[-0.057576, 0.049901]$

TABLE 2. Certified 140-digit SU(2) Dobrushin confinement threshold brackets $[\beta_t^{\mathrm{lo}}, \beta_t^{\mathrm{hi}}]$ at $\beta_s = 0$ (Theorem 3.2): because $\beta_t \mapsto I_{k+1}(\beta_t)/I_1(\beta_t)$ is strictly increasing on $(0,\infty)$ (by the monotonicity of modified Bessel ratios $\beta_t \mapsto I_{\nu+1}(\beta_t)/I_\nu(\beta_t)$), $\alpha_{\mathrm{Dob}}(d, \beta_t, N_\tau)$ is strictly increasing in β_t, so every $\beta_t \le \beta_t^{\mathrm{lo}}$ is provably confined with a unique center-symmetric Gibbs measure ($\alpha_{\mathrm{Dob}}(\beta_t^{\mathrm{lo}}) < 1 < \alpha_{\mathrm{Dob}}(\beta_t^{\mathrm{hi}})$).

d	N_τ	β_t^{lo} (confined)	β_t^{hi} (witness)	$\alpha_{\mathrm{Dob}}(\beta_t^{\mathrm{lo}}) \le$	$\alpha_{\mathrm{Dob}}(\beta_t^{\mathrm{hi}}) \ge$
3	2	1.01035	1.01036	0.999987	1.000005
3	4	2.41265	2.41266	0.999992	1.000003
3	6	3.60717	3.60718	0.999995	1.000002
3	8	4.72831	4.72832	0.999995	1.000001
4	2	0.82214	0.82215	0.999987	1.000010
4	4	2.09996	2.09997	0.999989	1.000002
4	6	3.18131	3.18132	0.999995	1.000004
4	8	4.18147	4.18148	0.999997	1.000005

TABLE 3. Certified 140-digit full anisotropic 4D SU(N) Bakry–Émery curvature floor κ_{aniso}, spatial-volume-uniform $O(1/N^2)$ Polyakov-loop variance bound $N^2 \operatorname{Var}(\operatorname{Re} L)$, and Herbst concentration rate $c_P = \kappa_{\mathrm{aniso}}/(2N_\tau)$ valid for all $N \ge 2$ (Theorem 4.2).

N_τ	γ_t	γ_s	$\kappa_{\mathrm{aniso}} \ge$	$N^2 \operatorname{Var}(\operatorname{Re} L) \le$	$c_P \ge$
2	0	0	0.500000	4.000000	0.125000
4	1/100	1/100	0.260000	15.384616	0.032500
4	1/80	1/200	0.199999	20.000001	0.024999
4	1/60	1/60	0.099999	40.000001	0.012499
6	1/100	1/100	0.260000	23.076924	0.021666
6	1/60	0	0.099999	60.000001	0.008333
8	1/100	1/100	0.260000	30.769231	0.016250
8	1/50	1/50	0.019999	400.000001	0.001249



(a) Dobrushin confinement thresholds **(b) 4D Polyakov confinement mass floor**

FIGURE 1. Certified finite-temperature SU(2) confinement profiles at $\beta_s = 0$ (Theorem 3.2). (a) Certified Dobrushin confinement thresholds $\beta_t^{\mathrm{Dob}}(d, N_\tau)$ growing nearly linearly with N_τ up to 4.18147 (4D) and 4.72831 (3D) at $N_\tau = 8$. (b) Certified Polyakov-loop confinement screening mass lower bound $m_{\mathrm{conf}} \geq \ln(1/\alpha_{\mathrm{Dob}})$ in 4D.

- *Zero-axiom Lean 4.36.0 benchmark kernel:* The file `FiniteTempPolyakovCerts.lean` compiles cleanly with `-DwarningAsError=true`, 0 warnings, 0 `sorry`, and `#print axioms` reporting *zero axioms* across all five theorems certifying the 25 integer-scaled (10^{12}) inequalities and coordination/Hessian combinatorics (the analytical proofs are given in Sections 2–4).

6. Scope and Limitations

(1) *Strong-spatial-coupling limit $\beta_s = 0$ in Theorems 2.1–4.1:* The exact Svetitsky–Yaffe reduction (3)–(4) and resulting high-β_t Dobrushin confinement thresholds hold at $\beta_s = 0$. Extending the Dobrushin confinement threshold to isotropic coupling $\beta_s = \beta_t$ requires controlling non-local spatial plaquette couplings induced after integrating out spatial links.

(2) *Deconfined phase and continuum critical temperature:* Theorem 3.2 proves confinement below $\beta_t^{\mathrm{Dob}}(d, N_\tau)$, while Theorem 4.2 controls the full anisotropic theory at $\beta_s \geq 0$ on the Bakry–Émery strong-coupling window $\max(24\gamma_t, 8\gamma_t + 16\gamma_s) < 1/2$. Rigorous determination of the 4D continuum deconfinement temperature $T_c/\sqrt{\sigma}$ remains open.

Data and Code Availability

All verification scripts (`certify_finite_temp_polyakov.py`, `generate_lean.py`), certified 140-digit data (`finite_temp_polyakov_certs.json`), independent tests (`test_finite_temp_polyakov.py`), and the standalone zero-axiom Lean 4.36.0 benchmark kernel (`FiniteTempPolyakovCerts.lean`) are archived online at:

https://airr.science/papers/ARR-2026-7GQC2S7HEJ8BEV94/

References

1. Dominique Bakry and Michel Émery, *Diffusions hypercontractives*, Séminaire de Probabilités XIX 1983/84, Lecture Notes in Mathematics, vol. 1123, Springer, 1985, pp. 177–206.
2. Dominique Bakry, Ivan Gentil, and Michel Ledoux, *Analysis and geometry of Markov diffusion operators*, Grundlehren der mathematischen Wissenschaften, vol. 348, Springer, 2014.
3. Christian Borgs and Erhard Seiler, *Lattice Yang–Mills theory at nonzero temperature and the confinement problem*, Communications in Mathematical Physics **91** (1983), no. 3, 329–380.

4. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Computer Science, vol. 12699, Springer, 2021, pp. 625–635.
5. Roland L. Dobrushin, *The description of a random field by means of conditional probabilities and conditions of its regularity*, Theory of Probability and its Applications **13** (1968), no. 2, 197–224.
6. Hans-Otto Georgii, *Gibbs measures and phase transitions*, 2nd ed., De Gruyter Studies in Mathematics, vol. 9, De Gruyter, 2011.
7. Jens Langelage and Owe Philipsen, *The deconfinement transition of finite density QCD with heavy quarks from strong coupling series*, Journal of High Energy Physics **2010** (2010), no. 1, 089.
8. Alexander M. Polyakov, *Thermal properties of gauge fields and quark liberation*, Physics Letters B **72** (1978), no. 4, 477–480.
9. Barry Simon, *The statistical mechanics of lattice gases*, vol. 1, Princeton University Press, 1993.
10. Leonard Susskind, *Lattice models of quark confinement at high temperature*, Physical Review D **20** (1979), no. 10, 2610–2618.
11. Benjamin Svetitsky and Laurence G. Yaffe, *Critical behavior at finite-temperature confinement transitions*, Nuclear Physics B **210** (1982), no. 4, 423–447.
12. E. T. Tomboulis and L. G. Yaffe, *Finite temperature $SU(2)$ lattice gauge theory*, Communications in Mathematical Physics **100** (1985), no. 3, 313–341.
13. Kenneth G. Wilson, *Confinement of quarks*, Physical Review D **10** (1974), no. 8, 2445–2459.

Independent Researcher, Östersund, Sweden
Email address: https://airr.science/papers/ARR-2026-7GQC2S7HEJ8BEV94/